



CLINIC-*SIM*

Simulating a Patient-Centered Approach to Optimize Early
Glycemic Control and Weight Management in Type 2 Diabetes

Disclaimer

- This presentation was commissioned by Lilly Medical and is intended to be used by HCPs for medical, scientific, and educational purposes.

Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making




SUPPORTING EARLY GLYCEMIC CONTROL AND WEIGHT MANAGEMENT THROUGH SHARED DECISION-MAKING

- Decision Point 1
- Long-Term Benefits of Early Glycemic Control
- Decision Point 2
- Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy
- Decision Point 3
- Supporting Weight Loss Early in Type 2 Diabetes Management

Supporting Early Glycemic Control for Patients With T2D

HCPs can support their patients **by assessing goals, advising, and escalating** to appropriate T2D pharmacotherapy when necessary

The **ADA-EASD Consensus Report** recommends several strategies for patients with T2D, including:



Assessing HbA_{1c} every 3 months for patients who are not at glycemic targets



Intensifying treatments when therapy goals are not met



In the absence of other comorbidities requiring more frequent monitoring, patients who have achieved glycemic targets can be assessed every 6 months



SUPPORTING EARLY GLYCEMIC CONTROL AND WEIGHT MANAGEMENT THROUGH SHARED DECISION-MAKING



Decision Point 1



Long-Term Benefits of Early Glycemic Control



Decision Point 2



Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy



Decision Point 3



Supporting Weight Loss Early in Type 2 Diabetes Management

ADA=American Diabetes Association; EASD=European Association for the Study of Diabetes; T2D=Type 2 Diabetes;
HCP=Health Care Professional; HbA_{1c}=Glycosylated hemoglobin
Davies MJ, et al. *Diabetes Care*. 2022;45(11):2753-2786.

Strategies to Foster Empathy and Trust with Patients: Co-Create Goals

Rationale: Using common language to discuss treatment goals results in greater clarity and impact




ASK

Your patients questions about their goals



PROBE

To clarify the questions



REPEAT

Words or expressions you heard back to the patients



RECORD

In chart for future exchanges



SUPPORTING EARLY GLYCEMIC CONTROL AND WEIGHT MANAGEMENT THROUGH SHARED DECISION-MAKING



Decision Point 1



Long-Term Benefits of Early Glycemic Control



Decision Point 2



Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy



Decision Point 3



Supporting Weight Loss Early in Type 2 Diabetes Management

5As of Weight Management

Obesity is a chronic, relapsing, and progressive disease, characterized by abnormal or excessive accumulation of adiposity that presents a risk to health^{1,2}.

How an individual develops obesity and how they will respond to interventions involves a complex interaction of genetic, biological, environmental, and behavioral factors³.

Successful obesity management requires identifying and addressing the 'root causes' of weight gain and barriers to weight management⁴.



ASK⁴
for permission to discuss weight and explore readiness



ASSESS⁴
obesity-related risks and potential "root causes" of weight gain



ADVISE⁴
on health risks and treatment options



AGREE⁴
on health outcomes and behavioral goals



ASSIST⁴
in accessing appropriate resources and providers and arrange follow-up



SUPPORTING EARLY GLYCEMIC CONTROL AND WEIGHT MANAGEMENT THROUGH SHARED DECISION-MAKING



Decision Point 1



Long-Term Benefits of Early Glycemic Control



Decision Point 2



Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy



Decision Point 3



Supporting Weight Loss Early in Type 2 Diabetes Management

1. https://www.who.int/health-topics/obesity#tab=tab_1 (Accessed August 28, 2023). 2. Sbraccia P. *Intern Emerg Med*. 2023;18:1-5.
3. <https://obesitycanada.ca/resources/tools-accessing-health-care/#> (Accessed August 28, 2023). 4. <https://obesitycanada.ca/wp-content/uploads/2025/03/Canadian-Adult-Obesity-CPG-Appendix-2-5As-Framework-for-Obesity-Management.pdf> (Accessed August 28, 2023)

Use of Appropriate Language with Patients^{1,2}

Appropriate use of language is a simple, implementable technique clinicians can use to lower the patient's anxiety, build confidence, and support self-care.²



Fosters patient-clinician collaboration

Ask open-ended questions that are not directly related to body weight

“What concerns you about your health today?”

[Click here to see an example](#)

Encourages engagement

When body weight is raised while addressing the patient's health concerns, it enhances patient engagement

Obesity is associated with several medical conditions, allowing for an easier transition to discuss body weight

Supports empathy and trust

Use non-medical language to discuss diabetes treatment goals to maintain patient comfort and engagement

E.g., how does the patient describe the impact of diabetes on their lives?



SUPPORTING EARLY GLYCEMIC CONTROL AND WEIGHT MANAGEMENT THROUGH SHARED DECISION-MAKING

Decision Point 1

Long-Term Benefits of Early Glycemic Control

Decision Point 2

Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

1. American Diabetes Association. *Diabetes Care*. 2022;45(1):S1-S264.

2. <https://www.england.nhs.uk/wp-content/uploads/2018/06/language-matters.pdf> (Accessed September 28, 2022).

Use of Appropriate Language with Patients^{1,2}

An example of patient-clinician communication



Clinician

What concerns do you have about your health today?



Patient

My knees have been bothering me lately and I've been having trouble sleeping...



Clinician

Thank you for letting me know about your knee pain and sleep issues. It's important to address these symptoms. They can sometimes be connected to underlying health conditions. Would it be okay if we discussed potential factors that might contribute to your symptoms?



Patient

Yes.



Clinician

It's worth considering that maintaining a healthy weight can often have a positive impact on knee pain and sleep quality. Many people have similar concerns, and they find that addressing them can improve their overall well-being. Would you be willing to talk about your weight? What are your thoughts about your weight right now?



Patient

Yes, when I was younger and my body weight was lower than it is now, I had more energy. Now I am too exhausted to get up and exercise.



Clinician

Would you be willing to work together and create a new treatment plan that feels manageable to you? You'll have less knee pain, better sleep, and more energy again.



Patient

Okay, I'm willing to try.



SUPPORTING EARLY GLYCEMIC CONTROL AND WEIGHT MANAGEMENT THROUGH SHARED DECISION-MAKING



Decision Point 1



Long-Term Benefits of Early Glycemic Control



Decision Point 2



Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy



Decision Point 3



Supporting Weight Loss Early in Type 2 Diabetes Management

Patient Profile at Visit #1 *(at time of diagnosis)*

Patient Profile

- 47-year-old African American female
- Diagnosed with T2D 9 months ago
- HbA_{1c} since diagnosis: 8.7%
- Current treatment: 1000 mg metformin daily at diagnosis
- Followed by registered dietitian/nurse educator

Medical History


- Prediabetes for 2 years preceding diagnosis, tried to manage with diet and exercise
- Gestational diabetes with her second child
- Hypertension
- Obesity
- Mixed dyslipidemia

Family History

- T2D in father, maternal aunt
- CAD in father
- Hypertension in mother and father

Social History

- Married, has 2 children, ages 10 and 15
- Never smoked or vaped, no illicit substances, rarely drinks alcohol
- High school math teacher



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

DECISION POINT 1

Long-Term Benefits of Early Glycemic Control


Decision Point 2

Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

Physical Exam and Labs



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

DECISION POINT 1

Long-Term Benefits of Early Glycemic Control

Decision Point 2

Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy

Decision Point 3


Supporting Weight Loss Early in Type 2 Diabetes Management

BMI=Body Mass Index; eGFR=Estimated Glomerular Filtration Rate; HDL=High-Density Lipoprotein; LDL=Low-Density Lipoprotein.

Visit #2 (9 months)

- The patient was diagnosed with type 2 diabetes, and her HbA_{1c} was 8.7% at the time. She was started on 1000 mg metformin daily. During that visit, she was advised on lifestyle modifications, including dietary changes to help both her diabetes and hypertension management, as well as recommendations to incorporate exercise. She was encouraged to start a statin for dyslipidemia but was too overwhelmed by her diabetes and declined statin therapy.

- At visit 2, her HbA_{1c} decreased to 7.9%. She again declined a statin and agreed to start an ACE inhibitor. Her metformin was titrated to 1500 mg total daily dose.



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

DECISION POINT 1

Long-Term Benefits of Early Glycemic Control

Decision Point 2

Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

ACE=Angiotensin-Converting Enzyme; HbA_{1c}=Glycosylated hemoglobin

Visit #3 - 4-month follow-up (13 months since diagnosis)

- She returns 4 months later.
- She is taking a dose of 1500 mg metformin daily (500 mg in the morning and 1000 mg in the evening) without missing doses. She was not able to tolerate a dose of 1000 mg twice daily. Her HbA_{1c} has decreased to 7.7% from 7.9%, and her body weight remained stable.
 - BMI: 33 kg/m² (195 lbs)
 - Blood pressure: 130/85 mmHg
 - Total cholesterol: 205 mg/dL
 - LDL cholesterol: 110 mg/dL
 - HDL cholesterol: 43 mg/dL
 - Triglycerides: 260 mg/dL

- She increased her physical activity by parking farther and trying to walk more but often struggles to find the time to get more exercise in.
- She tries to eat more balanced meals with protein, vegetables, and fewer carbohydrates; she finds it difficult to make multiple meals for her family members, who prefer to keep to their old dietary routine.
- She is motivated to continue working on lifestyle changes and has agreed to start a statin.



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

DECISION POINT 1

Long-Term Benefits of Early Glycemic Control

Decision Point 2

Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

Decision Point 1: In addition to continuing metformin 1500 mg daily, which of the following may be the next best step?

A

Continue lifestyle modifications – she is very motivated and prefers not to start another medication

B

Start a sulfonylurea and continue lifestyle modifications

C

Start an incretin receptor agonist and continue lifestyle modifications

D

Start a DPP-4 inhibitor



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

DECISION POINT 1

Long-Term Benefits of Early Glycemic Control

Decision Point 2

Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

Decision Point 1: In addition to continuing metformin 1500 mg daily, which of the following may be the next best step?

Incorrect -there may be a better option for this patient. Lifestyle modifications, including healthy nutrition, physical activity, and weight management, are important, and she has room for improvement. However, it is important to consider these changes take time, and biologic factors contributing to obesity and high HbA_{1c} may necessitate the addition of pharmacotherapy with lifestyle changes to achieve targets. To optimally reduce her potential risk of future diabetes complications, achieving glycemic control as early as possible is very important.

HbA_{1c}=Glycosylated hemoglobin.
Davies MJ, et al. *Diabetes Care*. 2022;45(11):2753-2786

Start an incretin receptor agonist and continue lifestyle modifications

Start a DPP-4 inhibitor



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

DECISION POINT 1

Long-Term Benefits of Early Glycemic Control

Decision Point 2

Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

DPP-4=Dipeptidyl Peptidase-4.

Decision Point 1: In addition to continuing metformin 1500 mg daily, which of the following may be the next best step?

Incorrect - there may be a better option for this patient. While sulfonylureas are effective for glucose control, the potential disadvantages of this therapy may outweigh the benefits for this patient. Specifically, sulfonylureas:

- Lack a durable effect
- Increase the risk of hypoglycemia
- Are associated with weight gain

The guidelines currently favor treatments with additional non-glucose-lowering effects, and an overall better safety profile. Given that this patient has obesity as well as taking into consideration current guideline recommendations for pharmacotherapy, an agent with additional benefits beyond glycemic control is likely to be more beneficial.

Davies MJ, et al. *Diabetes Care*. 2022;45(11):2753-2786

Davies MJ, et al. *Diabetes Care*. 2022;45[11]:2753-2786



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

DECISION POINT 1

Long-Term Benefits of Early Glycemic Control

Decision Point 2

Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

Decision Point 1: In addition to continuing metformin 1500 mg daily, which of the following may be the next best step?

Correct -this may be an appropriate option for this patient.

According to the ADA/EASD Consensus Report, incretin receptor agonists have high to very high efficacy for glucose lowering. Concomitant use of an incretin receptor agonist with an insulin secretagogue or insulin may increase the risk of hypoglycemia.

ADA=American Diabetes Association; EASD=European Association for the Study of Diabetes.
Davies MJ, et al. *Diabetes Care*. 2022;45(11):2753-2786.

C

Start an incretin receptor agonist and continue lifestyle modifications

D

Start a DPP-4 inhibitor



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

DECISION POINT 1

Long-Term Benefits of Early Glycemic Control

Decision Point 2

Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

Decision Point 1: In addition to continuing metformin 1500 mg daily, which of the following may be the next best step?

Incorrect - there may be a better option for this patient. According to the ADA/ EASD Consensus Report, DPP-4 inhibitors have modest glucose-lowering efficacy and a neutral effect on weight. Therefore, in this individual with uncontrolled type 2 diabetes and obesity, a DPP-4 inhibitor would not be the best choice.

ADA=American Diabetes Association; EASD=European Association for the Study of Diabetes; DPP-4=Dipeptidyl Peptidase-4.
Davies MJ, et al. *Diabetes Care*. 2022;45(11):2753-2786.

C

Start an incretin receptor agonist and continue lifestyle modifications

D

Start a DPP-4 inhibitor



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

DECISION POINT 1

Long-Term Benefits of Early Glycemic Control

Decision Point 2

Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

Long-Term Benefits of Early Glycemic Control

Learnings from UKPDS and Diabetes & Aging



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

Decision Point 1

LONG-TERM BENEFITS OF EARLY GLYCEMIC CONTROL

Decision Point 2

Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

The ADA-EASD Guidelines Recommend Early Intervention to Achieve Glycemic Targets

“Attaining recommended glycemic targets yields substantial and enduring reductions in the onset and progression of microvascular complications, and **early intervention** is essential”

“A reasonable HbA_{1c} to see microvascular benefits (generally 10 years) is around 53 mmol/mol (7%) or less. **Aiming for a lower HbA_{1c} level than this may have value if it can be achieved safely** without significant hypoglycemia or other adverse treatment effects.”



- Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making
- Decision Point 1
- LONG-TERM BENEFITS OF EARLY GLYCEMIC CONTROL**
- Decision Point 2
- Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy
- Decision Point 3
- Supporting Weight Loss Early in Type 2 Diabetes Management

ADA-EASD Guidelines Outline Medications for Glycemic Control and Weight Management as Adjuncts to Lifestyle Interventions^{1,2}

Goal | Achievement and Maintenance of Glycemic and Weight Management Goals

Glycemic Management: Choose approaches that provide the efficacy to achieve goals	Achievement and Maintenance of Weight Management Goals:
Efficacy for Glucose Lowering¹	Efficacy for Weight Loss¹
VERY HIGH - Dulaglutide (high dose)^a - Semaglutide^a - Tirzepatide^a - Insulin - Combination oral - Combination injectable (GLP-1 RA/insulin)	VERY HIGH - Semaglutide^a - Tirzepatide^a
HIGH - GLP-1 RA (not listed above) - Metformin^a - SGLT-2 inhibitor - SU - TZD	HIGH - Dulaglutide^a - Liraglutide^a
INTERMEDIATE DPP-4 inhibitor	INTERMEDIATE - GLP-1 RA (not listed above) - SGLT-2 inhibitor
	NEUTRAL - DPP-4 inhibitor - Metformin^a


Medications for the management of T2D can have varying effects on clinical outcomes, including body weight, that vary between, and sometimes within, drug classes²

^a For the prescribing information of these products, kindly refer to the [USPI References Slide](#) at the end.

DPP-4=Dipeptidyl Peptidase-4; GLP-1 RA=Glucagon-Like Peptide-1 Receptor Agonist; SGLT-2=Sodium-Glucose Cotransporter-2; SU=Sulfonylurea; TZD=Thiazolidinedione; ADA=American Diabetes Association; EASD=European Association for the Study of Diabetes; T2D=Type 2 Diabetes; USPI=United States Prescribing Information.


1. Davies MJ, et al. *Diabetes Care*. 2022;45(11):2753-2786. 2. Apovian CM, et al. *Adv Ther*. 2019;36:44-58.

VV-MED-148828 © 2025 Lilly USA, LLC. All rights reserved



Early Intensive Glycemic Control Is Associated With Decreased Risk of Type 2 Diabetes-Related Complications¹⁻³

Long-term Follow-up and Legacy Effect



In the UKPDS trial:

Historical HbA_{1c} values have a **greater impact** than recent values on all-cause mortality, myocardial infarction, and microvascular disease

Results from the 10-year post-trial follow-up showed:

13% risk reduction in all-cause mortality (p=0.007)

15% risk reduction in myocardial infarction (p=0.01)

24% risk reduction in microvascular disease (p=0.001)



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

Decision Point 1

LONG-TERM BENEFITS OF EARLY GLYCEMIC CONTROL

Decision Point 2

Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

UKPDS=The UK Prospective Diabetes Study; HbA_{1c}=Glycosylated hemoglobin..

1. UK Prospective Diabetes Study (UKPDS) Group. *Lancet*. 1998;352(9131):837-853. 2. Holman RR, et al. *N Engl J Med*. 2008;359(15):1577-1589.

3. Lind M, et al. *Diabetes Care*. 2021 Oct 1;44(10):2231-7

Improved Glycemic Control Lowers the Risk of Microvascular and Macrovascular Complications

DCCT/EDIC study^{1,2}

Among patients with T1D, initial intensive glycemic therapy and comprehensive care improved glycemia and overall outcomes during long-term follow-up



Although DCCT/EDIC was conducted in patients with T1D, the consequences of hyperglycemia are similar in T1D and T2D. Similar to UKPDS, this trial highlights **the importance of early glycemic control on long-term outcomes in diabetes.**



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

Decision Point 1

LONG-TERM BENEFITS OF EARLY GLYCEMIC CONTROL

Decision Point 2

Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

DCCT/EDIC=Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications; T1D=Type 1 Diabetes; T2D=Type 2 Diabetes.


1. Nathan DM, et al. *N Engl J Med.* 2005;353:2643-53. 2. DCCT/EDIC study research group. *Diabetes Care.* 2016;39:686-93.

Improved Glycemic Control Lowers the Risk of Microvascular and Macrovascular Complications



DCCT/EDIC study

To determine whether intensive therapy compared with conventional therapy during the Diabetes Control and Complications Trial affected the incidence of cardiovascular disease over 30 years of follow-up



Decision Point 1

LONG-TERM BENEFITS OF EARLY GLYCEMIC CONTROL

Decision Point 2

Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

DCCT=Diabetes Control and Complications Trial; EDIC=Epidemiology of Diabetes Interventions and Complications.

1. Nathan OM, et al. *N Engl J Med*. 2005;353:2643-53. 2. DCCT/EDIC study research group. *Diabetes Care*. 2016;39:686-93.

In DCCT/EDIC, Improved Glycemic Control Lowered the Risk of Microvascular and Macrovascular Complications



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

Decision Point 1

LONG-TERM BENEFITS OF EARLY GLYCEMIC CONTROL

Decision Point 2

Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management


During 30 years of follow-up:



149 CV events
in 82 former intensive treatment participants¹


217 CV events
in 102 former conventional treatment participants¹

Macrovascular risk reduction with intensive versus conventional treatment:




↓ **30%** risk reduction for the first CV event¹

↓ **32%** risk reduction for the first occurrence of MACE (nonfatal MI, stroke or CV death)¹



Intensive glycemic therapy is associated with microvascular benefit:

↑ prevalence of microalbuminuria in the conventional group at EDIC year 11²




CV=Cardiovascular; DCCT/EDIC=Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications; MACE=Major Adverse Cardiovascular Event; MI=Myocardial Infarction.

1. DCCT/EDIC study research group. *Diabetes Care*. 2016;39:686-93. 2. Nathan DM, et al. *N Engl J Med*. 2005;353:2643-53.

Achieving HbA_{1c} <6.5% in the First Year After Diagnosis Was Associated With a Lower Risk of T2D Complications

In the **Diabetes & Aging study**: Compared with HbA_{1c} levels <6.5%, patients with HbA_{1c} ≥6.5% in the first year had a higher risk for microvascular and macrovascular events




- Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making
- Decision Point 1
- LONG-TERM BENEFITS OF EARLY GLYCEMIC CONTROL**
- Decision Point 2
- Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy
- Decision Point 3
- Supporting Weight Loss Early in Type 2 Diabetes Management

T2D=Type 2 Diabetes; HbA_{1c}=Glycosylated hemoglobin.
Laiteerapong N, et al. *Diabetes Care*. 2019;42(3):416-426.



Achieving HbA_{1c} <6.5% in the First Year After Diagnosis Was Associated With a Lower Risk of T2D Complications

In the Diabetes & Aging study: Compared with HbA levels <6.5%, patients with HbA




- Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making
- Decision Point 1
- LONG-TERM BENEFITS OF EARLY GLYCEMIC CONTROL**
- Decision Point 2
- Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy
- Decision Point 3
- Supporting Weight Loss Early in Type 2 Diabetes Management

T2D=Type 2 Diabetes.
Laiteerapong N, et al. *Diabetes Care*. 2019;42[3]:416-426..



Visit #4 – 3-month follow-up (16 months after diagnosis)



- Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making
- Decision Point 1
- long-term benefits of early glycemic control
- DECISION POINT 2**
- Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy
- Decision Point 3
- Supporting Weight Loss Early in Type 2 Diabetes Management

BMI=Body Mass Index; HbA_{1c}=Glycosylated hemoglobin.


Decision Point 2: Which of the following may be the next best step?

A

She is again reluctant to add another medication. Continue metformin and lifestyle modifications with additional counseling

B

Review the potential benefits of early glycemic control and potential additional non-glycemic benefits again at this visit and suggest adding an incretin receptor agonist to her current regimen of 1500 mg metformin daily



Decision Point 2: Which of the following may be the next best step?

Incorrect -there may be a better option for this patient. While lifestyle modifications are a cornerstone to optimizing blood glucose and weight management in people with type 2 diabetes, many metabolic and hormonal processes make achieving target glycemic and weight control with lifestyle alone difficult and sometimes impossible without pharmacotherapy.

Evidence suggests the patient may likely experience disease progression on metformin monotherapy. In addition to missing the potential key benefits of early effective glycemic control, metformin alone may not provide the added weight loss benefit that may be achieved with one of the other agents.

Kahn SE, et al. *N Engl J Med*. 2006 Dec 7;355[23]:2427-43.

metformin daily



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

Decision Point 1

long-term benefits of early glycemic control

DECISION POINT 2

Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

Decision Point 2: Which of the following may be the next best step?

Correct - this may be an appropriate option for this patient.

According to the ADA/EASD Consensus Report, glucose-lowering therapies with high to very high dual glucose and weight efficacy are preferred for patients with type 2 diabetes, given the importance of early and sustained glycemic control.

ADA=American Diabetes Association; EASD=European Association for the Study of Diabetes.
Davies MJ, et al. *Diabetes Care*. 2022;45(11):2753-2786.

metformin and lifestyle
modifications with additional
counseling

glycemic benefits again at this
visit and suggest adding an
incretin receptor agonist to her
current regimen of 1500 mg
metformin daily



- Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making
- Decision Point 1
- long-term benefits of early glycemic control
- DECISION POINT 2**
- Optimizing Early Glycemic Control Using Combination Therapy or Very High-Efficacy Monotherapy
- Decision Point 3
- Supporting Weight Loss Early in Type 2 Diabetes Management

Optimizing Early Glycemic Control Using Combination Therapy or Very High- Efficacy Monotherapy



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

Decision Point 1

Long-Term Benefits of Early Glycemic Control

Decision Point 2

OPTIMIZING EARLY GLYCEMIC CONTROL USING COMBINATION THERAPY OR VERY HIGH-EFFICACY MONOTHERAPY

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

Type 2 Diabetes Progression on Monotherapy

Learnings from UKPDS¹⁴

Among **newly diagnosed** patients with type 2 diabetes receiving **diet, metformin, sulfonylurea or insulin**:



HbA_{1c} progressively increases over time on monotherapy despite an initial decrease in the first year^{1,2}




Progressive loss of β -cell function, regardless of which therapy was used^{3,4}

After 9 years of monotherapy, only 9% (diet), 24% (sulfonylurea), and 28% (insulin) achieved HbA_{1c} <7%⁵

Learnings from ADOPT⁶

Among newly diagnosed patients with type 2 diabetes receiving **rosiglitazone, metformin^a, or glyburide^a** as initial treatment:

Cumulative incidence of monotherapy failure using historic therapies at 5 years:



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

Decision Point 1

Long-Term Benefits of Early Glycemic Control

Decision Point 2

OPTIMIZING EARLY GLYCEMIC CONTROL USING COMBINATION THERAPY OR VERY HIGH-EFFICACY MONOTHERAPY

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

*p<0.001.

^aFor the prescribing information of these products, kindly refer to the [USPI References Slide](#) at the end.

ADOPT=A Diabetes Outcome Progression Trial; UKPDS=The UK Prospective Diabetes Study; HbA_{1c}= Glycosylated hemoglobin; CI=Confidence Interval.

1. DeFronzo RA. *Diabetes*. 2009;58(4):773-95. 2. UK Prospective Diabetes Study (UKPDS) Group. *The Lancet*. 1998;352(9131):837-53. 3. UK Prospective Diabetes Study Group. *Diabetes*. 1995;44(11):1249-58. 4. Matthews DR, et al. *Diabetic medicine*. 1998;15(4):297-303. 5. Turner RC, et al. *JAMA*. 1999;281(21):2005-12. 6. Kahn SE, et al. *N Engl J Med*. 2006;355(23):2427-43.


Type 2 Diabetes Progression on Monotherapy

Learnings from UKPDS¹⁴

Learnings from ADOPT⁶

UKPDS

Effect of intensive vs. conventional glucose control on risk of micro and macrovascular complications



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

Decision Point 1

Long-Term Benefits of Early Glycemic Control

Decision Point 2

OPTIMIZING EARLY GLYCEMIC CONTROL USING COMBINATION THERAPY OR VERY HIGH-EFFICACY MONOTHERAPY

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

^a The median follow-up for the sulfonylurea-insulin and metformin groups was 10.0 years and 10.7 years, respectively.

FPG=Fasting Plasma Glucose; SU=Sulfonylureas; T2D=Type 2 Diabetes; UKPDS=The UK Prospective Diabetes Study.

1. UK Prospective Diabetes Study (UKPDS) Group. *Lancet*. 1998;352(9131):837-853. 2. Holman RR, et al. *N Engl J Med*. 2008;359 (15):1577-1589.


Type 2 Diabetes Progression on Monotherapy

Learnings from UKPDS¹⁴

Learnings from ADOPT⁶

ADOPT

To compare the effectiveness of rosiglitazone (TZD), as compared with glyburide (SU) and metformin, in maintaining long-term glycemic control in T2D.



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

Decision Point 1

Long-Term Benefits of Early Glycemic Control

Decision Point 2

OPTIMIZING EARLY GLYCEMIC CONTROL USING COMBINATION THERAPY OR VERY HIGH-EFFICACY MONOTHERAPY

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

Note: Patients who withdraw from treatment for any reason will continue to be followed in nontreatment observational follow up.

^aFor the prescribing information of these products, kindly refer to the [USPI References Slide](#) at the end.

ADOPT =A Diabetes Outcome Progression Trial; HbA1 c=Glycated Hemoglobin; SU=Sulfonylurea; T2D= Type 2 Diabetes; TZD= Thiazolidinedione.

Kahn SE, et al. *N Engl J Med*. 2006;355(23):2427-43.

Early Combination Treatment Reduces Time to Treatment Failures^{1,2}


VERIFY Trial Results

Number of patients with good glycemic control over 5 years

Time before treatment intensification is needed

Initial treatment failure

Secondary treatment failure



>2x

More than twice as many patients had an **extended time with good glycemic control** following early combination therapy

26%

reduction in the risk of **time to secondary treatment failure** following early combination therapy

Adapted from Matthews et al.²

These results indicate that long-term clinical benefits may be achieved more frequently, and potentially without additional tolerability issues, with the **early initiation of medication(s) that target multiple pathways** compared with traditional stepwise therapy



- Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making
- Decision Point 1
- Long-Term Benefits of Early Glycemic Control
- Decision Point 2
- OPTIMIZING EARLY GLYCEMIC CONTROL USING COMBINATION THERAPY OR VERY HIGH-EFFICACY MONOTHERAPY**
- Decision Point 3
- Supporting Weight Loss Early in Type 2 Diabetes Management

VERIFY=Vildagliptin Efficacy in Combination With Metformin For Early Treatment of Type 2 Diabetes.

1. Del Prato S, et al. *Diabet Med*. 2014;31(10):1178-1184. 2. Matthews D, et al. *Diabetes Ther*. 2020;11(11):2465-2476

Early Combination Treatment Reduces Time to Treatment Failures^{1,2}

VERIFY Trial Results

Number of patients with good glycemic

Time before treatment


Initial treatment

Secondary treatment

VERIFY

To determine whether initiation of combination therapy would result in more durable glycemic control compared to metformin monotherapy in treatment-naïve patients with T2D

If HbA_{1c} was maintained at <7%, then treatment did not change



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

Decision Point 1

Long-Term Benefits of Early Glycemic Control

Decision Point 2

OPTIMIZING EARLY GLYCEMIC CONTROL USING COMBINATION THERAPY OR VERY HIGH-EFFICACY MONOTHERAPY

Decision Point 3

Supporting Weight Loss Early in Type 2 Diabetes Management

^aFor the prescribing information of these products, kindly refer to the [USPI References Slide](#) at the end.

DPP4=Diipeptidyl Peptidase-4 Inhibitor; HbA_{1c}=Glycated Hemoglobin; T2D=Type 2 Diabetes; VERIFY=Vildagliptin Efficacy in Combination With Metformin For Early Treatment of Type 2 Diabetes.

Matthews D et al, *Diabetes Ther.* 2020;11:2465-2476.

ADA-EASD Guidelines Recommend Incorporating Glucose-Lowering Therapies With Complementary Mechanisms of Action

Traditionally, a stepwise approach was advocated, in which a new agent is added to the existing regimen, but **evidence is growing to support a more proactive approach** of combining glucose-lowering agents from the time of initial diagnosis

When targets are not met, the intensification of glucose-lowering medication by combining **agents with complementary mechanisms of action** should be considered



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

Decision Point 1

Long-Term Benefits of Early Glycemic Control

Decision Point 2

OPTIMIZING EARLY GLYCEMIC CONTROL USING COMBINATION THERAPY OR VERY HIGH-EFFICACY MONOTHERAPY

Decision Point 3


Supporting Weight Loss Early in Type 2 Diabetes Management

Available Medications Can Treat Multiple Metabolic Mechanisms That Drive Type 2 Diabetes Progression¹⁻³

Class ^{2,3}	Primary Physiological Action(s) ^{2,3}	Core Defects ¹⁻³
Biguanides	↓ Hepatic glucose production	 Liver
SUs	↑ Insulin secretion	 Pancreatic β-cell
TZDs	↑ Insulin sensitivity	 Whole body
DPP-4 inhibitors	↑ Insulin secretion (glucose-dependent) ↓ Glucagon secretion (glucose-dependent)	 Pancreatic α-cell  Pancreatic β-cell
SGLT-2 inhibitors	Blocks glucose reabsorption by the kidney, increasing glucosuria 	 Kidney
Incretin receptor agonists*	↑ Insulin secretion (glucose-dependent) ↓ Glucagon secretion (glucose-dependent) ↓ Delays gastric emptying ↓ Weight	 Pancreatic α-cell  Pancreatic β-cell  GI tract


*Only actions consistent in both classes of incretin receptor agonists shown; the GIP/GLP-1 RA class acts on additional defects. Schematic is intended to provide an overview of T2D drugs and is not specific to only one product within each class listed. It is not limited to make any express or implied comparison among products. Classes shown are from the ADA guidelines and do not represent all T2D classes available to treat hyperglycemia. DPP-4=Dipeptidyl Peptidase-4; GIP=Glucose-Dependent Insulinotropic Polypeptide; GLP-1RA=Glucagon-Like Peptide-1 Receptor Agonist; SGLT2=Sodium-Glucose Cotransporter-2; SU=Sulfonylurea; TZD=Thiazolidinedione.

1. DeFronzo RA. *Diabetes*. 2009;58(4):773-795. 2. Inzucchi SE, et al. *Diabetes Care*. 2015;38:140-149. 3. Mounjaro (tirzepatide once weekly) [Summary of Product Characteristics]. Houten, Utrecht, Netherlands: Eli Lilly and Company



Visit 5 - She returns in 4 months (20 months after diagnosis)


- She is very pleased with her progress in lifestyle modifications, including improving her diet and exercising.
- Her HbA_{1c} remains at 7.4% and she is very frustrated by the lack of weight change despite lifestyle changes.
- She is still somewhat reluctant to try additional therapy, though.



- Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making
- Decision Point 1
- Long-Term Benefits of Early Glycemic Control
- Decision Point 2
- Optimizing early glycemic control using combination therapy or very high-efficacy monotherapy
- DECISION POINT 3**
- Supporting Weight Loss Early in Type 2 Diabetes Management

HbA_{1c}=Glycosylated hemoglobin.

Decision Point 3: In addition to continuing metformin 1500 mg daily, which of the following may be the next best step?



SGLT-2=Sodium-Glucose Cotransporter-2.



- Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making
- Decision Point 1
- Long-Term Benefits of Early Glycemic Control
- Decision Point 2
- Optimizing early glycemic control using combination therapy or very high-efficacy monotherapy
- DECISION POINT 3**
- Supporting Weight Loss Early in Type 2 Diabetes Management

Decision Point 3: In addition to continuing metformin 1500 mg daily, which of the following may be the next best step?

Incorrect - there may be a better option for this patient. Starting an SGLT-2 inhibitor is a reasonable consideration, especially given that she has room for improvement in her glycemic control. However, according to the ADA/EASD Consensus Report, a regimen with high to very high dual glucose and weight efficacy is recommended in individuals needing to achieve glycemic and weight management goals.

ADA=American Diabetes Association; EASD=European Association for the Study of Diabetes; SGLT-2=Sodium-Glucose Cotransporter-2.
Davies MJ, et al. *Diabetes Care*. 2022;45(11):2753-2786.

Start an incretin receptor agonist with high glycemic efficacy and weight reduction



- Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making
- Decision Point 1
- Long-Term Benefits of Early Glycemic Control
- Decision Point 2
- Optimizing early glycemic control using combination therapy or very high-efficacy monotherapy
- DECISION POINT 3**
- Supporting Weight Loss Early in Type 2 Diabetes Management

Decision Point 3: In addition to continuing metformin 1500 mg daily, which of the following may be the next best step?

Incorrect - there may be a better option for this patient. Lifestyle counseling and modifications, including physical activity, are critically important. However, considering that biologic contributors of obesity and high HbA_{1c} are often difficult to overcome with lifestyle modifications alone, it is important to remember that pharmacotherapy is often also needed to help get patients to glycemic and weight targets. Early weight loss is associated with delayed disease progression, and incorporating a glucose-lowering therapy with high efficacy for weight loss may improve long-term outcomes.

HbA_{1c}= Glycosylated hemoglobin.
Davies MJ, et al. *Diabetes Care*. 2022;45(11):2753-2786.

Start an incretin receptor agonist with high glycemic efficacy and weight reduction



- Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making
- Decision Point 1
- Long-Term Benefits of Early Glycemic Control
- Decision Point 2
- Optimizing early glycemic control using combination therapy or very high-efficacy monotherapy
- DECISION POINT 3**
- Supporting Weight Loss Early in Type 2 Diabetes Management

Decision Point 3: In addition to continuing metformin 1500 mg daily, which of the following may be the next best step?

Correct - this may be an appropriate option for this patient.

According to the ADA/EASD Consensus Report, incretin receptor agonists have high to very high efficacy for glycemic control and intermediate to very high efficacy for weight reduction.

ADA=American Diabetes Association; EASD=European Association for the Study of Diabetes.
Davies MJ, et al. *Diabetes Care*. 2022;45(11):2753-2786.

Start an incretin receptor agonist with high glycemic efficacy and weight reduction



- Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making
- Decision Point 1
- Long-Term Benefits of Early Glycemic Control
- Decision Point 2
- Optimizing early glycemic control using combination therapy or very high-efficacy monotherapy
- DECISION POINT 3**
- Supporting Weight Loss Early in Type 2 Diabetes Management

Supporting Weight Loss Early in Type 2 Diabetes Management

Learnings from UKPDS, DiRECT, and Counterbalance



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

Decision Point 1

Long-Term Benefits of Early Glycemic Control

Decision Point 2


optimizing early glycemic control using combination therapy or very high-efficacy monotherapy

Decision Point 3

SUPPORTING WEIGHT LOSS EARLY IN TYPE 2 DIABETES MANAGEMENT

ADA-EASD Guidelines on Achievement of Weight Goals for Type 2 Diabetes Management^{1,2}

“**Weight loss of 5%-10%** confers metabolic improvement; weight loss of **10%-15%** or more may have a disease-modifying effect²”



“Weight loss may exert benefits that extend **beyond glycemic management** to improve risk factors for cardiometabolic disease²”



- Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making
- Decision Point 1
- Long-Term Benefits of Early Glycemic Control
- Decision Point 2
- optimizing early glycemic control using combination therapy or very high-efficacy monotherapy
- Decision Point 3
- SUPPORTING WEIGHT LOSS EARLY IN TYPE 2 DIABETES MANAGEMENT**

1. American Diabetes Association. *Diabetes Care*. 2022;45(suppl 1):S83-S96. 2. Davies MJ, et al. *Diabetes Care*. 2022;45(11):2753-2786.

Excess Weight and Glycemic Control



UKPDS

UKPDS demonstrated that fasting plasma glucose and HbA_{1c} increase progressively to a greater extent in participants with obesity compared to those without obesity.¹




DiRECT

The DiRECT trial demonstrated that early weight reduction is associated with improved HbA_{1c} levels and participants achieving ≥15 kg weight loss were more likely to achieve HbA_{1c} <6.5%.²



COUNTER BALANCE


The Counterbalance study is a small study that demonstrated improvement in glycemic control following weight loss was greater in those with shorter duration of T2D than in those with longer duration of T2D.³



Counterbalance=Counteracting Beta Cell Failure By Long Term Action to Normalize Calorie Intake; DiRECT=Diabetes Remission Clinical Trial; UKPDS=The UK Prospective Diabetes Study; HbA_{1c}=Glycosylated hemoglobin; T2D=Type 2 Diabetes.

1. United Kingdom Prospective Diabetes Study Group. *BMJ: British Medical Journal*. 1995;83-8. 2. Lean MEJ, et al. *Lancet Diabetes Endocrinol*. 2019;7(5):344-355. 3. Steven S et al. *Diabetes Care*. 2016;39:808-815.

Excess Weight and Glycemic Control




- Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making
- Decision Point 1
- Long-Term Benefits of Early Glycemic Control
- Decision Point 2
- optimizing early glycemic control using combination therapy or very high-efficacy monotherapy
- Decision Point 3
- SUPPORTING WEIGHT LOSS EARLY IN TYPE 2 DIABETES MANAGEMENT**

^aThe median follow-up for the sulfonylurea-insulin and metformin groups was 10.0 years and 10.7 years, respectively. FPG=Fasting Plasma Glucose; SU=Sulfonylureas; T2D=Type 2 Diabetes; UKPDS=The UK Prospective Diabetes Study.

1. UK Prospective Diabetes Study (UKPDS) Group. *Lancet*. 1998;352(9131):837-853. 2. Holman RR, et al. *N Engl J Med*. 2008;359(15):1577-1589

Excess Weight and Glycemic Control



- Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making
- Decision Point 1
- Long-Term Benefits of Early Glycemic Control
- Decision Point 2
- optimizing early glycemic control using combination therapy or very high-efficacy monotherapy
- Decision Point 3
- SUPPORTING WEIGHT LOSS EARLY IN TYPE 2 DIABETES MANAGEMENT**

BMI=Body Mass Index; DiRECT=Diabetes Remission Clinical Trial; T2D=Type 2 Diabetes; VLCD=Very Low-Calorie Diet.

Lean MEJ et al. *Lancet Diabetes Endocrinol.* 2019;7[5]:344-355.

Excess Weight and Glycemic Control

COUNTER BALANCE

To study the effect of weight loss, achieved through a VLCD, on control of fasting plasma glucose levels in T2D



N=30
With T2D of either short duration (<4 years) or long duration (>8 years) and BMI 27-45 kg/m²

Very-low-calorie diet (VLCD)

Isocaloric diet

Structured, individualized weight maintenance program

8 weeks

2 weeks

6 months Follow-up



Supporting Early Glycemic Control and Weight Management Through Shared Decision-Making

Decision Point 1

Long-Term Benefits of Early Glycemic Control

Decision Point 2

optimizing early glycemic control using combination therapy or very high-efficacy monotherapy

Decision Point 3

SUPPORTING WEIGHT LOSS EARLY IN TYPE 2 DIABETES MANAGEMENT

BMI=Body Mass Index; Counterbalance=Counteracting Beta Cell Failure By Long Term Action to Normalize Calorie Intake; T2D= Type 2 Diabetes; VLCD=Very Low-Calorie Diet.
Steven Set al. *Diabetes Care*. 2016;39:808-815.

USPI References^a



- **Dulaglutide (TRULICITY®)**
[Trulicity® Prescribing Information](#)
- **Glyburide (DiaBeta®)**
[DiaBeta® Prescribing Information](#)
- **Liraglutide (SAXENDA®)**
[Saxenda® Prescribing Information](#)
- **Metformin (GLUMETZA®)**
[Glumetza® Prescribing Information](#)
- **Semaglutide (OZEMPIC®)**
[Ozempic® Prescribing Information](#)
- **Tirzepatide (MOUNJARO®)**
[Mounjaro® Prescribing Information](#)
- **Rosiglitazone (AVANDIA)**
[Avandia Prescribing Information](#)

^aOther product/company names mentioned herein are the trademarks of their respective owners.



CLINIC-*SIM*

Simulating a Patient-Centered Approach to Optimize Early
Glycemic Control and Weight Management in Type 2 Diabetes