TAMPER EVIDENT: DO NOT USE IF POUCH IS OPEN

Drug Facts

Active Ingredient (in each softgel) Purpose Bismuth subsalicylate 262 mg......Upset stomach reliever and anti-diarrheal

Uses relieves • travelers' diarrhea • diarrhea • upset stomach due to overindulgence in food and drink, including: • heartburn • indigestion • nausea • gas • belching • fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert Contains salicylate.

Do not take if you are

allergic to salicylates (including aspirin) • taking other salicylate products
 Do not use if you have • an ulcer • a bleeding problem • bloody or

black stool

Ask a doctor before use if you have • fever • mucus in the stool

• rever • mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for • anticoagulation (thinning the blood) • diabetes • gout • arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

 symptoms get worse or last more than 2 days

 ringing in the ears or loss of hearing occurs
 diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions • swallow with water, do not chew • adults and children 12 years and over: • 2 softgels (1 dose) every 1/2 hour or 4 softgels (2 doses) every hour as needed for diarrhea • 2 soft gels (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, hearthurn.indigestion, nausea) • do not exceed 8 doses (16 softgels) in 24 hours • use until diarrhea stops but not more than 2 days • children under 12 years: ask a doctor • drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other Information

 each softgel contains: Salicylate 99 mg • very low sodium • store between 15–30°C (59–86°F)

Inactive ingredients Aerosil, beeswax, FD&C Red No. 33, FD&C Vellow No. 6, gelatin, glycerin, light liquid paraffin, purified water, soya lecithin, sodium carboxy methyl cellulose, sorbitol solution, titanium dioxide

Questions or comments? 1-888-333-9792

* This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark Pepto-Bismol[®]



Less than the limit of detection and consistent with gluten-free diet abeling per FDA

NDC: 68210-4170-1



ACC-1-10020 • Distributed by Cabinet Health P.B.C. Question/Comments? 1-888-333-9792 • www.cabinethealth.con

PACKAGE NOT CHILD-RESISTANT

CABINET:

 \gg

Stomach Relief

Compare to the active ingredient in: Pepto-Bismol^{®*}

Active ingredient: BISMUTH SUBSALICYLATE 262 MG

Relieves: Nausea Heartburn Indigestion Upset Stomach Diarrhea

COMPOSTABLE REFILL POUCH



