

Home Care - Physical Therapy (PT) and Occupational Therapy (OT)

Disclaimer

Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.

Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.

Summary

Members recently discharged from the hospital and/or those diagnosed with certain medical conditions may require short-term skilled care in the home for rehabilitation. When medically necessary, such services can be used to restore or improve functional independence. Physical therapy (PT) and occupational therapy (OT) are examples of these skilled home care services.

PT is designed to improve functioning, relieve disease symptoms, and prevent disability in individuals with acute and chronic disease. Treatments may consist of heat and cold therapy, electric stimulation, a variety of exercise regimens, and functional training for ambulatory activities. PT may be performed by a qualified, licensed physical therapist or by a physical therapy assistant (PTA) under the supervision of a qualified, licensed physical therapist.

OT is designed to provide individuals with purposeful activities and training to regain skills of daily living that were lost through disease or injury. OT may be performed by a qualified, licensed occupational therapist or by an occupational therapy assistant (OTA) under the supervision of a qualified, licensed occupational therapist.

PT and OT are often coordinated by a multidisciplinary team of licensed therapists, nurses, and prescribing clinicians. Home PT and OT require a prescription and clear documentation of progress,

goals, and ongoing necessity. Home PT and OT should also include a home exercise and activity program designed for the member to participate in alone or with the help of caregivers that do not require skilled personnel present. This guideline provides criteria regarding the indications and exclusions for PT and OT.

Information about coverage and benefit limitations can be found in the member's plan contract at hioscar.com/forms.

Definitions

"Homebound" refers to members who have normal inability to leave home without considerable and taxing effort (i.e., requires an assistive device or the assistance of another person to leave home) AND one of the following:

- Members who cannot leave home due to a medical condition, chronic disease, or injury; *or*
- Members advised by a treating provider not to leave home for various reasons (e.g. safety, ongoing medical treatment needs, etc); *or*
- Members who need the aid of supportive devices such as crutches, canes, wheelchairs, and walkers, special transportation (when the member is unable to use common transportation such as private automobile, bus, taxi due to medical condition), or the assistance of others to leave their place of residence.

When the member does leave home, the absence of the member from the home is infrequent or for short periods of time, such as to receive health treatment or adult daycare (non-residential program providing services during the day).

"Physical Therapy (PT)" refers to supervised therapeutic procedures performed by licensed healthcare professionals which are intended to relieve disease symptoms, prevent disability and restore clinical function. PT is often one of many components in a multidisciplinary treatment plan following injury or in chronic disease. Physical therapy may include, but is not limited to:

- Ambulation and mobility training
- Gait and balance training
- Strength training
- Joint mobilization
- Neuromuscular reeducation
- Therapeutic exercises
- Assistive device and adaptive equipment training
- Orthotic or prosthetic training
- Transfer training

"Occupational Therapy" refers to a therapeutic intervention program designed and supervised by a team of physicians and occupational therapists to assist members in regaining skills of daily living that

have been lost or impaired. Such programs are individualized to each member to help improve quality of life by restoring independence. Occupational therapy may include, but is not limited to:

- Activities of Daily Living (ADL) training
- Muscle re-education
- Cognitive or neurodevelopmental training
- Perceptual motor training
- Fine motor coordination/strength training
- Assistive device and adaptive equipment training
- Environment modification recommendations and training
- Transfer training
- Functional mobility training
- Manual therapy

“Activities of Daily Living (ADLs)” are defined as routine activities that most healthy persons perform daily without requiring assistance. These include, but are not limited to: bathing, communication, dressing, feeding, grooming, mobility, personal hygiene, self-maintenance, skin management, and toileting.

“Instrumental Activities of Daily Living (IADLs)” are defined as activities that may be performed daily but are not fundamental for daily functioning. These include, but are not limited to: the use of public transportation, balancing a checkbook, community living activities, meal preparation, laundry, leisure activities and sports, and motor vehicle operation.

“Rehabilitative Treatments” are OT or PT treatments provided with the goal of restoring or improving upon functions that have been lost or impaired due to injury, disease, or congenital abnormality. Rehabilitative treatments are differentiated from “habilitative treatments” in that the individual has previously met these functional milestones but has lost them due to some process.

“Habilitative Treatments” are OT or PT treatments provided with the primary goal of developing skills needed to perform ADLs or IADLs which, as a result of injury, disease, or congenital abnormality, are not developed to the normal level of functioning. This is different from rehabilitative treatment in that habilitative treatments are for individuals that have not developed to the expected level of function or have not yet met a development milestone; an example includes therapy for a child who is not talking at the expected age.

“Custodial Care” or “Long-term Care” are non-skilled, personal care to maintain the member’s ADLs or IADLS over a long-term duration and do not require oversight or skilled services by trained health professionals or technical personnel. These services are not part of a medical treatment plan for recovery, rehabilitation, habilitation, or improvement in sickness or injury. Custodial services may be provided in the home, assisted living facilities, or nursing homes, etc. This type of custodial or long-term care typically does not apply for plan benefits, please see the member’s plan benefit.

Hospice Care / End-of-Life Care - are interdisciplinary and holistic care when curative or life-prolonging treatments are no longer beneficial and services may focus on symptom control, psychosocial and spiritual care, nursing, or short-term acute services. Trained clinicians and support staff support individual and family quality-of-life goals. Hospice care can be provided in the home, skilled nursing facility, or hospital setting (for acute symptom management and stabilization to return to previous level of hospice care).

Palliative Care - are interdisciplinary and holistic care that focuses on symptom management, relieving suffering in all stages of disease, supporting communication, assessing psychosocial and spiritual resources, social and economic resources. Members may receive curative or life-prolonging treatment, and may not choose to receive hospice care or end-of-life care. Furthermore, palliative care provides support for individual and family quality-of-life goals.

Clinical Indications

Physical Therapy and Occupational Therapy in the home is medically necessary for initial requests when ALL of the following criteria are met:

1. The treatment plan is prescribed and monitored by a licensed provider(MD, DO, NP, or PA) as per individual state law and must be provided by a licensed physical or occupational therapist; *and*
2. The member meets the definition of homebound (see *Definitions* section above); *and*
3. Medical necessity criteria in the appropriate MCG Home Care Optimal Recovery Guidelines or MCG Home Care General Recovery Guideline are met; *and/or*
4. For members requesting hospice care/end-of-life care or palliative care (please check plan benefits to verify hospice or palliative care benefit timeframes), the following criteria must be met to meet medical necessity:
 - a. The member is terminally ill, presenting with functional decline, and certified by a medical practitioner for life expectancy less than twelve months for palliative care and less than six months for hospice/end-of-life care; *and*
 - b. The PT/OT services are rendered as part of a hospice care program; *and*
 - c. The member may receive curative treatment while receiving palliative care; *and/or*
 - d. The member is not receiving curative treatment while in hospice care; *and*
5. Therapy is aimed at establishing or restoring function that was lost or impaired *as a result of* disease, injury, or procedure; *and*
6. Rehab potential is evident based on a review of the member's condition, and the member's function is not expected to improve in the absence of therapy; *and*
7. Improvement can be expected with sustainable benefit in range of motion, strength, function, reduced pain level, and independence of ADLs; *and*
8. The written plan of care includes an initial evaluation and is sufficient to determine the necessity of therapy, including the following elements:
 - a. A medical evaluation (qualified provider or practitioner within scope of state specific licensure) has been conducted within 30 days of the service dates; *and*

- b. The diagnosis, the date of onset or exacerbation of the disorder/diagnosis, the duration, the severity, the anticipated course (stable, progressive or, improving), and the prognosis; *and*
 - c. Prior level of functioning and current level of functioning; *and*
 - d. Long-term and short-term goals that are specific, quantitative, objective, and attainable in no more than 3 months; *and*
 - e. The frequency and duration of proposed treatment; *and*
 - f. The specific treatment techniques and/or exercises to be used; *and*
 - g. Discharge plan.
9. Documentation of medical necessity should be reviewed when ANY of the following occur:
- a. The plan of care exceeds the expected duration and/or estimated frequency of care; *or*
 - b. There is a change in the member's condition that may impact the plan of care; *or*
 - c. The specific goals are no longer expected to be achieved in a reasonable or expected duration of time.

Duration of Therapy

The duration of therapy is dependent on the treatment plan of care and the severity of the member's condition. The Plan utilizes MCG home care criteria for the recommended visits per episode.

Extension Requests

A Plan member who requires continued PT or OT visits, beyond the original treatment plan of care, may receive extended treatment when the follow criteria are met:

1. A completed re-evaluation documented by a qualified provider/practitioner/physical or occupational therapist practicing within the scope of state specific licensure, which has been conducted within 30 days of the service dates; *and*
 - a. The appropriate MCG Home Care Optimal Recovery Guidelines with the Extended Visits criteria are met; *or*
 - b. The appropriate MCG Home Care General Recovery Guideline and the member is still in General Treatment Course Stage 2 or has not met all of the milestones in Stage 3; *or*
2. For an extension request/recertification for hospice or palliative care, please see plan benefits and requirements. The member meets medical necessity when the member continues to meet criteria as listed under Clinical Indications above in criterion 4.

Experimental or Investigational / Not Medically Necessary

Skilled care, and thus physical and occupational therapy, should be discontinued when one of the following is present:

- Homebound status is no longer met; *or*
- The member reaches the predetermined goals or skilled treatment is no longer required; *or*
- The member has reached maximum rehab potential; *or*
- The goals will not be met and there is no expectation of meeting them in reasonable time; *or*

- The member can safely and effectively continue their rehabilitation in a home exercise program; *or*
- The member's medical condition prevents further therapy; *or*
- The member refuses treatment.

Physical/Occupational therapy is not considered medically necessary for the following:

- Asymptomatic members or those without an identifiable clinical condition; *or*
- Improvement in functioning is not expected over a reasonable and predictable period of time (i.e. a "stable deficit"); *or*
- Cases of transient or easily reversible loss or reduction in function which could be reasonably expected to improve spontaneously as the member gradually resumes normal activities; *or*
- Chronic illness / chronic flare-ups or exacerbations that did not result in a decline in function or related to an acute exacerbation; *or*
- Long-term maintenance therapy, as it is aimed to preserve the present level of function or to prevent regression below an acceptable level of functioning; *or*
- Custodial care or Long-term care services; *or*
- Duplicative therapy services or programs; *or*
- Treatment modalities or home exercises that do not require a licensed PT/OT and can safely be conducted by the member alone or with the help of family or caregivers; *or*
- Therapy aimed at improving or restoring only IADLs; *or*
- Occupational or recreational programs aiming to augment or improve upon normal human functioning; this includes services considered as routine, conditioning, educational, for employment or job training, or as part of a fitness or sports program; *or*
- Pilates or general exercises to promote fitness or flexibility are not medically necessary; *or*
- Transcutaneous electrical nerve stimulator (TENS) in the home setting is not medically necessary, please refer to the Plan Clinical Guideline: Outpatient Physical Therapy & Occupational Therapy (CG044) for outpatient setting; *or*
- Neuromuscular electrical stimulation (NMES) / Electrical Muscle Stimulation in the home setting is not medically necessary, please refer to Plan Clinical Guideline: Outpatient Physical Therapy & Occupational Therapy (CG044) for outpatient setting; *or*
- Treatment modalities where the benefits of PT/OT are not adequately supported by peer literature or accepted standards of practice to be safe and effective and therefore, considered experimental or investigational include, but are not limited to:
 - Augmented soft tissue mobilization is considered investigational due to limited evidence of improved outcomes over standard techniques for soft tissue mobilization
 - Cognitive skills training to improve memory or problem solving
 - Driver/safety training
 - Equestrian therapy (hippotherapy)
 - Group therapy (criteria require *individualized* plans)
 - Kinesio taping for back pain or radicular pain is considered investigational and not clearly established in the literature

- Low level laser therapy (LLLT)
- MEDEK Therapy (Metodo Dinamico de Estimulacion Kinesica or Dynamic Method for Kinetic Stimulation)
- Microcurrent electrical nerve stimulation (MENS)
- Percutaneous electrical nerve stimulation (PENS) involves electroacupuncture (needle electrodes connected to a power source) and inserted into the skin near the site of pain. These needles are not permanently implanted, but tunneled percutaneously under ultrasound guidance into the subcutaneous tissue. Due to a lack of peer-reviewed reviewed literature supporting the effectiveness of PENS and pain reduction, ongoing clinical trials, and clinical practice guidelines stating there is insufficient evidence to recommend PENS for lower back pain, this type of procedure is considered experimental or investigational.
- Interferential stimulation or interferential current therapy (IF) is the superficial application of a medium-frequency alternating current, modulated to produce low frequencies up to 150 Hz and is considered experimental and investigational due to the lack of high grade quality literature. Devices such as neoGEN-Series System (RST-Sanexas) as a form of interferential current therapy for neuropathic pain provide ultra-high digital frequency to produce pulsed electronic signal energy waves that are delivered transcutaneously via contact electrodes. As per Hayes 2023, there are no relevant clinical studies, systematic reviews, or guidance documents supporting use of RST-Sanexas.
- The Interactive Metronome Program
- Sensory integrative techniques
- Vertebral axial decompression and motorized traction devices
- Conditions where the benefits of PT/OT are not adequately supported by peer literature or accepted standards of practice to be safe and effective and therefore, considered experimental or investigational include, but are not limited to:
 - Constipation
 - Vaginismus
 - Social functioning
 - Sexual dysfunction (erectile dysfunction, premature ejaculation)
 - Scoliosis
 - TMJ pain

Applicable Billing Codes

Codes considered medically necessary when clinical criteria are met:

CPT/HCPCS Codes considered medically necessary if criteria are met:	
<i>Code</i>	<i>Description</i>
97161	Physical therapy evaluation: low complexity, requiring these components: A history with no personal factors and/or comorbidities that impact the plan of care; An examination of body system(s) using standardized tests and measures addressing 1-2

	elements from any of the following: body structures and functions, activity limitations, and/or participation restrictions; A clinical presentation with stable and/or uncomplicated characteristics; and Clinical decision making of low complexity using standardized patient assessment instrument and/or measurable assessment of functional outcome. Typically, 20 minutes are spent face-to-face with the patient and/or family.
97162	Physical therapy evaluation: moderate complexity, requiring these components: A history of present problem with 1-2 personal factors and/or comorbidities that impact the plan of care; An examination of body systems using standardized tests and measures in addressing a total of 3 or more elements from any of the following: body structures and functions, activity limitations, and/or participation restrictions; An evolving clinical presentation with changing characteristics; and Clinical decision making of moderate complexity using standardized patient assessment instrument and/or measurable assessment of functional outcome. Typically, 30 minutes are spent face-to-face with the patient and/or family.
97163	Physical therapy evaluation: high complexity, requiring these components: A history of present problem with 3 or more personal factors and/or comorbidities that impact the plan of care; An examination of body systems using standardized tests and measures addressing a total of 4 or more elements from any of the following: body structures and functions, activity limitations, and/or participation restrictions; A clinical presentation with unstable and unpredictable characteristics; and Clinical decision making of high complexity using standardized patient assessment instrument and/or measurable assessment of functional outcome. Typically, 45 minutes are spent face-to-face with the patient and/or family.
97164	Re-evaluation of physical therapy established plan of care, requiring these components: An examination including a review of history and use of standardized tests and measures is required; and Revised plan of care using a standardized patient assessment instrument and/or measurable assessment of functional outcome Typically, 20 minutes are spent face-to-face with the patient and/or family.
97165	Occupational therapy evaluation, low complexity, requiring these components: An occupational profile and medical and therapy history, which includes a brief history including review of medical and/or therapy records relating to the presenting problem; An assessment(s) that identifies 1-3 performance deficits (ie, relating to physical, cognitive, or psychosocial skills) that result in activity limitations and/or participation restrictions; and Clinical decision making of low complexity, which includes an analysis of the occupational profile, analysis of data from problem-focused assessment(s), and consideration of a limited number of treatment options. Patient presents with no comorbidities that affect occupational performance. Modification of tasks or assistance (eg, physical or verbal) with

	assessment(s) is not necessary to enable completion of evaluation component. Typically, 30 minutes are spent face-to-face with the patient and/or family.
97166	Occupational therapy evaluation, moderate complexity, requiring these components: An occupational profile and medical and therapy history, which includes an expanded review of medical and/or therapy records and additional review of physical, cognitive, or psychosocial history related to current functional performance; An assessment(s) that identifies 3-5 performance deficits (ie, relating to physical, cognitive, or psychosocial skills) that result in activity limitations and/or participation restrictions; and Clinical decision making of moderate analytic complexity, which includes an analysis of the occupational profile, analysis of data from detailed assessment(s), and consideration of several treatment options. Patient may present with comorbidities that affect occupational performance. Minimal to moderate modification of tasks or assistance (eg, physical or verbal) with assessment(s) is necessary to enable patient to complete evaluation component. Typically, 45 minutes are spent face-to-face with the patient and/or family.
97167	Occupational therapy evaluation, high complexity, requiring these components: An occupational profile and medical and therapy history, which includes review of medical and/or therapy records and extensive additional review of physical, cognitive, or psychosocial history related to current functional performance; An assessment(s) that identifies 5 or more performance deficits (ie, relating to physical, cognitive, or psychosocial skills) that result in activity limitations and/or participation restrictions; and Clinical decision making of high analytic complexity, which includes an analysis of the patient profile, analysis of data from comprehensive assessment(s), and consideration of multiple treatment options. Patient presents with comorbidities that affect occupational performance. Significant modification of tasks or assistance (eg, physical or verbal) with assessment(s) is necessary to enable patient to complete evaluation component. Typically, 60 minutes are spent face-to-face with the patient and/or family.
97168	Re-evaluation of occupational therapy established plan of care, requiring these components: An assessment of changes in patient functional or medical status with revised plan of care; An update to the initial occupational profile to reflect changes in condition or environment that affect future interventions and/or goals; and A revised plan of care. A formal reevaluation is performed when there is a documented change in functional status or a significant change to the plan of care is required. Typically, 30 minutes are spent face-to-face with the patient and/or family.
G0129	Occupational therapy services requiring the skills of a qualified occupational therapist, furnished as a component of a partial hospitalization treatment program, per session (45 minutes or more)

G0151	Services performed by a qualified physical therapist in the home health or hospice setting, each 15 minutes
G0152	Services performed by a qualified occupational therapist in the home health or hospice setting, each 15 minutes
G0157	Services performed by a qualified physical therapist assistant in the home health or hospice setting, each 15 minutes
G0158	Services performed by a qualified occupational therapist assistant in the home health or hospice setting, each 15 minutes
G0159	Services performed by a qualified physical therapist, in the home health setting, in the establishment or delivery of a safe and effective physical therapy maintenance program, each 15 minutes
G0160	Services performed by a qualified occupational therapist, in the home health setting, in the establishment or delivery of a safe and effective occupational therapy maintenance program, each 15 minutes
S9129	Occupational therapy, in the home, per diem
S9131	Physical therapy; in the home, per diem

Codes not considered medically necessary for indications listed in this Guideline:

CPT/HCPCS Codes <u>not considered medically necessary</u>	
<i>Code</i>	<i>Description</i>
97169	Athletic training evaluation, low complexity, requiring these components: A history and physical activity profile with no comorbidities that affect physical activity; An examination of affected body area and other symptomatic or related systems addressing 1-2 elements from any of the following: body structures, physical activity, and/or participation deficiencies; and Clinical decision making of low complexity using standardized patient assessment instrument and/or measurable assessment of functional outcome. Typically, 15 minutes are spent face-to-face with the patient and/or family
97170	Athletic training evaluation, moderate complexity, requiring these components: A medical history and physical activity profile with 1-2 comorbidities that affect physical activity; An examination of affected body area and other symptomatic or related systems addressing a total of 3 or more elements from any of the following: body structures, physical activity, and/or participation deficiencies; and Clinical decision making of moderate complexity using standardized patient assessment instrument and/or measurable assessment of functional outcome. Typically, 30 minutes are spent face-to-face with the patient and/or family.

97171	Athletic training evaluation, high complexity, requiring these components: A medical history and physical activity profile, with 3 or more comorbidities that affect physical activity; A comprehensive examination of body systems using standardized tests and measures addressing a total of 4 or more elements from any of the following: body structures, physical activity, and/or participation deficiencies; Clinical presentation with unstable and unpredictable characteristics; and Clinical decision making of high complexity using standardized patient assessment instrument and/or measurable assessment of functional outcome. Typically, 45 minutes are spent face-to-face with the patient and/or family.
97172	Re-evaluation of athletic training established plan of care requiring these components: An assessment of patient's current functional status when there is a documented change; and A revised plan of care using a standardized patient assessment instrument and/or measurable assessment of functional outcome with an update in management options, goals, and interventions. Typically, 20 minutes are spent face-to-face with the patient and/or family.
E0746	Electromyography (EMG), biofeedback device
S8990	Physical or manipulative therapy performed for maintenance rather than restoration
S9117	Back school, per visit

Codes considered experimental or investigational:

CPT/HCPCS Codes considered experimental or investigational	
<i>Code</i>	<i>Description</i>
97150	Therapeutic procedure(s), group (2 or more individuals)
S8940	Equestrian/hippotherapy, per session
S8948	Application of a modality (requiring constant provider attendance) to one or more areas, low-level laser; each 15 minutes
S9090	Vertebral axial decompression, per session

Codes for electrical stimulation considered Not Medically Necessary (TENS, NMES/electrical muscle stimulation) for the home setting:

CPT/HCPCS Codes <u>not considered medically necessary</u>	
<i>Code</i>	<i>Description</i>
97014	Application of a modality to 1 or more areas; electrical stimulation (unattended)

97032	Application of a modality to 1 or more areas; electrical stimulation (manual), each 15 minutes
A4558	Conductive gel or paste, for use with electrical device (e.g., TENS, NMES), per oz
A4595	Electrical stimulator supplies, 2 lead, per month, (e.g., TENS, NMES)
E0720	Transcutaneous electrical nerve stimulation (TENS) device, two-lead, localized stimulation.
E0730	Transcutaneous electrical nerve stimulation (TENS) device, four or more leads, for multiple nerve stimulation
E0731	Form-fitting conductive garment for delivery of TENS or NMES (with conductive fibers separated from the patient's skin by layers of fabric)
E0745	Neuromuscular stimulator, electronic shock unit [NMES]
E0769	Electrical stimulation or electromagnetic wound treatment device, not otherwise classified
G0281	Electrical stimulation, (unattended), to one or more areas, for chronic Stage III and Stage IV pressure ulcers, arterial ulcers, diabetic ulcers, and venous stasis ulcers not demonstrating measurable signs of healing after 30 days of conventional care, as part of a therapy plan of care
G0282	Electrical stimulation, (unattended), to one or more areas, for wound care other than described in G0281
G0283	Electrical stimulation (unattended), to one or more areas for indication(s) other than wound care, as part of a therapy plan of care

CPT/HCPCS Codes for electrical stimulation considered experimental or investigational (Interferential therapy, MENS, PENS):	
<i>Code</i>	<i>Description</i>
64999	<p>Unlisted procedure, nervous system</p> <ul style="list-style-type: none"> • <u>Due to the broad nature of this code and lack of specificity in certain scenarios, clarification is provided below:</u> • When this code is billed for percutaneous electrical nerve stimulation (PENS), it is considered experimental or investigational
E1399	<p>Durable medical equipment, miscellaneous</p> <ul style="list-style-type: none"> • <u>Due to the broad nature of this code and lack of specificity in certain scenarios, clarification is provided below:</u>

	<ul style="list-style-type: none"> When this code is billed for microcurrent electrical nerve stimulation (MENS), it is considered experimental or investigational
S8130	Interferential current stimulator, 2 channel
S8131	Interferential current stimulator, 4 channel

References

- The American College of Obstetrics and Gynecologists. (July 2019). Female Sexual Dysfunction. Practice Bulletin Number 213.
<https://www.acog.org/clinical/clinical-guidance/practice-bulletin/articles/2019/07/female-sexual-dysfunction>
- CMS Medicare Benefit Policy Manual Chapter 7 - Home Health Services. CMS.gov.
<https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c07.pdf>
Revised 05/7/2021.
- Centers for Medicare & Medicaid Services. Medicare Benefit Policy Manual Chapter 9 - Coverage of Hospice Services Under Hospital Insurance. Revised 10/21/21.
<https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c09.pdf>
- CMS LCD Guidelines. <http://www.cms.gov/mcd/index>. Accessed on Feb 23, 2016.
 - Local Coverage Determination (LCD) for PHYSICAL THERAPY - Home Health (L33942)
 - Local Coverage Determination (LCD) for PHYSICAL THERAPY - Home Health (L34564)
 - Local Coverage Determination (LCD) for Outpatient PHYSICAL and Occupational THERAPY Services Health (L34049)
 - Local Coverage Determination (LCD) for Medicine: PHYSICAL THERAPY - Outpatient (L34310)
 - Local Coverage Determination (LCD) for Home Health OCCUPATIONAL THERAPY (L34560)
 - Local Coverage Determination (LCD) for Medicine: OCCUPATIONAL THERAPY - Outpatient (L34308)
- American Occupational Therapy Association. (2021). Standards of practice for occupational therapy. *The American Journal of Occupational Therapy*, 75(Supplement_3).
- American Physical Therapy Association. Standards of practice for physical therapy. Last updated Aug 12,2020. Available at:
<https://www.apta.org/apta-and-you/leadership-and-governance/policies/standards-of-practice-p-t>
- Bavarian R, Khawaja SN, Ajisafe AH, Sultan AS. The efficacy of microcurrent electrical nerve stimulation in treating masticatory myofascial pain: A systematic review and meta-analysis. *The Journal of Craniomandibular & Sleep Practice*. 2021;
doi: <https://doi.org/10.1080/08869634.2021.2020006>
- Cejudo P, Bautista J, Montemayor T, et al. Exercise training in mitochondrial myopathy: A randomized controlled trial. *Muscle Nerve*. 2005;32(3):342-350.

9. Chou R. Subacute and chronic low back pain: Nonpharmacologic and pharmacologic treatment. UpToDate.com. Last updated: Jul 18, 2023. Accessed from https://www.uptodate.com/contents/subacute-and-chronic-low-back-pain-nonpharmacologic-and-pharmacologic-treatment?search=Nonpharmacologic%20and%20pharmacologic%20treatment&source=search_result&selectedTitle=1%7E150&usage_type=default&display_rank=1
10. Cospser SM, Lee GP, Peters SB, Bishop E. Interactive Metronome training in children with attention deficit and developmental coordination disorders. *Int J Rehabil Res.* 2009;32(4):331-336.
11. de Freitas, G. R., Szpoganicz, C., & Ilha, J. (2018). Does neuromuscular electrical stimulation therapy increase voluntary muscle strength after spinal cord injury? A systematic review. *Topics in spinal cord injury rehabilitation.* *Top Spinal Cord Inj Rehabil.* 24(1), 6-17.
12. De Oliveira, G.R., & Vidal MF. (2021). Developmental outcomes in a child with corpus callosum abnormalities and congenital heart disease after Cuevas Medek Exercises: A case report. *Clinical Case Reports.* Doi: <https://doi.org/10.1002/ccr3.4637>
13. Department of Veterans Affairs. (2022). VA/DoD CLINICAL PRACTICE GUIDELINE FOR DIAGNOSIS AND TREATMENT OF LOW BACK PAIN. Version 3.0. Retrieved from <https://www.healthquality.va.gov/guidelines/Pain/lbp/VADoDLBPCPGFinal508.pdf>
14. Feinstein A. Conversion Disorder. *Continuum (Minneapolis, Minn).* 2018 Jun;24(3, BEHAVIORAL NEUROLOGY AND PSYCHIATRY):861-872.
15. Fitzgerald MP, Anderson RU, Potts J, et al; Urological Pelvic Pain Collaborative Research Network. Randomized multicenter feasibility trial of myofascial physical therapy for the treatment of urological chronic pelvic pain syndromes. *J Urol.* 2013;189(1 Suppl):S75-S85.
16. Frontera W, Silver J, Rizzo TD editors. *Essentials of Physical Medicine and Rehabilitation.* 4th ed. Philadelphia, PA: Elsevier, 2018.
17. Fu TC, Wong AM, Pei YC, Wu KP, Chou SW, Lin YC. Effect of Kinesio taping on muscle strength in athletes-a pilot study. *J Sci Med Sport.* 2008 Apr;11(2):198-201.
18. Yu GH, Lee JS, Kim SK, & Cha TH. (2017). Effects of interactive metronome training on upper extremity function, ADL and QOL in stroke patients. *NeuroRehabilitation.* 2017; 41(1), pp. 161-168. DOI: 10.3233/NRE-171468
19. García-Collado A, Valera-Calero JA, Fernández-de-Las-Peñas C, Arias-Burúa JL. Effects of ultrasound-guided nerve stimulation targeting peripheral nerve tissue on pain and function: a scoping review. *J Clin Med.* 2022;11(13):3753. doi:10.3390/jcm11133753
20. González-Iglesias J, Fernández-de-Las-Peñas C, Cleland JA, et al. Short-term effects of cervical kinesio taping on pain and cervical range of motion in patients with acute whiplash injury: A randomized clinical trial. *J Orthop Sports Phys Ther.* 2009;39(7):515-521.
21. Gulick DT. Comparison of tissue heating between manual and hands-free ultrasound techniques. *Physiother Theory Pract.* 2010;26(2):100-106.
22. Hayes, Inc. Evolving Evidence Review. NeoGEN-Series System (RST-Sanexas) for Treatment of Neuropathic Pain. Lansdale, PA: Hayes, Inc.; Annual review: Jan 2023.
23. Hayes, Inc. Evidence Analysis Research Brief. *Percutaneous Electrical Nerve Stimulation for Treatment of Low Back Pain.* Lansdale, PA: Hayes, Inc.; Dec 22, 2022.

24. Jansen M, de Groot IJ, van Alfen N, Geurts ACh. Physical training in boys with Duchenne muscular dystrophy: The protocol of the No Use is Disuse study. *BMC Pediatr.* 2010;10:55.
25. Katalinic OM, Harvey LA, Herbert RD. Effectiveness of stretch for the treatment and prevention of contractures in people with neurological conditions: A systematic review. *Phys Ther.* 2011;91(1):11-24.
26. Kong LJ, Zhan HS, Cheng YW, Yuan WA, Chen B, Fang M. Massage therapy for neck and shoulder pain: a systematic review and meta-analysis. *Evid Based Complement Alternat Med.* 2013;2013:613279.
27. Krauss I, Steinhilber B, Haupt G, et al. Efficacy of conservative treatment regimes for hip osteoarthritis -- evaluation of the therapeutic exercise regime "Hip School": A protocol for a randomised, controlled trial. *BMC Musculoskelet Disord.* 2011;12:270.
28. Landers S, Madigan E, Leff B, Rosati RJ, McCann BA, et al. The Future of Home Health Care. A Strategic Framework for Optimizing Value. *Home Health Care Manag Pract.* 2016; 28(4): 262–278. doi: 10.1177/1084822316666368
29. Latham NK, Harris BA, Bean JF, et al. Effect of a Home-Based Exercise Program on Functional Recovery Following Rehabilitation After Hip Fracture: A Randomized Clinical Trial. *JAMA.* 2014;311(7):700–708. doi:10.1001/jama.2014.469
30. Melham TJ, Sevier TL, Malnofski MJ, et al. Chronic ankle pain and fibrosis successfully treated with a new noninvasive augmented soft tissue mobilization technique (ASTM): A case report. *Med Sci Sports Exerc.* 1998;30(6):801-804.
31. Michaleff ZA, Maher CG, Lin CW, et al. Comprehensive physiotherapy exercise programme or advice for chronic whiplash (PROMISE): A pragmatic randomised controlled trial. *Lancet.* 2014;384(9938):133-141.
32. Minor MA, Sanford MK. The role of physical therapy and physical modalities in pain management. *Rheum Dis Clin North Am.* 1998;25(1):233-248, viii
33. National Association for Home Care & Hospice. *What is Hospice and Palliative Care?* <https://www.nahc.org/about/faq/#111> Accessed: March 8, 2022.
34. National Hospice and Palliative Care Organization. (2021). *NHPCO Facts and Figures Report, 2021 edition.* <https://www.nhpco.org/hospice-care-overview/hospice-facts-figures/> Accessed March 8, 2022.
35. National Institute for Health and Care Excellence. (2016). Low back pain and sciatica in over 16s: assessment and management [NG59]. Retrieved from <https://www.nice.org.uk/guidance/ng59>
36. Nair HKR. Microcurrent as an adjunct therapy to accelerate chronic wound healing and
 - a. 2 reduce patient pain. *J Wound Care.* 2018 May 2;27(5):296-306.
37. Pontari M, Giusto L. New developments in the diagnosis and treatment of chronic prostatitis/chronic pelvic pain syndrome. *Curr Opin Urol.* 2013;23(6):565-569.
38. Rajpurohit B, Khatri SM, Metgud D, Bagewadi A. Effectiveness of transcutaneous
 - a. 38 electrical nerve stimulation and microcurrent electrical nerve stimulation in bruxism
 - b. 39 associated with masticatory muscle pain--a comparative study. *Indian J Dent Res.* 2010
 - c. 40 Jan-Mar;21(1):104-6.

39. Rampazo ÉP, Liebano RE. Analgesic Effects of Interferential Current Therapy: A Narrative Review. *Medicina*. 2022; 58(1):141. <https://doi.org/10.3390/medicina58010141>
40. Sakzewski L, Ziviani J, Boyd R. Systematic review and meta-analysis of therapeutic management of upper-limb dysfunction in children with congenital hemiplegia. *Pediatrics*. 2009 Jun;123(6):e1111-22.
41. Scherl SA. Adolescent idiopathic scoliosis: Management and prognosis. UpToDate, Waltham, MA. Reviewed February 2017.
42. Shank TM, & Harron W. A retrospective outcomes study examining the effect of interactive metronome on hand function. *Journal of Hand Therapy*. 2015; 28 (4): 396-402. Doi: <https://doi.org/10.1016/j.jht.2015.06.003>
43. Sheon RP, Duncombe AM. Rehabilitation program for the low back. UpToDate. Waltham, MA: UpToDate; reviewed February 2017.
44. Taylor NF, Dodd KJ, Shields N, Bruder A. Therapeutic exercise in physiotherapy practice is beneficial: a summary of systematic reviews 2002-2005. *Aust J Physiother*. 2007;53(1):7-16
45. Thomaz, S. R., Cipriano Jr, G., Formiga, M. F., Fachin-Martins, E., Cipriano, G. F. B., Martins, W. R., & Cahalin, L. P. (2019). Effect of electrical stimulation on muscle atrophy and spasticity in patients with spinal cord injury—a systematic review with meta-analysis. *Spinal cord*, 57(4), 258-266.
46. Trenell MI, Sue CM, Kemp GJ, et al. Aerobic exercise and muscle metabolism in patients with mitochondrial myopathy. *Muscle Nerve*. 2006;33(4):524-531.
47. Urological aspects of chronic pelvic pain. In: Engeler D, Baranowski AP, Elneil S, et al. Guidelines on chronic pelvic pain. Arnhem, The Netherlands: European Association of Urology (EAU); February 2012.
48. Veerbeek JM, van Wegen E, van Peppen R, et al. What is the evidence for physical therapy poststroke? A systematic review and meta-analysis. *PLoS One*. 2014;9(2):e87987.
49. Legg LA, Drummond AE, Langhorne P. Occupational therapy for patients with problems in activities of daily living after stroke. *Cochrane Database Syst Rev*. 2006;(4):CD003585.
50. Reitz SM, Austin DJ, Brandt LC, et al. Guidelines to the Occupational Therapy Code of Ethics. *Am J Occup Ther*. 2006; 60(6):652-668.
51. Steultjens EM, Dekker JJ, Bouter LM, et al. Evidence of the efficacy of occupational therapy in different conditions: an overview of systematic reviews. *Clin Rehabil*. 2005; 19(3):247-254.
52. Steultjens EM, Dekker JJ, Bouter LM, et al. Occupational therapy for rheumatoid arthritis. *Cochrane Database of Syst Rev*. 2004;(1):CD003114.
53. Trombly CA, Ma HI. A synthesis of the effects of occupational therapy for persons with stroke, Part I: Restoration of roles, tasks, and activities. *Am J Occup Ther*. 2002 May-Jun;56(3):250-9. 20.
54. Walker MF, Leonardi-Bee J, Bath P, Langhorne P, Dewey M, Corr S, et al. Individual patient data metaanalysis of randomized controlled trials of community occupational therapy for stroke patients. *Stroke*. 2004 Sep;35(9):2226-32.
55. Cheatham SW, Lee M, Cain M, Baker R. The efficacy of instrument assisted soft tissue mobilization: a systematic review. *J Can Chiropr Assoc*. 2016;60(3):200-211.

56. Halseth T, McChesney JW, DeBeliso M, et al. The Effects of Kinesio Taping on proprioception at the ankle. *J Sports Sci Med*. 2004;3:1-7.
57. Yasukawa A, Patel P, Sisung C. Pilot study: Investigating the effects of Kinesio Taping in an acute pediatric rehabilitation setting. *Am J Occup Ther*. 2006;60(1):104-110.
58. Fu TC, Wong AM, Pei YC, et al. Effect of Kinesio taping on muscle strength in athletes-a pilot study. *J Sci Med Sport*. 2008;11(2):198-201.
59. Thelen MD, Dauber JA, Stoneman PD. The clinical efficacy of kinesio tape for shoulder pain: A randomized, double-blinded, clinical trial. *J Orthop Sports Phys Ther*. 2008;38(7):389-395.
60. Scherl SA. Adolescent idiopathic scoliosis: Management and prognosis. UpToDate. <https://www.uptodate.com/contents/adolescent-idiopathic-scoliosis-management-and-prognosis> Accessed April 3, 2017.
61. Meier DE, Back AL, Berman A, Block SD, et al. A National Strategy For Palliative Care. *Health Affairs*. 2017; 36(7). Doi: <https://doi.org/10.1377/hlthaff.2017.0164>
62. Meier DE, McCormick E & Lagman RL. (Dec 7, 2020). *Hospice: Philosophy of care and appropriate utilization in the United States*. UpToDate.com. Accessed March 4, 2022.
63. McNeely ML, Armijo Olivo S, Magee DJ. A systematic review of the effectiveness of physical therapy interventions for temporomandibular disorders. *Phys Ther*. 2006; 86:710.
64. McKivigan, J., Krisak, B., Vogel, B., Harrington, A., & Nguyen, V. (2019). Evaluating the Efficacy of Vertebral Axial Decompression Therapy in Treatment of Patients with Chronic Lower Back Pain: A Systematic Review. *Rehabilitation*, 4(2), 25-34.
65. Nazligul T, Akpınar P, Aktas I, Unlu Ozkan F, Cagliyan Hartevioglu H. The effect of interferential current therapy on patients with subacromial impingement syndrome: a randomized, double-blind, sham-controlled study. *Eur J Phys Rehabil Med*. 2018 Jun;54(3):351-357. doi: 10.23736/S1973-9087.17.04743-8. Epub 2017 Sep 11. PMID: 28895673.
66. Scrivani SJ. Temporomandibular disorders in adults. UpToDate. <https://www.uptodate.com/contents/temporomandibular-disorders-in-adults>. Accessed April 3, 2017.
67. STECKO, Monika Joanna, WAWRYKÓW, Agata, KORABIUSZ, Katarzyna, KORDEK, Agnieszka, FABIAN - DANIELEWSKA, Anna, WILCZYŃSKA, Agnieszka, JANIK, Inga & MACIEJEWSKA, Martyna. The methods of therapy of children suffering from cerebral palsy. *Journal of Education, Health and Sport* [online]. 9 September 2018, T. 8, nr 9, s. 562–569. [accessed March 9, 2022].
68. King, PM, Myers, CA, Ling, FW, Rosenthal, RH. Musculoskeletal factors in chronic pelvic pain. *J Psychosom Obstet Gynaecol*. 1991; 12:87.
69. Romano M, Minozzi S, Bettany-Saltikov J, et al. Exercises for adolescent idiopathic scoliosis. *Cochrane Database Syst Rev*. 2012; :CD007837.
70. Bylund P, Aaro S, Gottfries B, Jansson E. Is lateral electric surface stimulation an effective treatment for scoliosis? *J Pediatr Orthop*. 1987; 7:298.
71. Ye L, Kalichman L, Spittle A, Dobson F, Bennell K. Effects of rehabilitative interventions on pain, function and physical impairments in people with hand osteoarthritis: a systematic review. *Arthritis Res Ther*. 2011 Feb 18;13(1):R28.

72. van Reijn-Baggen DA, Han-Geurts IJM, Voorham-van der Zalm PJ, Pelger RCM, Hagenaars-van Miert CHAC, Laan ETM. Pelvic Floor Physical Therapy for Pelvic Floor Hypertonicity: A Systematic Review of Treatment Efficacy. *Sex Med Rev.* 2022 Apr;10(2):209-230. doi: 10.1016/j.sxmr.2021.03.002. Epub 2021 Jun 12. PMID: 34127429.
73. Wong MS, Mak AF, Luk KD, et al. Effectiveness of audio-biofeedback in postural training for adolescent idiopathic scoliosis patients. *Prosthet Orthot Int.* 2001; 25:60.

[Clinical Guideline Revision / History Information](#)

Original Date: 4/11/2017

Reviewed/Revised: 1/18/2018, 4/13/2018, 4/15/2019, 7/21/2020, 04/21/2021, 12/01/2021, 04/25/2022, 05/02/2023, 04/16/2024