

Pain Management: Facet Joint Injections and Radiofrequency Facet Denervation

Disclaimer

Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Oscar may delegate utilization management decisions of certain services to third-party delegates, who may develop and adopt their own clinical criteria.

The clinical guidelines are applicable to all commercial plans. Services are subject to the terms, conditions, limitations of a member's plan contracts, state laws, and federal laws. Please reference the member's plan contracts (e.g., Certificate/Evidence of Coverage, Summary/Schedule of Benefits) or contact Oscar at 855-672-2755 to confirm coverage and benefit conditions.

Summary

Oscar members with chronic back and neck pain may qualify for diagnostic and therapeutic procedures to further characterize or treat their pain. When the back pain is non-radiating and not associated with any neurologic deficits, the facet joints may be implicated as the cause. Facet joints are small joints between the bones that make up the spine that allow for bending and twisting motions. Small nerves run through these joints and may cause pain when the joint is inflamed or irritated. To determine whether the facet joint is involved, small amounts of anesthetic can be injected directly into the joint or to the surrounding nerves. If the joint is the cause of the pain, the injection should result in relief. If the pain recurs after this initial injection, a special procedure to destroy the nerve causing the pain may be indicated.

For pain management with steroid injections and nerve blocks, please see CG048: Pain Management: Epidural Steroid Injections, Selective Nerve Root Blocks (SNRB), and Intradiscal Steroid Injections.

Definitions

"Facet Joints" are where two bones of the vertebra in the spine interlock. They are thought to be a major cause of chronic back pain. Injections with steroids or anesthetics into the facet joint may help reduce pain and/or inflammation in this area, known as facet nerve block. Symptoms consistent with "facet joint syndrome" include:

- Absence of radiculopathy
- Predominantly axial low back pain
- Pain aggravated by extension, rotation, or lateral bending of the back or neck

- Tenderness to palpation in the paravertebral areas (over the facet region)
- An absence of neurologic deficits (in a dermatomal distribution)
- Physical exam findings and diagnostic imaging reveal no other obvious cause of the pain

“Facet Joint Injection” (i.e., intra-articular facet injection) is where medications to reduce swelling and/or pain are injected into the facet joint. Injections can be done for therapeutic purposes, where the medication is injected with goal of long-term pain alleviation, or diagnostic purposes, where the injection is done to confirm the facet joint as the cause of the patient’s pain.

“Medial Branch Nerve Block” is very similar to a facet joint injection and often used for the same indications, except instead of injecting into the joint space, the anesthetic is injected into the space around the supplying nerve to the facet joint known as the “median nerve”.

“Straight Leg Raise Test” is a simple office test where the patient lies on his or her back, and one leg is raised upwards while held straight. The test is deemed positive when there is radiating pain down the leg as it is raised between 30 degrees and 70 degrees. This suggests a herniated disk or other radiculopathy, and is not typically observed in facet joint syndromes.

“Radicular Pain” (i.e., radiculopathy) refers to pain that radiates along the course of a spinal nerve root. Signs of radicular pain include positive straight leg test, dermatomal pattern of sensory loss, pain or numbness radiating below the level of the knee (lumbar), pain or numbness in the shoulder, arm, hand, or fingers (cervical), or diminished reflexes. Radicular pain is not characteristic of facet joint syndromes.

“Non-Radicular Pain” is pain that does not radiate and is typically most intense local to the source, rather than spreading in a predictable distribution like radicular pain. Facet joint syndrome typically cause this type of pain.

“Denervation” (i.e., ablation) is a procedure where a nerve supply is destroyed or impaired, often to treat pain syndromes. Examples of denervation include but are not limited to:

- Thermal (non-pulsed) radiofrequency denervation
- Chemical neurolysis (e.g., alcohol, phenol, hypertonic saline)
- Cooled or pulsed radiofrequency denervation
- Cryosurgery
- Surgical facet denervation (i.e., facet neurotomy, facet rhizotomy)
- Laser ablation

“Radiofrequency Facet Denervation” (also known as radiofrequency ablation or neurolysis) is a procedure for facet-mediated pain where a small radiofrequency probe is inserted to a nerve branch of the facet joint using fluoroscopic imaging. The probe heats up the nerve to denervate the supply to the painful joint. The heat can be supplied continuously or in an intermittent, pulsed fashion. Pulsed methods typically do not “ablate” the nerve.

“Fluoroscopy” is an x ray technique used to visualize internal structures.

“Activities of Daily Living (ADLs)” are defined as routine activities that most healthy persons perform daily without requiring assistance. These include, but are not limited to: bathing, communication, dressing, feeding, grooming, mobility, personal hygiene, self-maintenance, skin management, and toileting.

Covered Services and Clinical Indications

Diagnostic Facet Joint Injections

Oscar covers diagnostic facet joint injections when ALL of the following criteria are met:

1. Presence of non-radicular lower back and/or neck pain that is suggestive of facet joint syndromes, as defined above; *and*
2. Pain interferes with ADLs; *and*
3. Pain has not responded to at least 3 months of conservative therapy, as defined by the following:
 - a. Trial of appropriate medications (e.g., NSAIDs, analgesics, etc.); *and*
 - b. Physical therapy, spinal manipulation therapy, or other specific interventions tailored to the member’s unique presentation.
4. There are no current medical problems that may increase the risk of side effects, including but not limited to the following:
 - a. Local infection at the injection site; *or*
 - b. Systemic infection; *or*
 - c. Increased bleeding risk, whether medication induced or organic in etiology; *or*
 - d. Any other unstable medical condition such as cauda equina or vertebral fracture.
5. Alternative causes of the pain such as intraspinal tumor, other space occupying spinal lesion, or non-spinal etiologies of the pain have been ruled out and/or adequately addressed; *and*
6. There has been no vertebral fusion at the levels proposed for treatment; *and*
7. The injection meets ALL of the following criteria:
 - a. Performed as a diagnostic test with temporary, local anesthetic, to identify the source of the back pain; *and*
 - b. The injection is to the cervical or lumbar spine; *and*
 - c. Real time imaging guidance is used (CT scan or fluoroscopy) to ensure proper needle position; *and*
 - d. No more than 3 levels are injected per session; *and*
 - e. No other injections are performed the same day (e.g., epidural, trigger point injection).
8. Radiofrequency facet denervation is being considered should the diagnostic test be positive.

Initial Non-Pulsed Radiofrequency Facet Denervation

Oscar covers the initial non-pulsed radiofrequency facet denervation of the cervical and/or lumbar spine (with fluoroscopy) when ALL of the following criteria are met:

1. Member meets criteria 1-6 in the facet injection section above; *and*
2. The pain has been severe and limited ADLs for at least 6 months duration; *and*
3. Imaging studies show no disk herniation or obvious source of the pain; *and*
4. There is no significant narrowing of the vertebral canal or spinal instability requiring surgery; *and*
5. A diagnostic facet joint injection meeting the above criteria resulted in significant reduction in the pain, as defined by >50% reduction in pain and/or symptoms for at least 2 weeks; *and*
6. No more than 3 levels of facet joints are injected in a single session.

Additional Non-Pulsed Radiofrequency Facet Denervation

Oscar covers additional non-pulsed radiofrequency facet denervation of the spine (with fluoroscopy) when ALL of the following criteria are met:

1. All of the criteria for the original denervation continue to be met; *and*
2. The last successful procedure was less than 1 year ago, where another diagnostic facet joint injection would be required for durations >1 year; *and*
3. The previous treatment resulted in at least 50% pain relief for a duration of at least 12 weeks; *and*
4. No more than 2 procedures are performed in a rolling 12 month period, with at least 6 months interval between them; *and*
5. If different regions require neural blockade, these should be performed at intervals of no sooner than one week, and preferably 2 weeks for most blocks.

Coverage Exclusions

Facet joint injections for any other indication are *not covered* by Oscar, as they are considered experimental, investigational, or unproven. Non-covered indications include, but are not limited to, the following:

- Facet joint injections for therapeutic purposes
- Sacroiliac facet injection
- Thoracic facet injection
- Ultrasound or MRI guidance
- Diagnostic injection when RFA is not being considered
- Injections for radicular back pain

Radiofrequency denervation for any other indication is *not covered* by Oscar, as it is considered experimental, investigational, unproven, or not medically necessary. Non-covered indications include, but are not limited to, the following:

- Ablation without fluoroscopic guidance, as this is integral to identifying the proper location for injection
- Pulsed radiofrequency

- Endoscopic RFA or endoscopic dorsal ramus rhizotomy
- Ablation for SI joint pain
- Ablation for thoracic facet joints
- RFA for coccygodynia, dorsal root ganglia, terminal (peripheral) nerve endings
- Cooled RFA
- Ultrasound or MRI guidance

The following treatments and procedures for facet joint pain are *not covered* by Oscar, as they are considered experimental, investigational, unproven, or not medically necessary. Non-covered treatments include, but are not limited to, the following:

- Cryoablation (i.e., cryoneurolysis, cryodenervation)
- Chemical ablation or percutaneous alcohol ablation
- Laser ablation
- Monitored anesthesia care (MAC)
- Epidurography

Evidence for Non-Coverage of Above Indications

Facet joint Injections for therapeutic purposes: Facet joint injections are recommended as a diagnostic tool for patients meeting specific criteria. Injections for therapeutic or treatment purposes are not currently recommended given the existing literature. UpToDate clinical review states “Glucocorticoid injections into the facet joint have not been shown to be effective in the treatment of low back pain.” Further, a systematic review and technology assessment (AHRQ, 2015) looked at injection therapies for lower back pain, including 13 trials for facet joint injections specifically. They found no differences between facet joint corticosteroid injections and placebo interventions in the treatment of back pain. Further well designed randomized trials published in the New England Journal of Medicine have found no difference in pain relief following saline (placebo) and glucocorticoids. Expert consensus guidelines from the American Association of Neurological Surgeons, the American Pain Society, and others, overwhelmingly recommend against therapeutic facet joint injection.^{2,6,8-10,}

Sacroiliac facet injection: Similar to the data for SI joint RFA procedures, there is limited evidence for sacroiliac facet injections. An UpToDate review of the literature concluded that “Effective methods for diagnosing and treating sacroiliac joint pain in patients without spondyloarthropathy remain controversial”, citing that “[t]here are no randomized trials of intraarticular sacroiliac joint steroid injection versus a sham procedure”. Hansen et al conducted a systematic review of therapeutic sacroiliac joint interventions in 2012. They reviewed 56 studies, finding that 11 were of quality to be included (6 randomized, 5 non-randomized). They concluded that “the evidence for effectiveness of intraarticular steroid injections is poor. The evidence for periarticular injections of local anesthetic and steroid or botulinum toxin is poor. The evidence for effectiveness of conventional radiofrequency neurotomy is poor. The evidence for pulsed radiofrequency is poor.” Another review in 2015 by Kennedy et al looked at 39 studies, concluding that it was “unclear whether image-guided intra-articular diagnostic injections

of local anesthetic predict positive responses to therapeutic agents.” Another study by Scholten et al in 2015 found that SI joint injections showed pain reduction at 2 weeks but no difference in symptom scores compared to baseline at 8 weeks, demonstrating that long-term benefit was lacking. Furthermore, the best technique and approach to SIJ injections is not yet well established in the literature.^{10, 62-64}

Thoracic facet injection: A 2005 systematic review by Sehgal et al concluded that the diagnostic accuracy of thoracic facet joint injections was rated as “moderate”, compared to “strong” for lumbar due to a lack of evidence. Further studies by Manchikanti et al and Revel et al have found the false positive rate with single nerve blocks to be in the range of “17%- 47% in the lumbar spine, 27%-63% in the cervical spine, and 55%-58% in the thoracic spine”, as summarized in a recent review. The consistently >50% false positive rate in the thoracic spine is of particular concern. At the present time, there is limited randomized, controlled data on the efficacy, and major consensus guidelines do not currently offer recommendations for thoracic facet joint injections.⁵⁷⁻⁶¹

Ultrasound guidance: The data on ultrasound guidance for facet joint injections is limited. One study by Jee et al (2014) randomized patients to ultrasound guided vs. fluoroscopy guided SI joint injections and found greater accuracy in the fluoroscopy guided approach (98% vs. 87%). Further research is needed to identify any potential role for ultrasound.²⁴

Diagnostic injection when RFA is not being considered: Diagnostic facet nerve injection without consideration for treatment should not be performed as the results are used to guide treatment decisions.

Injections for radicular back pain: Facet joint pain has been defined as a non-radicular pain distribution, whereas radicular joint pain is typically due to other etiologies. Tachihara et al (1976) looked at a rat model of facet joint inflammation and found that it could potentially result in an inflammatory spread to nearby nerves, resulting in radiculopathy. However, the evidence for radicular back pain in human subjects with facet joint syndromes is limited.⁴⁷

Ablation without fluoroscopic guidance, as this is integral to identifying the proper location for injection.

Pulsed radiofrequency methods: Pulsed radiofrequency has been suggested as a non-ablative alternative to standard thermal radiofrequency ablation. The existing literature is limited to small or non-randomized studies. One study by Kroll et al (2008) randomized 50 patients to pulsed vs. continuous radiofrequency ablation for the treatment of lumbar back pain, but found no significant difference in outcomes between the two groups. Expert review highlights the lack of evidence for a mechanism of pain relief with pulse radiofrequency. The existing data is limited and insufficient to support the efficacy and duration of pulse radiofrequency for the treatment of chronic back pain.^{1, 29, 51, 55}

Endoscopic RFA or endoscopic dorsal ramus rhizotomy: The current data is limited to abstracts and small, retrospective studies. Haufe et al (2010) looked at the efficacy of this technique in retrospective form, finding that 68-78% of patients had >50% improvement in their pain, but conclude that further randomized data is needed to determine the clinical benefit.²²

Ablation for SI joint pain: The existing data is relatively limited to small patient populations. One study by Cohen et al (2008) looking at 28 patients found improved short-term response in patients treated with RFA for SI joint pain compared to local block alone, but the duration of efficacy was limited to only 14% showing continued pain relief at 1 year. Other studies have faced similar limitations and the existing evidence is insufficient to support the use of RFA for this indication.^{3, 13, 21, 23, 26-27, 46}

Ablation for thoracic facet joints: Systematic reviews have showed only two studies looking specifically at thoracic RFA, concluding that the evidence is currently limited and further research is needed to define the clinical role³⁴

RFA for coccygodynia, dorsal root ganglia, terminal (peripheral) nerve endings: There is a lack of published evidence supporting RF ablation for these indications

Cooled RFA: There is a lack of published evidence supporting cooled RF ablation in facet joint pain.⁷

Cryoablation (i.e., cryoneurolysis, cryodenervation): Several studies have looked at cryoablation for chronic back pain. One study by Birkenmaier et al (2007) looked at 50 patients with low back pain treated with cryoablation, finding improvements in back pain at different time intervals. However, failure rate at 12 months was 43%.⁵

Chemical ablation or percutaneous alcohol ablation: There is a lack of published evidence looking at the effectiveness and safety of chemodenervation in facet joint pain. Where radiofrequency ablation is targeted and local, chemical ablation may not be as focused and may theoretically lead to more toxicity.¹⁰

Laser ablation: There is a lack of published evidence supporting laser ablation for denervation in facet joint pain.⁵⁶

Monitored anesthesia care (MAC): MAC is not considered medically necessary for radiofrequency procedures or diagnostic facet nerve injections as these procedures can be managed with local anesthesia administered by an outpatient provider.

[Applicable Billing Codes \(HCPCS/CPT Codes\)](#)

<i>Facet Joint Injections</i>	
CPT/HCPCS Codes covered if criteria are met:	
<i>Code</i>	<i>Description</i>
64490	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with image guidance (fluoroscopy or CT), cervical or thoracic; single level [covered for cervical only]
64491	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with image guidance (fluoroscopy or CT), cervical or thoracic; second level (List separately in addition to code for primary procedure)
64492	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with image guidance (fluoroscopy or CT), cervical or thoracic; third and any additional level(s) (List separately in addition to code for primary procedure)
64493	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with image guidance (fluoroscopy or CT), lumbar or sacral; single level [covered for lumbar only]
64494	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with image guidance (fluoroscopy or CT), lumbar or sacral; second level (List separately in addition to code for primary procedure)
64495	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with image guidance (fluoroscopy or CT), lumbar or sacral; third and any additional level(s) (List separately in addition to code for primary procedure)
77002	Fluoroscopic guidance for needle placement (eg, biopsy, aspiration, injection, localization device) [should be billed with the integrated code above]
ICD-10 codes covered if criteria are met:	
<i>Code</i>	<i>Description</i>
M53.0 -M53.1	Cervicocranial - cervicobrachial syndrome
M53.81 - M53.83	Other specified dorsopathies [cervical and lumbar regions]

M53.85 - M53.87	
M54.2	Cervicalgia
M54.30 - M5.5	Lumbago and sciatica
M54.5	Low back pain
M54.6	Pain in thoracic spine
M53.84	Other specified dorsopathies, thoracic region
ICD-10 codes <i>not</i> covered:	
<i>Code</i>	<i>Description</i>
M53.3	Sacrococcygeal disorders, not elsewhere classified [coccygodynia]
M54.10 - M54.18	Radiculopathy

<i>Radiofrequency Facet Denervation</i>	
CPT/HCPCS Codes covered if criteria are met:	
<i>Code</i>	<i>Description</i>
64633	Destruction by neurolytic agent, paravertebral facet joint nerve(s), with imaging guidance (fluoroscopy or CT); cervical or thoracic, single facet joint
64634	Destruction by neurolytic agent, paravertebral facet joint nerve(s), with imaging guidance (fluoroscopy or CT); cervical or thoracic, each additional facet joint (List separately in addition to code for primary procedure)
64635	Destruction by neurolytic agent, paravertebral facet joint nerve(s), with imaging guidance (fluoroscopy or CT); lumbar or sacral, single facet joint
64636	Destruction by neurolytic agent, paravertebral facet joint nerve(s), with imaging guidance (fluoroscopy or CT); lumbar or sacral, each additional facet joint (List separately in addition to code for primary procedure)
ICD-10 codes covered if criteria are met:	
<i>Code</i>	<i>Description</i>

M53.0 -M53.1	Cervicocranial - cervicobrachial syndrome
M53.81 - M53.83 M53.85 - M53.87	Other specified dorsopathies [cervical and lumbar regions]
M54.2	Cervicalgia
M54.5	Low back pain
ICD-10 codes <i>not</i> covered:	
<i>Code</i>	<i>Description</i>
M53.3	Sacrococcygeal disorders, not elsewhere classified [coccygodynia]
M54.10 - M54.18	Radiculopathy

CPT/HCPCS codes <i>not</i> covered for indications in this guideline:	
<i>Code</i>	<i>Description</i>
00600 - 00670	Anesthesia for procedures on the spine and spinal cord [includes monitored anesthesia care; when administered with facet joint injections and radiofrequency facet denervation]
27035	Denervation, hip joint, intrapelvic or extrapelvic intra-articular branches of sciatic, femoral, or obturator nerves
0213T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, cervical or thoracic; single level
0214T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, cervical or thoracic; second level (List separately in addition to code for primary procedure)
0215T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, cervical or thoracic; third and any additional level(s) (List separately in addition to code for primary procedure)

0216T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, lumbar or sacral; single level
0217T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, lumbar or sacral; second level (List separately in addition to code for primary procedure)
0218T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, lumbar or sacral; third and any additional level(s) (List separately in addition to code for primary procedure)
64640	Destruction by neurolytic agent; other peripheral nerve or branch
64999	Unlisted procedure, nervous system [when used to bill for pulsed radiofrequency ablation]
72275	Epidurography, radiological supervision and interpretation
76942	Ultrasonic guidance for needle placement (eg, biopsy, aspiration, injection, localization device), imaging supervision and interpretation
77021	Magnetic resonance guidance for needle placement (eg, for biopsy, needle aspiration, injection, or placement of localization device) radiological supervision and interpretation
G0260	Injection procedure for sacroiliac joint; provision of anesthetic, steroid and/or other therapeutic agent, with or without arthrography

References

1. Abejon D. Pulsed radiofrequency in lumbar radicular pain: clinical effects in various etiological groups. *Pain Pract.* MAR-2007; 7(1): 21-6.
2. Agency for Healthcare Research and Quality (AHRQ). Pain management injection therapies for low back pain. 2015 March. Technology Assessment Report ESIB081. Available at: <http://www.cms.gov/Medicare/Coverage/DeterminationProcess/Downloads/id98TA.pdf>. Accessed on July 5, 2017.
3. American Society of Anesthesiologists Task Force on Chronic Pain Management, American Society of Regional Anesthesia and Pain Medicine. Practice guidelines for chronic pain management: An updated report by the American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. *Anesthesiology.* 2010;112(4):810-833.

4. Ayden SM, Gharibo CG, Mehnert M, Stitik TP. The role of radiofrequency ablation for sacroiliac joint pain: a meta-analysis. *PM R*. 2010; 2(9):842-851.
5. Birkenmaier C, Veihelmann C, Trouillier H, et al. Percutaneous cryodenervation of lumbar facet joints: a prospective clinical trial. *Int Orthop*. 2007; 31(4):525-530.
6. Carette S, Marcoux S, Truchon R, Grondin C, Gagnon J, Allard Y, Latulippe M. N Engl J Med. A controlled trial of corticosteroid injections into facet joints for chronic low back pain. 1991;325(14):1002.
7. Cheng J, Pope JE, Dalton JE, et al. Comparative outcomes of cooled versus traditional radiofrequency ablation of the lateral branches for sacroiliac joint point. *Clin J Pain*. 2013; 29(2):132-137.
8. Chou R, Atlas SJ, Stanos SP, Rosenquist RW. Nonsurgical interventional therapies for low back pain. *Spine*. 2009; 34(10):1078-1093.
9. Chou R, Loeser J, Owens D, et al. Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain – an evidence-based clinical practice guideline from the American Pain Society. *Spine*. 2009;34:1066-1077.
10. Chou R. UpToDate. Subacute and chronic low back pain: Nonsurgical interventional treatment. UpToDate.com. Waltham, MA.
11. Christensen S, Sehgal N. What is the correlation between facet joint radiofrequency outcome and response to comparative medial branch blocks? *Pain Physician*. 2016;19:163-172.
12. Cohen S, Raja S. Pathogenesis, diagnosis, and treatment of lumbar zygapophysial (facet) joint pain. *Anesthesiology*. 2007;106:591-614.
13. Cohen SP, Hurley RW, Buckenmaier CC 3rd, et al. Randomized placebo-controlled study evaluating lateral branch radiofrequency denervation for sacroiliac joint pain. *Anesthesiology*. 2008;109(2):279-288.
14. Cohen SP, Raja SN. Pathogenesis, diagnosis and treatment of lumbar zygapophysial (facet) joint pain. *Anesthesiology*. 2007; 106(3):591-614.
15. Cohen SP, Stojanovic MP, Crooks M, et al. Lumbar zygapophysial (facet) joint radiofrequency denervation success as a function of pain relief during diagnostic medial branch blocks: a multicenter analysis. *Spine J*. 2008; 8:498-504.
16. Cohen SP, Strassels SA, Kurihara C, et al. Establishing an optimal "cutoff" threshold for diagnostic lumbar facet blocks: A prospective correlational study. *Clin J Pain*. 2013; 29:382-391.
17. Cohen SP, Strassels SA, Kurihara C, et al. Outcome predictors for sacroiliac joint (lateral branch) radiofrequency denervation. *Reg Anesth Pain Med*. 2009;34(3):206-214.
18. Cohen SP. Sacroiliac joint pain: a comprehensive review of anatomy, diagnosis, and treatment. *Anesth Analg*. 2005; 101(5):1440-1453.
19. Ferrante FM, King LF, Roche EA, et al. Radiofrequency sacroiliac joint denervation for sacroiliac syndrome. *Reg Anesth Pain Med*. 2001; 26(2):137-142.
20. Geurts, J W, van Wijk RM, Stolker RJ, Groen GJ. Efficacy of radiofrequency procedures for the treatment of spinal pain: a systematic review of randomized clinical trials. *Reg Anesth Pain Med*. 2001; 26(5):394-400.

21. Hansen HC, McKenzie-Brown AM, Cohen SP, et al. Sacroiliac joint interventions: A systematic review. *Pain Physician*. 2007;10(1):165-184.
22. Haufe SMW, Mork AR. Endoscopic Facet Debridement for the treatment of facet arthritic pain - a novel new technique. *International Journal of Medical Sciences*. 2010;7(3):120-123.
23. Ho K-Y, Hadi MA, Pasuthamchat K, Tan KH. Cooled radiofrequency denervation for treatment of sacroiliac joint pain: Two-year results from 20 cases. *J Pain Res*. 2013;6:505-511.
24. Jee H, Lee JH, Park KD, et al. Ultrasound-guided versus fluoroscopy-guided sacroiliac joint intra-articular injections in the noninflammatory sacroiliac joint dysfunction: A prospective, randomized, single-blinded study. *Arch Phys Med Rehabil*. 2014;95(2):330-337.
25. Juch JN, Mass ET, Ostelo RW, et al. Effect of radiofrequency denervation on pain intensity among patients with chronic low back pain. The mint randomized clinical trials. *J Amer Med Assoc*. 2017; 318(1):68-81.
26. Kapural L, Nageeb F, Kapural M, et al. Cooled radiofrequency system for the treatment of chronic pain from sacroiliitis: The first case-series. *Pain Pract*. 2008;8(5):348-354.
27. Karaman H, Kavak GO, Tufek A, et al. Cooled radiofrequency application for treatment of sacroiliac joint pain. *Acta Neurochir (Wien)*. 2011;153(7):1461-1468.
28. Kornick C, Kramarich SS, Lamer TJ, et al. Complications of lumbar facet radiofrequency denervation. *Spine*. 2004; 29(12):1352-1354.
29. Kroll HR, Kim D, Danic MJ, et al. A randomized, double-blind, prospective study comparing the efficacy of continuous versus pulsed radiofrequency in the treatment of lumbar facet syndrome. *J Clin Anesth* 2008 Nov;20(7):534-7.
30. Lilius G, Laasonen EM, Myllynen P, Harilainen A, Grönlund G. Lumbar facet joint syndrome. A randomised clinical trial. *J Bone Joint Surg Br*. 1989 Aug;71(4):681-4.
31. Manchikanti KN, Atluri S, Singh V, Geffert S, Sehgal N, Falco FJ. An update of evaluation of therapeutic thoracic facet joint interventions. *Pain Physician* 2012;15(4):E463-81
32. Manchikanti L, Abdi S, Atluri S, et al. An update on comprehensive evidence-based guidelines for interventional techniques in chronic spinal pain. Part II: guidance and recommendations. *Pain Physician*. 2013; 16:S49-283.
33. Manchikanti L, Boswell MV, Singh V, et al. Comprehensive evidence-based guidelines for interventional techniques in the management of chronic spinal pain. *Pain Physician* 2009; 12(4):699-802.
34. Manchikanti L, Singh V, Falco FJ, et al. Comparative effectiveness of a one-year follow-up of thoracic medial branch blocks in management of chronic thoracic pain: a randomized, double-blind active controlled trial. *Pain Physician*. 2010a; 13(6):535-548.
35. Manchikanti L, Singh V, Falco FJ, et al. Comparative outcomes of a 2-year follow-up of cervical medial branch blocks in management of chronic neck pain: a randomized, double-blind controlled trial. *Pain Physician*. 2010b; 13(5):437-450.
36. Manchikanti L, Singh V, Falco FJ, et al. Evaluation of lumbar facet joint nerve blocks in managing chronic low back pain: a randomized, double-blind, controlled trial with a 2-year follow-up. *Int J Med Sci*. 2010c; 7(3):124-135.

37. Mikeladze G, Espinal R, Finnegan R, et al. Pulsed radiofrequency application in treatment of chronic zygapophyseal joint pain. *Spine J.* 2003; 3(5):360-362.
38. Muhlner S. Review article: radiofrequency neurotomy for the treatment of sacroiliac joint syndrome. *Curr Rev Musculoskelet Med.* 2009; 2(1):10–14.
39. Nath S, Nath CA, Pettersson K. Percutaneous lumbar zygapophysial (Facet) joint neurotomy using radiofrequency current in the management of chronic low back pain: a randomized double-blind trial. *Spine (Phila Pa 1976).* 2008; 33(12):1291-1297.
40. Patel N, Gross A, Brown L, Gekht G. A randomized, placebo-controlled study to assess the efficacy of lateral branch neurotomy for chronic sacroiliac joint pain. *Pain Med.* 2012;13(3):383-98.
41. Patel N. Twelve-month follow-up of a randomized trial assessing cooled radiofrequency denervation as a treatment for sacroiliac region pain. *Pain Pract.* 2016; 16(2):154-167.
42. Rambaransingh, B, Stanford G, Burnham R. The effect of repeated zygapophyseal joint radiofrequency neurotomy on pain, disability and improvement duration. *Pain Med.* 2010; 11(9):1343-1347.
43. Ribeiro LH, Furtado RN, Konai MS, Andreo AB, Rosenfeld A, Natour J. Effect of facet joint injection versus systemic steroids in low back pain: a randomized controlled trial. *Spine (Phila Pa 1976).* 2013;38(23):1995.
44. Schofferman J, Kine G. Effectiveness of repeated radiofrequency neurotomy for lumbar facet pain. *Spine (Phila Pa 1976).* 2004; 29(21):2471-2473.
45. Slipman CW, Bhat AL, Gilchrist RV, et al. A critical review of the evidence for the use of zygapophysial injections and radiofrequency denervation in the treatment of low back pain. *Spine J.* 2003; 3(4):310-316.
46. Stelzer W, Aiglesberger M, Stelzer D, Stelzer V. Use of cooled radiofrequency lateral branch neurotomy for the treatment of sacroiliac joint-mediated low back pain: A large case series. *Pain Med.* 2013;14(1):29-35.
47. Tachihara et al. Does facet joint inflammation induce radiculopathy?: an investigation using a rat model of lumbar facet joint inflammation. *Spine (Phila Pa 1976).* 2007 Feb 15;32(4):406-12.
48. UpToDate, Inc. Overview of the treatment of non-cancer chronic pain. <http://www.uptodate.com>. Accessed October 8, 2017
49. UpToDate, Inc. Subacute and chronic low back pain: nonsurgical interventional treatment. <http://www.uptodate.com>. Accessed October 8, 2017
50. UpToDate, Inc. Treatment of neck pain. <http://www.uptodate.com>. Accessed October 8, 2017
51. Vallejo R, Benyamin RM, Kramer J, et al. Pulsed radiofrequency denervation for the treatment of sacroiliac joint syndrome. *Pain Med.* 2006;7(5):429-434
52. Van Boxem K, de Meij N, Kessels A, et al. Pulsed radiofrequency for chronic intractable lumbosacral radicular pain: a six-month cohort study. *Pain Med.* 2015; 16(6):1155-1162.
53. Van Boxem K, van Eerd M, Brinkhuize T, et al. Radiofrequency and pulsed radiofrequency treatment of chronic pain syndromes: the available evidence. *Pain Pract.* 2008; 8(5):385-393.

54. Van Wijk RM, Guerts JW, Wynne HJ, et al. Radiofrequency denervation of lumbar facet joints in the treatment of chronic low back pain: a randomized, double-blind, sham lesion-controlled trial. *Clin J Pain*. 2005; 21(4):335-344.
55. Van Zundert J, Patijn J, Kessels A, et al. Pulsed radiofrequency adjacent to the cervical root ganglion in chronic cervical radicular pain: a double blind sham controlled randomized clinical trial. *Pain*. 2007; 127(1-2):173-182.
56. Iwatsuki K, Yoshimine T, Awazu K. Alternative denervation using laser irradiation in lumbar facet syndrome. *Lasers Surg Med*. 2007; 39(3):225-229.
57. Sehgal N, Shah RV, McKenzie-Brown AM, Everett CR. Diagnostic utility of facet (zygapophysial) joint injections in chronic spinal pain: a systematic review of evidence. *Pain Physician*. 2005;8(2):211–224.
58. Manchikanti L, Staats PS, Singh V, Schultz DM, Vilims BD, Jasper JF, Kloth DS, Trescot AM, Hansen HC, Falasca TD, Racz GB, Deer TR, Burton AW, Helm S, Lou L, Bakhit CE, Dunbar EE, Atluri SL, Calodney AK, Hassenbusch SJ, Feler CA. Evidence-based practice guidelines for interventional techniques in the management of chronic spinal pain. *Pain Physician*. 2003;6(1):3–81.
59. Revel M, Poiraudreau S, Auleley GR, Payan C, Denke A, Nguyen M, Chevrot A, Fermanian J. Capacity of the clinical picture to characterize low back pain relieved by facet joint anesthesia. Proposed criteria to identify patients with painful facet joints. *Spine*. 1998;23(8):1972–1977.
60. Manchikanti L, Boswell MV, Singh V, Pampati V, Damron KS, Beyer CD. Prevalence of facet joint pain in chronic spinal pain of cervical, thoracic, and lumbar regions. *BMC Musculoskelet Disord*. 2004;5:15
61. Atluri S, Dutta S, Falco FJ, Lee M. Systematic review of diagnostic utility and therapeutic effectiveness of thoracic facet joint interventions. *Pain Physician*. 2008;11(5):611–629
62. Scholten PM, Patel SI, Christos PJ, Singh JR. Short-term efficacy of sacroiliac joint corticosteroid injection based on arthrographic contrast patterns. *PM R*. 2015;7(4):385-91.
63. Kennedy DJ, Engel A, Kreiner DS, Nampiaparampil D, Duszynski B, Macvicar J. Fluoroscopically Guided Diagnostic and Therapeutic Intra-Articular Sacroiliac Joint Injections: A Systematic Review. *Pain Med*. 2015;16(8):1500-18.
64. Hansen H, Manchikanti L, Simopoulos TT, et al. A systematic evaluation of the therapeutic effectiveness of sacroiliac joint interventions. *Pain Physician*. 2012;15(3):E247-78.

Clinical Guideline Revision / History Information

Original: Review/Revise Dates	Approval Signature/ Title
Original Date: Reviewed/Revised: Signed:	7/31/2018 Sean Martin, MD, Medical Director