

Kesimpta (ofatumumab)

Disclaimer

Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.

Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.

Kesimpta (ofatumumab)	1
Summary	1
Definitions	2
Clinical Indications	4
Medical Necessity Criteria for Clinical Review	4
General Medical Necessity Criteria	4
Medical Necessity Criteria for Initial Clinical Review	4
Initial Indication-Specific Criteria	4
Multiple Sclerosis - Adults	4
Medical Necessity Criteria for Subsequent Clinical Review	5
Subsequent Indication-Specific Criteria	5
Multiple Sclerosis - Adults	5
Experimental or Investigational / Not Medically Necessary	6
References	6
Clinical Guideline Revision / History Information	9

Summary

Multiple sclerosis (MS) is a chronic, inflammatory, demyelinating disease of the central nervous system. It typically presents in young adults (generally diagnosed before 50 years of age) with symptoms such as vision problems, muscle weakness, numbness, and difficulty with balance and coordination. The most

common form is relapsing-remitting MS (occurring in about 85% of patients), characterized by acute attacks followed by periods of remission. Treatment goals include reducing relapses, slowing disability progression, and managing symptoms. Disease-modifying therapies (DMTs) are the primary treatment approach and include injectable medications (e.g., interferons, glatiramer acetate), oral medications (e.g., dimethyl fumarate, fingolimod, teriflunomide, etc.), and infusion therapies (e.g., natalizumab, ocrelizumab).

MS is a progressive disease, meaning that symptoms tend to worsen over time, and it can be classified into several types, including relapsing-remitting MS (RRMS), primary progressive MS (PPMS), and secondary progressive MS (SPMS).

Currently, there is no cure for MS, but various treatment options are available to manage symptoms, slow the progression of the disease, and improve quality of life.

- Disease-modifying therapies (DMTs) are a class of medications that target the immune system to reduce inflammation and slow down the progression of the disease. The type of DMT prescribed will depend on the type and severity of MS, as well as the individual's medical history and preferences. Some common DMTs include interferon beta, glatiramer acetate, dimethyl fumarate, and fingolimod.
- High dose corticosteroids, such as high dose intravenous methylprednisolone or oral prednisone can be prescribed to reduce inflammation during acute MS relapses.
- Symptomatic treatments are also available to manage specific symptoms of MS, such as muscle spasms, bladder problems, and depression. Physical therapy, occupational therapy, and speech therapy can help individuals with MS maintain mobility, independence, and communication skills.

Kesimpta (ofatumumab) is a recombinant human anti-CD20 monoclonal antibody indicated for treating relapsing forms of MS, to include clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease, in adults. It acts by binding to CD20 on B lymphocytes, causing B-cell depletion through antibody-dependent cellular cytotoxicity and complement-dependent cytotoxicity. Kesimpta (ofatumumab) is administered subcutaneously, offering a self-administered option amongst high-efficacy therapies.

Definitions

"Clinically isolated syndrome" refers to a first episode of neurologic symptoms lasting at least 24 hours caused by inflammation or demyelination in the central nervous system.

"Compendia" are summaries of drug information and medical evidence to support decision-making about the appropriate use of drugs and medical procedures. Examples include, but are not limited to:

1. American Hospital Formulary Service Drug Information

2. Clinical pharmacology
3. National Comprehensive Cancer Network Drugs and Biologics Compendium
4. Thomson Micromedex DrugDex
5. United States Pharmacopeia-National Formulary (USP-NF)

"Disease-modifying therapy" is a medication that modifies the course of MS by reducing relapses and slowing disability progression.

"Documentation" refers to written information, including but not limited to:

- Up-to-date chart notes, relevant test results, and/or relevant imaging reports to support diagnoses; or
- Prescription claims records, and/or prescription receipts to support prior trials of formulary alternatives.

"MRI" or "Magnetic Resonance Imaging" refers to a medical imaging technique that creates detailed three-dimensional (3D) images of the organs and tissues in your body. A brain MRI can reveal areas of active MS disease called lesions within the central nervous system.

"Multiple sclerosis" is a chronic autoimmune disease of the central nervous system characterized by inflammation, demyelination, and neurodegeneration.

"No evidence of" indicates that the reviewer has not identified any records of the specified item or condition within the submitted materials or claims history. In the absence of such evidence, the member is considered eligible. If any evidence of the item or condition is present upon review of the request, the member does not qualify.

"Primary progressive MS" refers to worsening neurologic function from the onset of symptoms, without early relapses or remissions.

"Relapse" is defined as the appearance of new symptoms or the worsening of existing symptoms lasting at least 24 hours in the absence of fever or infection.

"Relapsing-remitting MS" refers to a disease course characterized by clearly defined attacks of new or increasing neurologic symptoms followed by periods of partial or complete recovery.

"[s]" indicates state mandates may apply.

"Secondary progressive MS" is a disease course following relapsing-remitting MS that is characterized by a progressive worsening of neurologic function over time with or without relapses.

Clinical Indications

Medical Necessity Criteria for Clinical Review

General Medical Necessity Criteria

The Plan considers Kesimpta (ofatumumab) medically necessary when ONE of the following criteria are met:

1. Authorization may be granted for pediatric members less than 18 years of age with multiple sclerosis when there is documentation that the benefits outweigh the risks; *OR*
Note: If approved, the requested product will be authorized for up until the member reaches 18 years of age.
2. The member meets the applicable [Medical Necessity Criteria for Initial Clinical Review](#) or [Subsequent Clinical Review](#) listed below.

Medical Necessity Criteria for Initial Clinical Review

Initial Indication-Specific Criteria

Multiple Sclerosis - Adults

The Plan considers Kesimpta (ofatumumab) medically necessary when recent (within the last 3 months) clinical chart documentation provided indicates the member meets ALL of the following:

1. Prescribed by or in consultation with a neurologist or physician who specializes in the treatment of multiple sclerosis; *AND*
2. The member is 18 years of age or older; *AND*
3. The member has ONE (1) of the following forms of multiple sclerosis:
 - a. Relapsing-remitting (RRMS); *or*
 - b. Active secondary progressive disease (SPMS); *or*
 - c. Clinically isolated syndrome (CIS); *AND*
4. The member meets ONE (1) of the following:
 - a. Documentation of highly active or aggressive disease, as demonstrated by at least ONE (1) of the following:
 - i. Frequent relapses (≥ 2 in the past year); *or*
 - ii. At least 1 relapse with incomplete recovery and MRI activity; *or*
 - iii. Rapidly advancing disability or cognitive impairment; *or*
 - iv. Disabling relapse with suboptimal response to corticosteroids; *or*
 - v. MRI findings showing high disease activity (e.g., new/enlarging T2 lesions, enhancing lesions); *or*
 - b. The member is unable to use, or has tried and failed ONE (1) of the following:^[a]
 - i. Dimethyl Fumarate (generic Tecfidera); *or*
 - ii. Fingolimod (generic Gilenya); *or*
 - iii. Teriflunomide (generic Aubagio); *AND*

5. The member has been screened for hepatitis B virus AND has no evidence of active hepatitis B infection; *AND*
6. Kesimpta (ofatumumab) will be used as monotherapy for multiple sclerosis (i.e., member is not using and will not use other disease-modifying MS therapies while on Kesimpta [ofatumumab]); *AND*
7. Kesimpta (ofatumumab) is being prescribed at a dose and frequency that is within FDA approved labeling OR is supported by compendia or evidence-based published dosing guidelines for the requested indication.
 - o *20 mg/0.4 mL solution in a single-dose prefilled Sensoready pen or prefilled syringe.*
 - i. *Initial dosing: 20 mg at weeks 0, 1, and 2.*
 - ii. *Maintenance dosing: 1 dose (20 mg) every 4 weeks starting at week 4.*

If the above prior authorization criteria are met, the requested medication will be approved for up to 12-months.^[a]

Continued Care

Medical Necessity Criteria for Subsequent Clinical Review

Subsequent Indication-Specific Criteria

Multiple Sclerosis - Adults

The Plan considers Kesimpta (ofatumumab) medically necessary when recent (within the last 6-months) clinical chart documentation provided indicates the member meets BOTH of the following:

1. The requested medication is prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis; *AND*
2. The member has experienced at least ONE (1) of the following:
 - a. Improvement in at least ONE (1) objective measure, such as:
 - i. Reduced disease activity on MRI; *and/or*
 - ii. Improved or stable disability scores; *and/or*
 - iii. Reduced relapse rate; *and/or*
 - iv. Improved fatigue or walking assessments; *AND/OR*
 - b. The member has shown stabilization or improvement in at least ONE (1) MS symptom, such as:
 - i. Motor function; *and/or*
 - ii. Fatigue; *and/or*
 - iii. Vision; *and/or*
 - iv. Bowel/bladder function; *and/or*
 - v. Spasticity; *and/or*
 - vi. Walking/gait; *and/or*
 - vii. Pain/numbness/tingling.

If the above reauthorization criteria are met, the requested product will be authorized for up to 12-months.^[5]

Experimental or Investigational / Not Medically Necessary^[5]

Kesimpta (ofatumumab) for any other indication or use is considered not medically necessary by the Plan, as it is deemed to be experimental, investigational, unproven, or not medically necessary.

Non-covered indications include, but are not limited to, the following:

- Use in combination with other disease-modifying therapies for MS. There is limited knowledge about the use of combining DMTs for MS.
- Treatment of non-relapsing forms of multiple sclerosis (e.g., primary progressive MS). The safety and efficacy of Kesimpta (ofatumumab) has not been established in those with PPMS.
- Treatment of other autoimmune conditions not FDA-approved (e.g., rheumatoid arthritis, systemic lupus erythematosus). While small controlled and open-label studies have shown some promise in improving the American College of Rheumatology 20% criteria for Improvement (ACR20), larger high quality studies have not yet been conducted. No high quality studies have supported the use of Kesimpta (ofatumumab) for systemic lupus erythematosus or other autoimmune conditions.
- Use for the treatment of nephrotic syndrome. There are no high quality studies to support the safety and efficacy of Kesimpta (ofatumumab) for the management of nephrotic syndromes. Several studies of low quality or small sample size show non-superiority over an alternative anti-CD20 monoclonal antibody (rituximab) or no benefit. Further studies are needed to support this indication.

References

1. Bainbridge JL, Miravalle A, Wong PS. Multiple Sclerosis. In DiPiro JT, Yee GC, Posey LM, et al, eds. *Pharmacotherapy: A Pathophysiologic Approach*. 11th ed. New York, NY: McGraw-Hill; 2019.
2. Bar-Or A, Grove RA, Austin DJ, et al,. Subcutaneous ofatumumab in patients with relapsing-remitting multiple sclerosis: The MIRROR study. *Neurology*. 2018 May 15;90(20):e1805-e1814. doi: 10.1212/WNL.0000000000005516. Epub 2018 Apr 25. Erratum in: *Neurology*. 2018 Sep 11;91(11):538. doi: 10.1212/WNL.0000000000005929.
3. Benallegue N, Rollot F, Wiertlewski S, Casey R, Debouverie M, Kerbrat A, De Seze J, Ciron J, Ruet A, Labauge P, Maillart E, Zephir H, Papeix C, Defer G, Lebrun-Frenay C, Moreau T, Berger E, Stankoff B, Clavelou P, Heinzlef O, Pelletier J, Thouvenot E, Al Khedr A, Bourre B, Casez O, Cabre P, Wahab A, Magy L, Vukusic S, Laplaud DA; OFSEP (Observatoire Français de la Sclérose en Plaques) Investigators. Highly Effective Therapies as First-Line Treatment for Pediatric-Onset Multiple Sclerosis. *JAMA Neurol*. 2024 Mar 1;81(3):273-282.
4. Bonanni A, Rossi R, Murtas C, Ghiggeri GM. Low-dose ofatumumab for rituximab-resistant nephrotic syndrome. *BMJ Case Rep*. 2015 Sep 16;2015:bcr2015210208. doi: 10.1136/bcr-2015-210208.
5. Burt RK, Balabanov R, Burman J, Sharrack B, Snowden JA, Oliveira MC, Fagius J, Rose J, Nelson F, Barreira AA, Carlson K, Han X, Moraes D, Morgan A, Quigley K, Yaung K, Buckley R, Alldredge C, Clendenan A, Calvario MA, Henry J, Jovanovic B, Helenowski IB. Effect of Nonmyeloablative

Hematopoietic Stem Cell Transplantation vs Continued Disease-Modifying Therapy on Disease Progression in Patients With Relapsing-Remitting Multiple Sclerosis: A Randomized Clinical Trial. *JAMA*. 2019 Jan 15;321(2):165-174. doi: 10.1001/jama.2018.18743. PMID: 30644983; PMCID: PMC6439765.

6. Chitnis T, Tenenbaum S, Banwell B, Krupp L, Pohl D, Rostasy K, Yeh EA, Bykova O, Wassmer E, Tardieu M, Kornberg A, Ghezzi A; International Pediatric Multiple Sclerosis Study Group. Consensus statement: evaluation of new and existing therapeutics for pediatric multiple sclerosis. *Mult Scler*. 2012 Jan;18(1):116-27.
7. Freedman MS, Devonshire V, Duquette P, Giacomini PS, Giuliani F, Levin MC, Montalban X, Morrow SA, Oh J, Rotstein D, Yeh EA; Canadian MS Working Group. Treatment Optimization in Multiple Sclerosis: Canadian MS Working Group Recommendations. *Can J Neurol Sci*. 2020 Jul;47(4):437-455. doi: 10.1017/cjn.2020.66. Epub 2020 Apr 6. PMID: 32654681.
8. Gärtner J, Hauser SL, Bar-Or A, et al. Efficacy and safety of ofatumumab in recently diagnosed, treatment-naive patients with multiple sclerosis: Results from ASCLEPIOS I and II. *Mult Scler*. 2022 Sep;28(10):1562-1575. doi: 10.1177/13524585221078825. Epub 2022 Mar 10.
9. Hua LH, Brown B, Camacho E, et al. Switch from intravenous anti-CD20 therapy to subcutaneous ofatumumab in patients with relapsing MS: results from the OLİKOS study. *J Neurol*. 2025 Oct 24;272(11):725. doi: 10.1007/s00415-025-13462-w.
10. Hauser SL, Bar-Or A, Cohen JA, et al. Ofatumumab versus Teriflunomide in Multiple Sclerosis. *N Engl J Med*. 2020 Aug 6;383(6):546-557. doi: 10.1056/NEJMoa1917246.
11. Hauser, S., & Cree, B. (2020). Treatment of Multiple Sclerosis: A Review.. *The American journal of medicine*. <https://doi.org/10.1016/j.amjmed.2020.05.049>.
12. Hauser SL, Freedman MS, Nakahara J, et al. Six-year safety and efficacy outcomes with first-line ofatumumab in recently diagnosed treatment-naive patients with relapsing multiple sclerosis. *Mult Scler Relat Disord*. 2026 Jan;105:106886. doi: 10.1016/j.msard.2025.106886. Epub 2025 Nov 25.
13. Hauser SL, Zielman R, Das Gupta A, et al. Efficacy and safety of four-year ofatumumab treatment in relapsing multiple sclerosis: The ALITHIOS open-label extension. *Mult Scler*. 2023 Oct;29(11-12):1452-1464. doi: 10.1177/13524585231195346. Epub 2023 Sep 11.
14. He A, Merkel B, Brown JW, et al. Timing of high-efficacy therapy for multiple sclerosis: a retrospective observational cohort study. *Lancet Neurol*. 2020 Apr;19(4):307-316. doi: 10.1016/S1474-4422(20)30067-3. Epub 2020 Mar 18.
15. Kesimpta (ofatumumab) [prescribing information]. East Hanover, NJ: Novartis Pharmaceuticals; August 2025.
16. Köhler M, Paul F, Janke K, et al. Comparative effectiveness of disease-modifying therapies for highly active relapsing-remitting multiple sclerosis despite previous treatment - a systematic review and network meta-analysis. *BMC Neurol*. 2025 Aug 9;25(1):328. doi: 10.1186/s12883-025-04338-7.
17. Krysko KM, Graves JS, Rensel M, et al; US Network of Pediatric MS Centers. Real-World Effectiveness of Initial Disease-Modifying Therapies in Pediatric Multiple Sclerosis. *Ann Neurol*. 2020 Jul;88(1):42-55.
18. Krysko KM, Graves J, Rensel M, et al; US Network of Pediatric MS Centers. Use of newer disease-modifying therapies in pediatric multiple sclerosis in the US. *Neurology*. 2018 Nov 6;91(19):e1778-e1787.
19. Kurrasch R, Brown JC, Chu M, et al. Subcutaneously administered ofatumumab in rheumatoid arthritis: a phase I/II study of safety, tolerability, pharmacokinetics, and pharmacodynamics. *J Rheumatol*. 2013 Jul;40(7):1089-96. doi: 10.3899/jrheum.121118. Epub 2013 Jun 1.
20. Larkins NG, Hahn D, Liu ID, Willis NS, Craig JC, Hodson EM. Non-corticosteroid immunosuppressive medications for steroid-sensitive nephrotic syndrome in children. *Cochrane Database Syst Rev*. 2024 Nov 8;11(11):CD002290. doi: 10.1002/14651858.CD002290.pub6.
21. Liu ID, Willis NS, Craig JC, Hodson EM. Interventions for idiopathic steroid-resistant nephrotic syndrome in children. *Cochrane Database Syst Rev*. 2025 May 8;5(5):CD003594. doi: 10.1002/14651858.CD003594.pub7.

22. McGinley MP, Goldschmidt CH, Rae-Grant AD. Diagnosis and Treatment of Multiple Sclerosis: A Review. *JAMA*. 2021;325(8):765–779. doi:10.1001/jama.2020.26858
23. Mohammadi I, Rajai Firouzabadi S, Aghajanian S, et al. Efficacy of alternative vs. standard dosing strategies of anti-CD20 monoclonal antibodies in multiple sclerosis: A systematic review and meta-analysis. *Mult Scler Relat Disord*. 2025 Nov;103:106668. doi: 10.1016/j.msard.2025.106668. Epub 2025 Aug 8.
24. Montalban X, Leist TP, Cohen BA et al. Cladribine tablets added to IFN- β in active relapsing MS: The ONWARD study. *Neurol Neuroimmunol Neuroinflamm*. 2018; 5:e477.
25. Montalban X, Lebrun-Frénay C, Oh J, et al. Diagnosis of multiple sclerosis: 2024 revisions of the McDonald criteria. *Lancet Neurol*. 2025 Oct;24(10):850-865. doi: 10.1016/S1474-4422(25)00270-4. Erratum in: *Lancet Neurol*. 2025 Nov;24(11):e13. doi: 10.1016/S1474-4422(25)00355-2.
26. Multiple Sclerosis Society of Canada. Disease-modifying therapies. <https://mssociety.ca/managing-ms/treatments/medications/disease-modifying-therapies-dmts>.
27. National Institute for Health and Care Excellence [NICE]. Multiple sclerosis in adults: management. NICE Guidelines [NG220]. 22 June 2022. Available at: <https://www.nice.org.uk/guidance/ng220/chapter/Recommendations#ms-symptom-management-and-rehabilitation>. Accessed 20 January 2026.
28. National MS Society. Disease-modifying therapies for MS (updated March 2022). Available from National MS Society website: <https://nms2cdn.azureedge.net/cmssite/nationalmssociety/media/msnationalfiles/brochures/brochure-the-ms-disease-modifying-medications.pdf>.
29. Østergaard M, Baslund B, Rigby W, et al. Ofatumumab, a human anti-CD20 monoclonal antibody, for treatment of rheumatoid arthritis with an inadequate response to one or more disease-modifying antirheumatic drugs: results of a randomized, double-blind, placebo-controlled, phase I/II study. *Arthritis Rheum*. 2010 Aug;62(8):2227-38. doi: 10.1002/art.27524.
30. Quattrocchi E, Østergaard M, Taylor PC, et al. Safety of repeated open-label treatment courses of intravenous ofatumumab, a human anti-CD20 monoclonal antibody, in rheumatoid arthritis: results from three clinical trials. *PLoS One*. 2016;11(6):e0157961. doi:10.1371/journal.pone.0157961.
31. Rae-Grant A, Day GS, Marrie RA, et al. Practice guideline recommendations summary: Disease-modifying therapies for adults with multiple sclerosis: Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology. *Neurology*. 2018;90(17):777-788.
32. Rashid W, Ciccarelli O, Leary SM, et al. Using disease-modifying treatments in multiple sclerosis: Association of British Neurologists (ABN) 2024 guidance. *Pract Neurol*. 2025 Jan 16;25(1):18-24. doi: 10.1136/pn-2024-004228.
33. Ravani P, Colucci M, Bruschi M, et al. Human or Chimeric Monoclonal Anti-CD20 Antibodies for Children with Nephrotic Syndrome: A Superiority Randomized Trial. *J Am Soc Nephrol*. 2021 Oct;32(10):2652-2663. doi: 10.1681/ASN.2021040561. Epub 2021 Sep 20.
34. Ravani P, Pisani I, Bodria M, Caridi G, Degl'Innocenti ML, Ghiggeri GM. Low-dose ofatumumab for multidrug-resistant nephrotic syndrome in children: a randomized placebo-controlled trial. *Pediatr Nephrol*. 2020 Jun;35(6):997-1003. doi: 10.1007/s00467-020-04481-y. Epub 2020 Jan 28.
35. Reich DS, Lucchinetti CF, Calabresi PA. 2018. Multiple sclerosis. *New England Journal of Medicine* 378(2):169-180.
36. Samjoo IA, Drudge C, Walsh S, et al.,. Comparative efficacy of therapies for relapsing multiple sclerosis: a systematic review and network meta-analysis. *J Comp Eff Res*. 2023 Jul;12(7):e230016. doi: 10.57264/cer-2023-0016. Epub 2023 Jun 2.
37. Śladowska K, Kawalec P, Holko P, Osiecka O. Comparative safety of high-efficacy disease-modifying therapies in relapsing-remitting multiple sclerosis: a systematic review and network meta-analysis. *Neurol Sci*. 2022 Sep;43(9):5479-5500. doi: 10.1007/s10072-022-06197-3. Epub 2022 Jun 17.

38. Sladowska K, Mocko P, Brzostek T, et al. Efficacy and safety of disease-modifying therapies in pediatric-onset multiple sclerosis: A systematic review of clinical trials and observational studies. *Mult Scler Relat Disord*. 2025. doi: 10.1016/j.msard.2025.106263
39. Taylor PC, Quattrocchi E, Mallett S, et al. Ofatumumab, a fully human anti-CD20 monoclonal antibody, in biological-naive, rheumatoid arthritis patients with an inadequate response to methotrexate: a randomised, double-blind, placebo-controlled clinical trial. *Ann Rheum Dis*. 2011 Dec;70(12):2119-25. doi: 10.1136/ard.2011.151522. Epub 2011 Aug 22.
40. The use of disease-modifying therapies in multiple sclerosis: principles and current evidence summary. Multiple Sclerosis Coalition. Available from the National MS Society Website: <https://www.nationalmssociety.org/>.
41. Tramacere I, Del Giovane C, Salanti G, et al. Immunomodulators and immunosuppressants for relapsing-remitting multiple sclerosis: a network meta-analysis. *Cochrane Database Syst Rev* 2015;9:CD011381.
42. Walsh R, Chitnis T. Therapeutic Advances in Pediatric Multiple Sclerosis. *Children*. 2025;12(3):259.
43. Wu X, Tan X, Zhang J, et al,. The Efficacy and Safety of Anti-CD20 Antibody Treatments in Relapsing Multiple Sclerosis: A Systematic Review and Network Meta-analysis. *CNS Drugs*. 2022 Nov;36(11):1155-1170. doi: 10.1007/s40263-022-00961-x. Epub 2022 Oct 16.
44. Yang, J., Rempe, T., Whitmire, N., Dunn-Pirio, A., & Graves, J. (2022). Therapeutic Advances in Multiple Sclerosis. *Frontiers in Neurology*, 13. <https://doi.org/10.3389/fneur.2022.824926>.

Clinical Guideline Revision / History Information

Original Date: 06/27/2024

Reviewed/Revised: 8/29/2024, 10/01/2025, 05/01/2026