

## Aripiprazole oral disintegrating tablet, solution

- Aripiprazole oral disintegrating tablet (Abilify Discmelt)
- Aripiprazole oral solution (Abilify 1mg/mL Solution)
- Aripiprazole oral film (Mezofy)
- Aripiprazole oral film (OPIPZA)

### Disclaimer

*Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.*

*Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.*

Aripiprazole oral disintegrating tablet, solution	1
Summary	2
Definitions	2
Medical Necessity Criteria for Clinical Review	4
General Medical Necessity Criteria	4
Medical Necessity Criteria for Initial Clinical Review	4
Initial Indication-Specific Criteria	4
Adjunctive Therapy of Major Depressive Disorder	4
Bipolar Disorder	4
Irritability Associated with Autistic Disorder	4
Postpartum Psychosis:	4
Schizophrenia	5
Tourette's Syndrome or chronic tic disorders	5

For other off-label uses	5
Experimental or Investigational / Not Medically Necessary	6
References	6
Clinical Guideline Revision / History Information	8

## Summary

Aripiprazole (Abilify, Mezofy) is an atypical antipsychotic medication (second generation antipsychotic) classified as a dopamine system stabilizer. Unlike most antipsychotics, which act as full dopamine antagonists, aripiprazole is a partial dopamine agonist and exhibits affinity to serotonin (5-HT) receptors 5-HT1A and 5-HT2A. This unique mechanism allows aripiprazole to reduce positive symptoms of schizophrenia and stabilize dopamine activity without causing significant extrapyramidal side effects.

- Abilify (aripiprazole) Oral Tablets, Orally-Disintegrating Tablets, and Oral Solution are indicated for the treatment of:
  - Schizophrenia.
  - Acute Treatment of Manic and Mixed Episodes associated with Bipolar I Disorder.
  - Adjunctive Treatment of Major Depressive Disorder.
  - Irritability Associated with Autistic Disorder.
  - Treatment of Tourette's Disorder.
- Mezofy (aripiprazole) oral film is indicated for the treatment of:
  - Schizophrenia in those 13 years of age and older.
- In addition to oral tablets, aripiprazole is available in several long-acting injectable formulations, including monthly, every 2 months, and every 1-2 month extended-release intramuscular injections for maintenance therapy of schizophrenia and bipolar I disorder.
- An ingestible digital sensor tablet was also approved to monitor adherence to oral aripiprazole treatment.

The major safety concerns with aripiprazole include increased mortality in older adult with dementia and increased suicidal ideation in pediatric and young adult patients, warranting boxed warnings for these populations. Like other second-generation antipsychotics or atypical antipsychotics, aripiprazole (Abilify, Mezofy), can cause metabolic syndrome including high cholesterol (hyperlipidemia, hypercholesterolemia), high blood sugar (hyperglycemia) and weight gain.

## Definitions

“Adjunctive therapy” is the use of a medication in combination with another medication, often to enhance the main therapy or treat additional symptoms. Aripiprazole is approved as adjunctive therapy for major depressive disorder.

"Atypical Antipsychotics" are a group of antipsychotic drugs used to treat psychiatric conditions. Atypical antipsychotics are less likely to produce extrapyramidal side effects (movement disorders) compared to the older typical antipsychotics.

"Bipolar disorder" is a mental health condition causing unusual shifts in mood, energy, and activity levels ranging from depressive lows to manic highs.

"Boxed Warnings", also known as a black box warning, is the most serious type of warning issued by the Food and Drug Administration (FDA). It is placed on the labeling of prescription drugs that have serious or life-threatening risks.

"Documentation" refers to written information, including but not limited to:

- Up-to-date chart notes, relevant test results, and/or relevant imaging reports to support diagnoses; or
- Prescription claims records, and/or prescription receipts to support prior trials of formulary alternatives.

"Irritability associated with autistic disorder" refers to irritability manifesting as aggression, deliberate self-injurious behavior, temper tantrums or quickly changing moods that is associated with autism spectrum disorder.

"Major depressive disorder" is a psychiatric disorder characterized by a persistent feeling of sadness, loss of interest in activities, and other debilitating symptoms.

"Postpartum Psychosis" is a rare but serious mental health emergency characterized by sudden onset of psychotic symptoms shortly after childbirth. It is often characterized by acute onset of delusions, disorganized thoughts, hallucinations, and/or agitation. While it may occur as a manifestation of bipolar disorder, it can also present in patients without prior psychiatric history. Early recognition and treatment are critical due to risks to both mother and infant.

"Schizophrenia" is a chronic and severe mental disorder characterized by disruptions in thinking, perception, emotions, and behavior. Hallucinations and delusions are common.

"Tourette's syndrome" is a neurological disorder characterized by sudden, repetitive, rapid, and unwanted movements or vocal sounds called tics.

"[s]" indicates state mandates may apply.

## Medical Necessity Criteria for Clinical Review

### General Medical Necessity Criteria

The Plan considers aripiprazole (Abilify) oral disintegrating tablet or solution, or Mezofy/Opipza (aripiprazole oral film) medically necessary when ALL of the following criteria are met:

1. Documentation indicating the member's inability or unwillingness to take the tablet form<sup>[s]</sup>; *AND*
2. IF the request is for Mezofy/Opipza (aripiprazole oral film), the member is unable to use, or has tried and failed aripiprazole tbdp/ODT<sup>[s]</sup>; *AND*
3. Aripiprazole (oral disintegrating tablet, solution, film) is being prescribed at a dose and frequency (i.e., once daily, up to a maximum daily dose of 30mg/day) that is within FDA approved labeling OR is supported by compendia or evidence-based published dosing guidelines for the requested indication; *AND*
4. The member meets the applicable indication-specific criteria listed below:

## Medical Necessity Criteria for Initial Clinical Review

### Initial Indication-Specific Criteria

#### Adjunctive Therapy of Major Depressive Disorder

4. The request is for aripiprazole (Abilify) oral disintegrating tablet or solution or Opipza (aripiprazole oral film); *AND*
5. The member is 18 years of age or older; *AND*
6. The member has a diagnosis of major depressive disorder.

#### Bipolar Disorder

4. The request is for aripiprazole (Abilify) oral disintegrating tablet or solution; *AND*
5. The member is 10 years of age or older; *AND*
6. The member has a diagnosis of bipolar disorder.

#### Irritability Associated with Autistic Disorder

4. The request is for aripiprazole (Abilify) oral disintegrating tablet or solution or Opipza (aripiprazole oral film); *AND*
5. The member is between ages 6 to 17 years; *AND*
6. The member has a diagnosis of irritability associated with autistic disorder (including symptoms of aggression, deliberate self-injurious behavior, temper tantrums, quickly changing moods).

#### Postpartum Psychosis:

4. The request is for aripiprazole (Abilify) oral disintegrating tablet or solution; *AND*
5. The requested medication is prescribed by or in consultation with a psychiatrist; *AND*
6. The member has postpartum psychosis as evidenced by ONE of the following:

- a. New onset psychotic symptoms (e.g., delusions, disorganized thoughts, hallucinations, agitation, bizarre behavior) in the postpartum period; *or*
- b. Bipolar disorder with postpartum psychotic symptoms; *or*
- c. Primary mental disorder with psychotic symptoms during the peripartum period.

#### Schizophrenia

4. The request is for aripiprazole (Abilify) oral disintegrating tablet or solution or Mezofy/Opipza (aripiprazole oral film); *AND*
5. The member is 13 years of age or older; *AND*
6. The member has a diagnosis of schizophrenia.

#### Tourette's Syndrome or chronic tic disorders

4. The request is for aripiprazole (Abilify) oral disintegrating tablet or solution or Opipza (aripiprazole oral film); *AND*
5. The member is 6 years of age or older; *AND*
6. The member has a diagnosis of Tourette disorder or chronic tic disorders.

#### For other off-label uses

4. The request is for aripiprazole (Abilify) oral disintegrating tablet or solution or Mezofy/Opipza (aripiprazole oral film); *AND*
5. The member is 18 years of age or older; *AND*
6. The medication is being requested for treatment of **ONE** of the following:
  - a. Agitation or aggression associated with psychiatric conditions (eg, schizophrenia, bipolar disorder), substance intoxication, or other organic causes; *or*
  - b. Agitation/aggression and psychosis related to dementia whose symptoms are dangerous, severe, or cause significant patient distress (i.e., severe behavioral or psychological symptoms of dementia); *or*
  - c. Delusion disorder; *or*
  - d. Huntington disease-associated chorea; *or*
  - e. Delusional infestation (also called delusional parasitosis) *AND* the member is unable to use, or has tried and failed olanzapine and risperidone<sup>[§]</sup>; *or*
  - f. Treatment-resistant obsessive-compulsive disorder (OCD) in members who have a partial response to initial treatment (e.g., antidepressants such as clomipramine, fluvoxamine, fluoxetine, paroxetine, sertraline)<sup>[§]</sup>; *or*

If the above prior authorization criteria are met, the requested product will be authorized for up to a lifetime.

## Experimental or Investigational / Not Medically Necessary<sup>[s]</sup>

Aripiprazole (oral disintegrating tablet, solution) for any other indication or use is considered not medically necessary by the Plan, as it is deemed to be experimental, investigational, or unproven.

Non-covered indications include, but are not limited to, the following:

- Attention Deficit Hyperactivity Disorder (ADHD). While several small, open-label studies have shown promising results in children and adolescents with ADHD (with or without a bipolar diagnosis), no large, controlled, randomized studies have demonstrated safety and efficacy for the management of ADHD with aripiprazole.
- Chronic Lower Back Pain (CLBP). There are no high quality published studies supporting the safety and efficacy of aripiprazole for the management of chronic lower back pain.
- Eating Disorders, including Anorexia Nervosa (AN). There are no high quality published studies supporting the safety and efficacy of aripiprazole for the management of eating disorders, including AN. In a small (n=22) unpublished study (NCT00685334), aripiprazole was compared to olanzapine in those with AN, and found to result in weight loss (versus weight gain with olanzapine) at 12-weeks.
- Generalized Anxiety Disorder (GAD)/Panic Disorder. While several small, open-label studies have shown some benefit with adjunct aripiprazole in those with anxiety or co-morbid anxiety, no large, controlled, randomized studies have demonstrated safety and efficacy for the management of GAD or panic disorder with aripiprazole. Postpartum Depression. There are no high quality published studies supporting the safety and efficacy of aripiprazole for the management of postpartum depression. Of note, aripiprazole may impact human milk product, and thus may not be ideal in breastfeeding women.

## References

1. Abilify (aripiprazole) [prescribing information]. Rockville, MD: Otsuka America Pharmaceutical Inc; November 2022.
2. Ahmed A, Affleck AG, Angus J, et al; British Association of Dermatologists' Clinical Standards Unit. British Association of Dermatologists guidelines for the management of adults with delusional infestation 2022. *Br J Dermatol.* 2022;187(4):472-480. doi:10.1111/bjd.21668
3. Ak M, Bulut SD, Bozkurt A, Ozsahin A. Aripiprazole augmentation of serotonin reuptake inhibitors in treatment-resistant obsessive-compulsive disorder: a 10-week open-label study. *Adv Ther.* 2011;28(4):341-348. doi:10.1007/s12325-011-0011-7
4. Aripiprazole solution [prescribing information]. Weston, FL: Apotex; November 2016.
5. Aripiprazole orally disintegrating tablets [prescribing information]. Cranford, NJ: Viona Pharmaceuticals; July 2022.
6. Arnold MJ. Management of Bipolar Disorder: Guidelines From the VA/DoD. *Am Fam Physician.* 2024 Jun;109(6):585-587. PMID: 38905567.
7. Arnold MJ. Management of First-Episode Psychosis and Schizophrenia: Guidelines From the VA/DoD. *Am Fam Physician.* 2024 May;109(5):482-483.
8. By the 2023 American Geriatrics Society Beers Criteria® Update Expert Panel. American Geriatrics Society 2023 updated AGS Beers Criteria® for potentially inappropriate medication use in older adults. *J Am Geriatr Soc.* 2023 Jul;71(7):2052-2081. doi: 10.1111/jgs.18372. Epub 2023 May 4. PMID: 37139824.
9. Ciammola A, Sassone J, Colciago C, et al. Aripiprazole in the treatment of Huntington's disease: a case series. *Neuropsychiatr Dis Treat.* 2009;5:1-4.

10. Cox JH, Cavanna AE. Aripiprazole for the treatment of Tourette syndrome. *Expert Rev Neurother.* 2021;21(4):381-391. doi:10.1080/14737175.2021.1893693.
11. Furukawa Y, Hamza T, Cipriani A, Furukawa TA, Salanti G, Ostinelli EG. Optimal dose of aripiprazole for augmentation therapy of antidepressant-refractory depression: preliminary findings based on a systematic review and dose-effect meta-analysis. *Br J Psychiatry.* 2022 Aug;221(2):440-447. doi: 10.1192/bjp.2021.165.
12. Garg A, Sinha P, Jyrwa S. Aripiprazole in difficult-to-treat delusional disorder with co-morbid epilepsy. *Psychiatry Clin Neurosci.* 2014;68(9):721-722. doi:10.1111/pcn.12200
13. Garriga M, Pacchiarotti I, Kasper S, et al. Assessment and management of agitation in psychiatry: expert consensus. *World J Biol Psychiatry.* 2016;17(2):86-128. doi:10.3109/15622975.2015.1132007
14. Guideline Development Panel for the Treatment of Depressive Disorders. Summary of the clinical practice guideline for the treatment of depression across three age cohorts. *Am Psychol.* 2022 Sep;77(6):770-780. doi: 10.1037/amp0000904. Epub 2021 Nov 29. PMID: 34843274.
15. Heller MM, Wong JW, Lee ES, et al. Delusional infestations: clinical presentation, diagnosis and treatment. *Int J Dermatol.* 2013;52(7):775-783. doi:10.1111/ijd.12067
16. Keepers GA, Fochtmann LJ, Anzia JM, Benjamin S, Lyness JM, Mojtabai R, Servis M, Walaszek A, Buckley P, Lenzenweger MF, Young AS, Degenhardt A, Hong SH; (Systematic Review). The American Psychiatric Association Practice Guideline for the Treatment of Patients With Schizophrenia. *Am J Psychiatry.* 2020 Sep 1;177(9):868-872. doi: 10.1176/appi.ajp.2020.177901. PMID: 32867516.
17. McQuaid JR, Buelt A, Capaldi V, et al. The Management of Major Depressive Disorder: Synopsis of the 2022 U.S. Department of Veterans Affairs and U.S. Department of Defense Clinical Practice Guideline. *Ann Intern Med.* 2022 Oct;175(10):1440-1451. doi: 10.7326/M22-1603. Epub 2022 Sep 20. PMID: 36122380.
18. Mezofy (aripiprazole) oral film [prescribing information]. Hamburg, Germany: Labtec Company GmbH. April 2025.
19. Mintzer JE, Tune LE, Breder CD, et al. Aripiprazole for the treatment of psychoses in institutionalized patients with Alzheimer dementia: a multicenter, randomized, double-blind, placebo-controlled assessment of three fixed doses. *Am J Geriatr Psychiatry.* 2007;15(11):918-931.
20. Miyamoto S, Miyake N, Ogino S, Endo T, Yamaguchi N. Successful treatment of delusional disorder with low-dose aripiprazole. *Psychiatry Clin Neurosci.* 2008;62(3):369. doi:10.1111/j.1440-1819.2008.01812.
21. Muscatello MR, Bruno A, Pandolfo G, et al. Effect of aripiprazole augmentation of serotonin reuptake inhibitors or clomipramine in treatment-resistant obsessive-compulsive disorder: a double-blind, placebo-controlled study. *J Clin Psychopharmacol.* 2011;31(2):174-179. doi:10.1097/JCP.0b013e31820e3db6.
22. Opipza (aripiprazole) oral film [prescribing information]. Hazlet, NJ: Carwin Pharmaceutical Associates, LLC. March 2025.
23. Qaseem A, Owens DK, Etxeandia-Ikobaltzeta I, Tuftte J, Cross JT Jr, Wilt TJ; Clinical Guidelines Committee of the American College of Physicians; Crandall CJ, Balk E, Cooney TG, Fitterman N, Hicks LA, Lin JS, Maroto M, Obley AJ, Tice JA, Yost J. Nonpharmacologic and Pharmacologic Treatments of Adults in the Acute Phase of Major Depressive Disorder: A Living Clinical Guideline From the American College of Physicians. *Ann Intern Med.* 2023 Feb;176(2):239-252. doi: 10.7326/M22-2056. Epub 2023 Jan 24. Erratum in: *Ann Intern Med.* 2023 Aug;176(8):1143-1144. doi: 10.7326/L23-0246. PMID: 36689752.
24. Reus VI, Fochtmann LJ, Eyler AE, et al. The American Psychiatric Association practice guideline on the use of antipsychotics to treat agitation or psychosis in patients with dementia. *Am J Psychiatry.* 2016;173(5):543-546. doi:10.1176/appi.ajp.2015.173501.
25. Roessner V, Eichele H, Stern JS, et al. European clinical guidelines for Tourette syndrome and other tic disorders-version 2.0. Part III: pharmacological treatment. *Eur Child Adolesc Psychiatry.* 2022;31(3):425-441. doi:10.1007/s00787-021-01899-z.

26. Roxanas MG. Pathological gambling and compulsive eating associated with aripiprazole. *Aust N Z J Psychiatry*. 2010;44(3):291. doi:10.3109/00048670903487282.
27. Sayyah M, Sayyah M, Boostani H, Ghaffari SM, Hoseini A. Effects of aripiprazole augmentation in treatment-resistant obsessive-compulsive disorder (a double blind clinical trial). *Depress Anxiety*. 2012;29(10):850-854. doi:10.1002/da.21996.
28. Seshadri A, Wermers ME, Habermann TJ, Singh B. Long-term Efficacy and Tolerability of Adjunctive Aripiprazole for Major Depressive Disorder: Systematic Review and Meta-analysis. *Prim Care Companion CNS Disord*. 2021 Jun 24;23(4):20r02799. doi: 10.4088/PCC.20r02799.
29. Swetlik C, Cohen LS, Kobylski LA, Sojka ET, Killenberg PC, Freeman MP, Viguera AC. Effects of Prenatal Exposure to Second-Generation Antipsychotics on Development and Behavior Among Preschool-Aged Children: Preliminary Results From the National Pregnancy Registry for Psychiatric Medications. *J Clin Psychiatry*. 2024 Mar 13;85(1):23m14965. doi: 10.4088/JCP.23m14965. PMID: 38488388.
30. Treatment and Management of Mental Health Conditions During Pregnancy and Postpartum: ACOG Clinical Practice Guideline No. 5. *Obstetrics & Gynecology* 141(6):p 1262-1288, June 2023. | DOI: 10.1097/AOG.0000000000005202.
31. World Federation of Societies of Biological Psychiatry; Ihl R, Frölich L, Winblad B, et al. World Federation of Societies of Biological Psychiatry (WFSBP) guidelines for the biological treatment of Alzheimer's disease and other dementias. *World J Biol Psychiatry*. 2011;12(1):2-32. doi:10.3109/15622975.2010.538083

#### Clinical Guideline Revision / History Information

Original Date: 9/21/2023

Reviewed/Revised: 12/19/2024, 10/01/2025, 12/01/2025, 07/01/2026