

## Mayzent (siponimod)

### Disclaimer

*Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.*

*Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.*

Mayzent (siponimod)	1
Summary	1
Definitions	2
Clinical Indications	4
Medical Necessity Criteria for Initial Clinical Review	4
Initial Indication-Specific Criteria	4
Multiple Sclerosis	4
Medical Necessity Criteria for Subsequent Clinical Review	5
Subsequent Indication-Specific Criteria	5
Multiple Sclerosis	5
Experimental or Investigational / Not Medically Necessary	6
References	6
Clinical Guideline Revision / History Information	7

### Summary

Multiple sclerosis (MS) is a chronic, inflammatory, demyelinating disease of the central nervous system. It typically presents in young adults (generally diagnosed before 50 years of age) with symptoms such as vision problems, muscle weakness, numbness, and difficulty with balance and coordination. The most common form is relapsing-remitting MS (occurring in about 85% of patients), characterized by acute attacks followed by periods of remission. Treatment goals include reducing relapses, slowing disability

progression, and managing symptoms. Disease-modifying therapies are the primary treatment approach and include injectable medications (e.g., interferons, glatiramer acetate), oral medications (e.g., dimethyl fumarate, fingolimod, teriflunomide, etc.), and infusion therapies (e.g., natalizumab, ocrelizumab).

MS is a progressive disease, meaning that symptoms tend to worsen over time, and it can be classified into several types, including relapsing-remitting MS (RRMS), primary progressive MS (PPMS), and secondary progressive MS (SPMS).

Currently, there is no cure for MS, but various treatment options are available to manage symptoms, slow the progression of the disease, and improve quality of life.

- Disease-modifying therapies (DMTs) are a class of medications that target the immune system to reduce inflammation and slow down the progression of the disease. The type of DMT prescribed will depend on the type and severity of MS, as well as the individual's medical history and preferences. Some common DMTs include interferon beta, glatiramer acetate, dimethyl fumarate, and fingolimod.
- High dose corticosteroids, such as high dose intravenous methylprednisolone or oral prednisone can be prescribed to reduce inflammation during acute MS relapses.
- Symptomatic treatments are also available to manage specific symptoms of MS, such as muscle spasms, bladder problems, and depression. Physical therapy, occupational therapy, and speech therapy can help individuals with MS maintain mobility, independence, and communication skills.

Mayzent (siponimod) is an oral sphingosine 1-phosphate receptor modulator approved for treating relapsing forms of MS, including clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease. It reduces inflammation and may have neuroprotective effects. Mayzent (siponimod) has demonstrated efficacy in reducing relapse rates and disability progression, particularly in those with active SPMS.

## Definitions

"Clinically isolated syndrome" refers to a first episode of neurologic symptoms lasting at least 24 hours caused by inflammation or demyelination in the central nervous system.

"Compendia" are summaries of drug information and medical evidence to support decision-making about the appropriate use of drugs and medical procedures. Examples include, but are not limited to:

1. American Hospital Formulary Service Drug Information
2. Clinical pharmacology
3. National Comprehensive Cancer Network Drugs and Biologics Compendium
4. Thomson Micromedex DrugDex
5. United States Pharmacopeia-National Formulary (USP-NF)

"CYP2C9 genotype" refers to the genetic variants of the cytochrome P450 2C9 enzyme, which affects the metabolism of siponimod and determines appropriate dosing.

"Disease-modifying therapy" is a medication that modifies the course of MS by reducing relapses and slowing disability progression.

"Documentation" refers to written information, including but not limited to:

- Up-to-date chart notes, relevant test results, and/or relevant imaging reports to support diagnoses; or
- Prescription claims records, and/or prescription receipts to support prior trials of formulary alternatives.

"Multiple sclerosis" is a chronic autoimmune disease of the central nervous system characterized by inflammation, demyelination, and neurodegeneration.

"No evidence of" indicates that the reviewer has not identified any records of the specified item or condition within the submitted materials or claims history. In the absence of such evidence, the member is considered eligible. If any evidence of the item or condition is present upon review of the request, the member does not qualify.

"Primary progressive MS" refers to worsening neurologic function from the onset of symptoms, without early relapses or remissions.

"Relapse" is defined as the appearance of new symptoms or the worsening of existing symptoms lasting at least 24 hours in the absence of fever or infection.

"Relapsing-remitting MS" refers to a disease course characterized by clearly defined attacks of new or increasing neurologic symptoms followed by periods of partial or complete recovery.

"[s]" indicates state mandates may apply.

"Secondary progressive MS" is a disease course following relapsing-remitting MS that is characterized by a progressive worsening of neurologic function over time with or without relapses.

## Clinical Indications

### Medical Necessity Criteria for Initial Clinical Review

#### Initial Indication-Specific Criteria

##### Multiple Sclerosis

The Plan considers Mayzent (siponimod) medically necessary when recent (within the last 3 months) clinical chart documentation provided indicates the member meets ALL of the following:

1. Prescribed by or in consultation with a neurologist or physician who specializes in the treatment of multiple sclerosis; *AND*
2. Is 18 years of age or older; *AND*
3. Has ONE (1) of the following forms of multiple sclerosis:
  - a. Relapsing-remitting (RRMS); *or*
  - b. Active secondary progressive disease (SPMS); *or*
  - c. Clinically isolated syndrome (CIS); *AND*
4. The member is unable to use, or has tried and failed at least TWO (2) of the following:
  - a. An interferon beta product (e.g., Avonex, Betaseron, Plegridy, or Rebif); *and/or*
  - b. Dimethyl Fumarate (generic Tecfidera); *and/or*
  - c. Fingolimod (generic Gilenya); *and/or*
  - d. Glatiramer acetate (Copaxone); *and/or*
  - e. Teriflunomide (generic Aubagio); *AND*
5. The member has been tested for CYP2C9 genotype and does not have the CYP2C9\*3/\*3 genotype; *AND*
6. The member meets ALL of the following criteria:
  - a. No evidence of recent (within the last 6 months) myocardial infarction, unstable angina, stroke, transient ischemic attack (TIA), or decompensated heart failure requiring hospitalization; *and/or*
  - b. No evidence of Class III or IV heart failure; *and/or*
  - c. No evidence of Mobitz type II second-degree or third-degree atrioventricular (AV) block or sick sinus syndrome, unless member has a functioning pacemaker; *AND*
7. Mayzent (siponimod) will be used as monotherapy for multiple sclerosis (i.e., member is not using and will not use other disease-modifying MS therapies while on Mayzent [siponimod]); *AND*
8. Mayzent (siponimod) is being prescribed at a dose and frequency that is within FDA approved labeling OR is supported by compendia or evidence-based published dosing guidelines for the requested indication.
  - o *For members with CYP2C9 genotypes \*1/\*1, \*1/\*2, or \*2/\*2:*
    - i. *Initiate treatment with 5-day titration: 0.25 mg once daily on days 1-2, 0.5 mg on day 3, 0.75 mg on day 4, and 1.25 mg on day 5*
    - ii. *Maintenance dose: 2 mg once daily starting on day 6 and thereafter*
  - o *For members with CYP2C9 genotypes \*1/\*3 or \*2/\*3:*

- i. *Initiate treatment with 4-day titration: 0.25 mg once daily on days 1-2, 0.5 mg on day 3, 0.75 mg on day 4*
- ii. *Maintenance dose: 1 mg once daily starting on day 5 and thereafter*

If the above prior authorization criteria are met, the requested medication will be authorized for up to 12-months.<sup>[s]</sup>

#### *Continued Care*

#### Medical Necessity Criteria for Subsequent Clinical Review

#### Subsequent Indication-Specific Criteria

#### Multiple Sclerosis

The Plan considers Mayzent (siponimod) medically necessary when recent (within the last 6-months) clinical chart documentation provided indicates the member meets BOTH of the following:

1. The requested medication is prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis; *AND*
2. The member has experienced at least ONE (1) of the following:
  - a. Improvement in at least ONE (1) objective measure, such as:
    - i. *Reduced disease activity on MRI; and/or*
    - ii. *Improved or stable disability scores; and/or*
    - iii. *Reduced relapse rate; and/or*
    - iv. *Improved fatigue or walking assessments; AND/OR*
  - b. The member has shown stabilization or improvement in at least ONE (1) MS symptom, such as:
    - i. *Motor function; and/or*
    - ii. *Fatigue; and/or*
    - iii. *Vision; and/or*
    - iv. *Bowel/bladder function; and/or*
    - v. *Spasticity; and/or*
    - vi. *Walking/gait; and/or*
    - vii. *Pain/numbness/tingling.*

If the above reauthorization criteria are met, the requested product will be authorized for up to 12-months.<sup>[s]</sup>

## Experimental or Investigational / Not Medically Necessary<sup>[5]</sup>

Mayzent (siponimod) for any other indication or use is considered not medically necessary by the Plan, as it is deemed to be experimental, investigational, or unproven. Non-covered indications include, but are not limited to, the following:

- Use for the treatment of other neurological conditions not related to multiple sclerosis (e.g., Alzheimer's disease). There are no high quality studies to support the safety and efficacy of Mayzent (siponimod) for the management of Alzheimer's disease; there are no additional on-going studies assessing Mayzent (siponimod) for other neurological conditions to support the safety and efficacy for conditions other than MS.
- Use for the treatment of primary progressive multiple sclerosis. The safety and efficacy of Mayzent (siponimod) has not been established in those with PPMS.
- Use in combination with other disease-modifying therapies for multiple sclerosis. There is limited knowledge about the use of combining DMTs for MS.
- Use in members under 18 years of age. The safety and efficacy in pediatric populations has not been established.
- Use for the treatment of intracerebral hemorrhage, Mayzent (siponimod) has only been studied for this indication in mouse models.
- Use for the treatment of the autoimmune diseases polymyositis and dermatomyositis. There are no high quality studies to support the safety and efficacy of Mayzent (siponimod) for the management of polymyositis and dermatomyositis.

## References

1. Bainbridge JL, Miravalle A, Wong PS. Multiple Sclerosis. In DiPiro JT, Yee GC, Posey LM, et al, eds. *Pharmacotherapy: A Pathophysiologic Approach*. 11th ed. New York, NY: McGraw-Hill; 2019.
2. Benedict RHB, Tomic D, Cree BA, et al,. Siponimod and Cognition in Secondary Progressive Multiple Sclerosis: EXPAND Secondary Analyses. *Neurology*. 2021 Jan 19;96(3):e376-e386. doi: 10.1212/WNL.0000000000011275. Epub 2020 Dec 16.
3. Fox RJ, Cohan S, Mao-Draayer Y, et al. Safety and tolerability of conversion to siponimod from other disease-modifying therapies in patients with advancing forms of relapsing MS: Results from the EXCHANGE study. *Mult Scler*. 2025 May;31(6):706-718. doi: 10.1177/13524585251330085. Epub 2025 Apr 18.
4. Hauser, S., & Cree, B. (2020). Treatment of Multiple Sclerosis: A Review.. *The American journal of medicine*. <https://doi.org/10.1016/j.amjmed.2020.05.049>.
5. He A, Merkel B, Brown JW, et al. Timing of high-efficacy therapy for multiple sclerosis: a retrospective observational cohort study. *Lancet Neurol*. 2020 Apr;19(4):307-316. doi: 10.1016/S1474-4422(20)30067-3. Epub 2020 Mar 18.
6. Kappos L, Bar-Or A, Cree BAC, et al,. Siponimod versus placebo in secondary progressive multiple sclerosis (EXPAND): a double-blind, randomised, phase 3 study. *Lancet*. 2018 Mar 31;391(10127):1263-1273. doi: 10.1016/S0140-6736(18)30475-6. Epub 2018 Mar 23. Erratum in: *Lancet*. 2018 Nov 17;392(10160):2170. doi: 10.1016/S0140-6736(18)32834-4.
7. Kappos L, Bar-Or A, Cree BAC, et al,. Siponimod versus placebo in secondary progressive multiple sclerosis (EXPAND): a double-blind, randomised, phase 3 study. *Lancet*. 2018 Mar 31;391(10127):1263-1273. doi: 10.1016/S0140-6736(18)30475-6. Epub 2018 Mar 23. Erratum in: *Lancet*. 2018 Nov 17;392(10160):2170. doi: 10.1016/S0140-6736(18)32834-4.
8. Mayzent (siponimod) [prescribing information]. East Hanover, NJ: Novartis Pharmaceuticals Corporation; August 2025.

9. McGinley MP, Goldschmidt CH, Rae-Grant AD. Diagnosis and Treatment of Multiple Sclerosis: A Review. *JAMA*. 2021;325(8):765–779. doi:10.1001/jama.2020.26858
10. Montalban X, Leist TP, Cohen BA et al. Cladribine tablets added to IFN- $\beta$  in active relapsing MS: The ONWARD study. *Neurol Neuroimmunol Neuroinflamm*. 2018; 5:e477.
11. Montalban X, Lebrun-Frénay C, Oh J, et al. Diagnosis of multiple sclerosis: 2024 revisions of the McDonald criteria. *Lancet Neurol*. 2025 Oct;24(10):850-865. doi: 10.1016/S1474-4422(25)00270-4. Erratum in: *Lancet Neurol*. 2025 Nov;24(11):e13. doi: 10.1016/S1474-4422(25)00355-2.
12. Multiple Sclerosis Society of Canada. Disease-modifying therapies. <https://mssociety.ca/managing-ms/treatments/medications/disease-modifying-therapies-dmts>.
13. National Institute for Health and Care Excellence [NICE]. Multiple sclerosis in adults: management. NICE Guidelines [NG220]. 22 June 2022. Available at: <https://www.nice.org.uk/guidance/ng220/chapter/Recommendations#ms-symptom-management-and-rehabilitation>. Accessed 20 January 2026.
14. National Institute for Health and Care excellence (NICE). (2020). Siponimod for treating secondary progressive multiple sclerosis (NICE guideline no. TA656). <https://www.nice.org.uk/guidance/ta656/> Accessed 7 April 2025.
15. National MS Society. Disease-modifying therapies for MS (updated March 2022). Available from National MS Society website: <https://nms2cdn.azureedge.net/cmssite/nationalmssociety/media/msnationalfiles/brochures/brochure-the-ms-disease-modifying-medications.pdf>.
16. Rae-Grant A, Day GS, Marrie RA, et al. Practice guideline recommendations summary: Disease-modifying therapies for adults with multiple sclerosis: Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology. *Neurology*. 2018;90(17):777-788.
17. Rashid W, Ciccarelli O, Leary SM, et al. Using disease-modifying treatments in multiple sclerosis: Association of British Neurologists (ABN) 2024 guidance. *Pract Neurol*. 2025 Jan 16;25(1):18-24. doi: 10.1136/pn-2024-004228.
18. Reich DS, Lucchinetti CF, Calabresi PA. 2018. Multiple sclerosis. *New England Journal of Medicine* 378(2):169-180
19. Shahzad F, Rasheed T, Siddiqui MR, et al. The Comparative Effectiveness and Tolerability of Sphingosine-1-Phosphate Receptor Modulators in Patients With Multiple Sclerosis: A Network Meta-Analysis of Randomized Controlled Trials. *Ann Clin Transl Neurol*. 2025 Oct;12(10):2002-2011. doi: 10.1002/acn3.70122. Epub 2025 Jul 4.
20. The use of disease-modifying therapies in multiple sclerosis: principles and current evidence summary. Multiple Sclerosis Coalition. Available from the National MS Society Website: <https://www.nationalmssociety.org/>.
21. Tramacere I, Del Giovane C, Salanti G, et al. Immunomodulators and immunosuppressants for relapsing-remitting multiple sclerosis: a network meta-analysis. *Cochrane Database Syst Rev* 2015;9:CD011381.
22. Wu X, Wang S, Xue T, et al. Disease-modifying therapy in progressive multiple sclerosis: a systematic review and network meta-analysis of randomized controlled trials. *Front Neurol*. 2024;15:1295770. doi:10.3389/fneur.2024.1295770
23. Yang, J., Rempe, T., Whitmire, N., Dunn-Pirio, A., & Graves, J. (2022). Therapeutic Advances in Multiple Sclerosis. *Frontiers in Neurology*, 13. <https://doi.org/10.3389/fneur.2022.824926>.

#### Clinical Guideline Revision / History Information

Original Date: 06/27/2024

Reviewed/Revised: 10/01/2025, 07/01/2026