

Deep Brain Stimulation (DBS) and Responsive Neurostimulation (RNS)

Disclaimer

Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.

Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.

Summary

The Plan members with certain movement disorders may be eligible for treatment with a type of neurostimulation known as deep brain stimulation (DBS). Deep brain stimulation starts with a neurosurgical procedure where electrodes are placed into the area of the brain suspected to be causing the abnormal movements. A separate device that creates the electrical pulses (often called a pulse generator) is also implanted. The device can then be adjusted to help with the symptoms of some intractable movement disorders, such as Parkinson's disease, essential tremor, and certain dystonias.

The Plan members with a seizure disorder may benefit from a different type of neurostimulation known as response neurostimulation (RNS). Instead of electrodes being placed in the deep brain regions, they are surgically implanted into an area of the brain cortex that has been determined to be the focus of the seizures. The RNS device then monitors for abnormal brain activity and when detected, delivers an electrical stimulation in an effort to prevent the seizure activity from propagating. This guideline provides criteria and exclusions for both deep brain stimulation and responsive neurostimulation devices.

Definitions

"Deep brain stimulation (DBS)" is a neurosurgical procedure used to treat a variety of movement disorders. An electrode is placed in the target area of the brain, either on both sides (bilateral) or one

side (unilateral), and electrical pulses are generated to decrease the movement symptoms. Deep brain structures that may be implanted include, but are not limited to:

- Ventral intermediate thalamic nucleus (thalamus)
- Globus Pallidus Internus
- Subthalamic Nucleus

“Responsive Neurostimulation (RNS)” is a neurosurgical procedure where an electrode connected to a pulse generator is implanted into an area of the brain suspected to be the focus of seizure activity. The RN stimulator device system monitors for abnormal electrical activity of the brain and then can deliver electrical activity in an effort to disrupt or lessen the seizure activity in patients with refractory disease.

“Parkinson’s Disease (PD)” is a neurodegenerative movement disorder that results from a loss of dopaminergic neurons in the regions of the brain controlling motor activity. Symptoms of Parkinson’s disease include resting tremor, slow shuffling gait, rigidity, and slow movements. PD is often treated with oral medications, but some patients may benefit more from invasive intervention such as deep brain stimulation when medication fails to improve the symptoms.

“Dystonia” is involuntary muscle contraction that can cause tremors, twisting, and abnormal postures. There are many different types of dystonia, including:

- Generalized dystonia: A more widespread dystonia affecting multiple muscles of the body
- Segmental dystonia: Similar to a focal dystonia, but it affects a segment of the body, such as the upper and lower arm, or the leg and hip
- Hemidystonia: A type of dystonia affecting half of body, such as the arm and leg on the same side
- Cervical dystonia: A type of focal dystonia where spasm of the neck muscles can cause the head to rotate to one side and/or pull towards the chest or back

“Essential Tremor” is a common movement disorder that can cause abnormal shaking of the hands, arms, fingers, or even the vocal cords or head. It usually differs from other movement disorders in that it is an “action tremor”, meaning that the shaking only happens during intentional movement of that body part. Essential tremor is most often treated with oral medications, but may require surgical intervention when the symptoms are disabling and not responsive to medication.

“Focal Seizures” (e.g., focal aware seizures, focal impaired awareness seizures, or partial seizures) are seizures that start in and only impact one part of the brain.

“Generalized Seizures” are seizures that affect both cerebral hemispheres and are more likely to cause impaired level of consciousness.

“Unified Parkinson’s Disease Rating Scale (UPDRS)” is a scale used to measure the severity of the motor complications of Parkinson’s Disease. It ranges from 0-108, with higher scores signifying greater disability.

Clinical Indications

Deep Brain Stimulation (DBS) for Parkinson’s Disease

the Plan considers deep brain stimulation for members with medically refractory Parkinson's Disease medically necessary when ALL of the following criteria are met:

1. The member has idiopathic Parkinson’s disease (i.e., excludes secondary or medication-induced causes; *and*
2. Unified Parkinson’s Disease Rating Scale (UPDRS) score of 30 or greater when without medication for 12 hours; *and*
3. Motor complications that cannot be adequately controlled with standard pharmacologic therapy; *and*
4. The motor impairment affects safety, functional status, or quality of life due to ONE or more of the following symptoms:
 - a. Bradykinesia; *or*
 - b. Levodopa-induced dyskinesia; *or*
 - c. Rigidity; *or*
 - d. Tremor.
5. Treatment with levodopa had previously resulted in improvement in symptoms; *and*
6. There is sufficient motor activity in the affected limb(s) such that the procedure can be reasonably expected to demonstrate improvement; *and*
7. The member has NONE of the following contraindications:
 - a. Evidence of dementia or cognitive impairment; *or*
 - b. Prior intracranial surgery at the target site; *or*
 - c. Focal lesion (such as space occupying lesion) that would impair stimulation or implant placement or may be causing the symptoms; *or*
 - d. High-risk for surgical complications (e.g., active systemic infection, coagulation disorders, etc.); *or*
 - e. Severe depression or suicidal ideation; *or*
 - f. Hoehn and Yahr Stage V Parkinson’s disease, as defined by the following:
 - i. Cachectic state; *or*
 - ii. Inability to stand or walk; *or*
 - iii. Bedbound; *or*
 - iv. Requiring continuous nursing care.
 - g. Cerebellar or pyramidal findings or dysautonomia that may be more suggestive of secondary parkinsonism.

Deep Brain Stimulation (DBS) for Essential Tremors

The Plan considers deep brain stimulation for members with medically refractory essential tremor medically necessary when ALL of the following criteria are met:

1. Alternative etiologies of the tremor have been excluded and/or adequately treated; *and*
2. Symptoms are disabling and/or impact ADLs; *and*
3. Symptoms have been continuously present for at least 3 months despite adequate trials of the following treatments, unless there are specific contraindications or adverse effects:
 - a. First line medications (e.g. propranolol and/or primidone); *and*
 - b. Second line medications (e.g. gabapentin, topiramate, nimodipine, or combinations of multiple agents); *and*
4. Brain imaging shows no evidence of structural abnormality or of another possible etiology; *and*
5. The member has NONE of the following contraindications:
 - a. Dementia or significant cognitive impairment (or Mini-Mental State Examination score of less than 24) ; *or*
 - b. Prior intracranial surgery at the target site; *or*
 - c. Focal lesion (such as space occupying lesion) that would impair stimulation or implant placement or may be causing the symptoms; *or*
 - d. High-risk for surgical complications (e.g., active systemic infection, coagulation disorders, etc.).
6. The stimulation target is the thalamus, globus pallidus, or subthalamic nuclei; *and*
7. The stimulation is unilateral in cases of one-sided tremors or bilateral in cases of tremors on both sides.

Deep Brain Stimulation (DBS) for Epilepsy

The Plan considers Deep Brain Stimulation for epilepsy medically necessary when ALL of the following criteria are met:

1. Member is 18 years or older; *and*
2. Treatment by FDA approved Medtronic DBS system for epilepsy; *and*
3. Presence of focal onset (partial) seizures despite trial of 3 or more antiepileptic medications; *and*
4. Presence of 6 or more partial seizures with or without secondary generalized seizures each month during the previous 3 months with no more than 30 days between seizures; *and*
5. The member does not meet criteria for focal resection epilepsy surgery; *and*
6. The member does not meet criteria for vagus nerve stimulation.

Deep Brain Stimulation (DBS) for Dystonia

The Plan considers deep brain stimulation for members with dystonia medically necessary when ALL of the following criteria are met:

1. The member is 7 years of age or older; *and*
2. The member has been diagnosed with any ONE of the following:
 - a. Primary dystonia, including:

- i. Generalized dystonia; *or*
 - ii. Segmental dystonia; *or*
 - iii. Hemidystonia; *or*
 - iv. Cervical dystonia; *or*
 - b. Secondary tardive dyskinesia/dystonia from ONE of the following:
 - i. Chronic levodopa treatment for Parkinson's Disease; *or*
 - ii. Chronic antidopaminergic effects from psychiatric disease treatment with motor response complications not controlled by medication; *and*
- 3. Dystonia is NOT caused by ANY of the following:
 - a. Stroke; *or*
 - b. Cerebral Palsy; *or*
 - c. Tumor; *or*
 - d. Trauma; *or*
 - e. Infection; *or*
 - f. Multiple sclerosis or other neurodegenerative disease; *or*
 - g. Medication(s); *and*
- 4. Motor complications are refractory to optimal medical management, including but not limited to botulinum toxin injections when applicable; *and*
- 5. Symptoms are disabling and/or impact ADLs; *and*
- 6. The member has NONE of the following contraindications:
 - a. Evidence of dementia or cognitive impairment; *or*
 - b. Prior intracranial surgery at the target site; *or*
 - c. Focal lesion (such as space occupying lesion) that would impair stimulation or implant placement or may be causing the symptoms; *or*
 - d. High-risk for surgical complications (e.g. active systemic infection, coagulation disorders, etc).

Responsive Neurostimulation (RNS)

The Plan considers responsive neurostimulation using an FDA approved device (e.g., NeuroPace RNS System) medically necessary when ALL of the following criteria are met:

- 1. The member is 18 years of age or older; *and*
- 2. Documented diagnosis of intractable focal seizures (simple or complex partial seizures; may also be documented as focal aware or focal impaired awareness seizures); *and*
- 3. Presence of only 1-2 epileptogenic foci with confirmed location; *and*
- 4. Condition is refractory to at least two appropriate antiepileptic drugs at therapeutic doses; *and*
- 5. At least three or more disabling seizures per month averaged over the immediately preceding three month period; where disabling is defined as:
 - a. Significant impairment of ADLs; *or*
 - b. Causing injury; *and*

6. The member is NOT a candidate for focal resective epilepsy surgery, such as for ANY of the following reasons:
 - a. Bitemporal foci, where resection would place the member at risk for memory and/or language deficits; *or*
 - b. Left-sided temporal lobe focus, where diagnostic testing suggests that removal of the target region may result in memory and/or language deficits; *or*
 - c. A focus with overlap of important anatomical regions, where removal may result in functional deficits pertaining to the specific region; *and*
7. Non-epileptic etiologies have been addressed and/or ruled out (e.g. cardiogenic syncope, psychogenic seizures); *and*
8. The member has NONE of the following contraindications:
 - a. Primary generalized epilepsy; *or*
 - b. Rapidly progressive neurologic disorder or evidence of cognitive impairment; *or*
 - c. 3 or more epileptogenic foci; *or*
 - d. Other implanted device(s) delivering electric energy to the brain; *or*
 - e. High-risk for surgical complications (e.g., active systemic infection, coagulation disorder); *or*
 - f. Focal lesion (such as space occupying lesion) that would impair stimulation or implant placement or may be causing the symptoms; *or*
 - g. Epileptogenic foci located inferior to the level of the subthalamic nucleus.

DBS or RNS Device Replacement

Replacement of deep brain stimulation or responsive neurostimulation devices is considered medically necessary when ALL of the following criteria are met:

1. Device is no longer functioning properly and cannot be safely repaired; *and*
2. Physician documents need for replacement; *and*
3. Device is no longer under the original warranty; *and*
4. Original placement was considered medically necessary per the above criteria; *and*
5. There are no contraindications to device replacement.

Experimental or Investigational / Not Medically Necessary

Deep brain stimulation for any other indication is considered experimental, investigational, or unproven.

Non-covered indications include, but are not limited to, the following:

- Addiction
- Alzheimer's disease
- Eating disorders, e.g. Anorexia nervosa
- Autism spectrum disorder
- Bipolar Disorder
- Blepharospasm
- Cerebral palsy

- Chronic cluster headache
- Chronic pain syndrome including complex regional pain syndrome/reflex sympathetic dystrophy, cancer-related pain, and post-stroke pain.
- Chronic vegetative state
- Depression
- Degenerative, metabolic and infectious disorders
- Dravet Syndrome
- Drug-induced movement disorders
- Explosive aggressive behavior
- Head or voice tremor, except when symptoms are due to covered essential tremor meeting the above criteria
- Huntington's disease or other choreiform disorders
- Minimally conscious state
- Multiple sclerosis
- Obesity
- Obsessive-compulsive disorder
- Orthostatic tremor
- Parkinson's disease-related camptocormia, dysarthria/speech deficits, and gait disorders (e.g., gait instability and freezing of gait)
- Post-traumatic tremor
- Secondary dystonias
- Self-injurious behavior
- Substance use disorders
- SUNCT syndrome
- Tourette syndrome
- Traumatic brain injury

Evidence for Experimental or Investigational / Not Medically Necessary

Addiction

The evidence for DBS in addiction is limited and the existing studies have yet to clearly define the optimal target for neurostimulation in patients with addiction.

Alzheimer's Disease

The existing literature for DBS in AD is limited to small, retrospective studies and case reports. Furthermore, the results have yet to show clear clinical efficacy and safety in this population. Further evidence is needed to define the role of DBS in patients with Alzheimer's disease.

Bipolar disorder

The evidence for DBS in bipolar disorder is limited to small studies and a review paper covering only 12 patients. An UpToDate review on the topic states that "It is not known if DBS is efficacious as adjunctive treatment for bipolar disorder due to the limited and low quality data that are available". There has been

some concern in the literature for the possibility of DBS causing treatment-related mania or hypomania, which may worsen in patients with underlying bipolar disorder. A study by Holtzheimer et al (2012) found no episodes of mania/hypomania in the 7 patients treated with DBS. Another review on 5 case reports found 1 patient with hypomanic symptoms. Other studies of DBS for other indications have suggested higher rates of the side effect, however. Further evidence is needed in the form of larger, preferably randomized studies, to determine the potential benefit and safety of DBS for this indication.

Chronic Pain

The existing evidence is limited to anecdotal reports and case studies. An older publication of the EFNS guidelines in 2007 reported that neurostimulation for neuropathic pain may be efficacious but should only be performed at experienced centers. Further evidence is needed to determine the potential efficacy and safety for DBS in this indication.

Cluster Headache

Deep brain stimulation has been considered for the use of medically refractory cluster headache. UpToDate considers DBS for this indication "investigational and [requiring] further study to confirm long-term benefit and safety". Fontaine et al (2010) looked at 11 patients randomized to periods of DBS for their cluster headache vs. sham treatment periods. They found that "during the randomized phase, no significant change in primary and secondary outcome measures was observed between active and sham stimulation". They also found three serious adverse events, and concluded "Randomized phase findings of this study did not support the efficacy of DBS in refractory CCH". Further evidence is needed to confirm any potential benefit with DBS for this indication.

Depression

The existing evidence for DBS in depression is limited. UpToDate reports that "Deep brain stimulation (DBS) is an investigational procedure for treatment resistant depression".

Dravet Syndrome

The evidence for DBS in Dravet Syndrome is limited to case reports without any significant large studies. An UpToDate review concluded that "It is therefore not yet possible to draw conclusions on the effectiveness of DBS in these patients."

Eating Disorders

The existing evidence for DBS in eating disorders is very limited. One study by Lipsman et al (2013) looked at six females with severe, refractory anorexia treated with DBS, with 50% of the patients showing weight gain and 4 out of 6 with improvement in their mood symptoms. That same research group later added 10 more patients to the study. That same year, Wu et al (2013) looked at four more patients with anorexia undergoing DBS. As of the present time, the literature regarding the safety and efficacy of DBS for this indication is limited to these very small, single institution series. Per UpToDate, DBS is experimental and investigational for the treatment of eating disorders.

Huntington's Disease

Per UpToDate, "The utility of deep brain stimulation in HD is unknown. Data are limited to case studies and small case series, which suggest some benefit in chorea...However, this may be at the expense of cognition." Further prospective evidence is needed to determine any potential benefit.

Other Choreiform disorders

The existing data for the use of DBS for choreiform disorders is limited to small, conflicting studies. One study by Edwards et al in 2012 looked at 16 case reports on patients treated with DBS for chorea. Further evidence in the form of randomized, prospective data is needed to define the clinical role and potential benefit of DBS for this indication.

Obsessive Compulsive Disorder

In Aug. 2022, Hayes rated D2 Deep Brain Stimulation (DBS) for the Treatment of Refractory Obsessive-Compulsive Disorder as an add-on therapy for obsessive-compulsive disorder (OCD) in adult patients with inadequate responses to ≥ 3 prior treatments and no contraindications to DBS. DBS has been inadequately studied for this indication. An UpToDate review on the topic concludes that "Deep brain stimulation is an investigational treatment for incapacitating, treatment-refractory OCD." A 2015 meta-analysis of several small randomized trials on the subject suggested that DBS may reduce the symptoms of OCD, but concluded that larger trials would be needed to confirm the findings. The study looked at 31 studies involving 116 subjects, finding an average symptom reduction on the Y-BOCS scale of 45.1% and an overall response rate of 60%. Despite these findings, larger, randomized evidence is needed to determine the optimal implant location, safety, and efficacy of DBS for this indication.

SUNCT Syndrome

The data on DBS for SUNCT syndrome is currently limited to very small, uncontrolled studies and case reports. Furthermore, the optimal location in the brain for DBS for SUNCT has not yet been fully determined. One study by Leone et al (2016) looked at 11 patients with refractory SUNCT treated with ipsilateral DBS. They reported a 78% median reduction in daily attack frequency. An UpToDate review of this study and topic reports that "Although these results are promising, further study is needed to evaluate the utility of DBS for patients with SUNCT and SUNA."

Tourette Syndrome

The existing evidence for DBS in Tourette syndrome is limited. A randomized, double blinded crossover trial published in The Lancet in 2015 looked at 15 patients with severe, medically refractory Tourette syndrome. The patients received surgery for DBS in the globus pallidus followed by 3 months of stimulation vs. 3 months of no stimulation, and then a cross over to the other group. Only 13 patients completed the randomization and crossover. The authors concluded that there was a significant improvement in tic severity, and that "Future research should concentrate on identifying the most effective target for DBS to control both tics and associated comorbidities, and further clarify factors that predict individual patient response". Given the limitation in the number of patients and available

evidence, further data is needed to determine the efficacy and safety of DBS for this indication. An UpToDate review on the topic concluded that “However, the available evidence is preliminary, and larger clinical trials are needed to determine whether DBS is beneficial for controlling tics in patients with TS.”

For indications not specifically listed above, the existing evidence is limited and/or insufficient to determine the potential efficacy and safety of DBS.

Applicable Billing Codes (HCPCS/CPT Codes)

<i>Deep Brain Stimulation and Responsive Neurostimulation</i>	
CPT/HCPCS Codes considered medically necessary if criteria are met:	
<i>Code</i>	<i>Description</i>
61850	Twist drill or burr hole(s) for implantation of neurostimulator electrodes, cortical
61860	Craniectomy or craniotomy for implantation of neurostimulator electrodes, cerebral, cortical
61863	Twist drill, burr hole, craniotomy, or craniectomy with stereotactic implantation of neurostimulator electrode array in subcortical site (e.g., thalamus, globus pallidus, subthalamic nucleus, periventricular, periaqueductal gray), without use of intraoperative microelectrode recording; first array
61864	Twist drill, burr hole, craniotomy, or craniectomy with stereotactic implantation of neurostimulator electrode array in subcortical site (eg, thalamus, globus pallidus, subthalamic nucleus, periventricular, periaqueductal gray), without use of intraoperative microelectrode recording; each additional array (List separately in addition to primary procedure)
61867	Twist drill, burr hole, craniotomy, or craniectomy with stereotactic implantation of neurostimulator electrode array in subcortical site (e.g., thalamus, globus pallidus, subthalamic nucleus, periventricular, periaqueductal gray), with use of intraoperative microelectrode recording; first array
61868	Twist drill, burr hole, craniotomy, or craniectomy with stereotactic implantation of neurostimulator electrode array in subcortical site (eg, thalamus, globus pallidus, subthalamic nucleus, periventricular, periaqueductal gray), with use of intraoperative microelectrode recording; each additional array (List separately in addition to primary procedure)
61880	Revision or removal of intracranial neurostimulator electrodes
61885	Insertion or replacement of cranial neurostimulator pulse generator or receiver, direct or inductive coupling; with connection to a single electrode array
61886	Insertion or replacement of cranial neurostimulator pulse generator or receiver, direct or inductive coupling; with connection to 2 or more electrode arrays

61888	Revision or removal of cranial neurostimulator pulse generator or receiver
95836	Electrocorticogram from an implanted brain neurostimulator pulse generator/transmitter, including recording, with interpretation and written report, up to 30 days
95970	Electronic analysis of implanted neurostimulator pulse generator/transmitter (eg, contact group[s], interleaving, amplitude, pulse width, frequency [Hz], on/off cycling, burst, magnet mode, dose lockout, patient selectable parameters, responsive neurostimulation, detection algorithms, closed loop parameters, and passive parameters) by physician or other qualified health care professional; with brain, cranial nerve, spinal cord, peripheral nerve, or sacral nerve, neurostimulator pulse generator/transmitter, without programming
95976	Electronic analysis of implanted neurostimulator pulse generator/transmitter (eg, contact group[s], interleaving, amplitude, pulse width, frequency [Hz], on/off cycling, burst, magnet mode, dose lockout, patient selectable parameters, responsive neurostimulation, detection algorithms, closed loop parameters, and passive parameters) by physician or other qualified health care professional; with simple cranial nerve neurostimulator pulse generator/transmitter programming by physician or other qualified health care professional
95977	Electronic analysis of implanted neurostimulator pulse generator/transmitter (eg, contact group[s], interleaving, amplitude, pulse width, frequency [Hz], on/off cycling, burst, magnet mode, dose lockout, patient selectable parameters, responsive neurostimulation, detection algorithms, closed loop parameters, and passive parameters) by physician or other qualified health care professional; with complex cranial nerve neurostimulator pulse generator/transmitter programming by physician or other qualified health care professional
95983	Electronic analysis of implanted neurostimulator pulse generator/transmitter (eg, contact group[s], interleaving, amplitude, pulse width, frequency [Hz], on/off cycling, burst, magnet mode, dose lockout, patient selectable parameters, responsive neurostimulation, detection algorithms, closed loop parameters, and passive parameters) by physician or other qualified health care professional; with brain neurostimulator pulse generator/transmitter programming, first 15 minutes face-to-face time with physician or other qualified health care professional
95984	Electronic analysis of implanted neurostimulator pulse generator/transmitter (eg, contact group[s], interleaving, amplitude, pulse width, frequency [Hz], on/off cycling, burst, magnet mode, dose lockout, patient selectable parameters, responsive neurostimulation, detection algorithms, closed loop parameters, and passive parameters) by physician or other qualified health care professional; with brain neurostimulator pulse generator/transmitter programming, each additional 15 minutes face-to-face time with physician or other qualified health care professional (List separately in addition to code for primary procedure)
C1767	Generator, neurostimulator (implantable), nonrechargeable
C1778	Lead, neurostimulator (implantable)
C1816	Receiver and/or transmitter, neurostimulator (implantable)

C1822	Generator, neurostimulator (implantable), high frequency, with rechargeable battery and charging system
C1883	Adaptor/ extension, pacing lead or neurostimulator lead (implantable)
C1897	Lead, neurostimulator test kit (implantable)
E0745	Neuromuscular stimulator, electronic shock unit
L8679	Implantable neurostimulator, pulse generator, any type
L8680	Implantable neurostimulator electrode, each
L8681	Patient programmer (external) for use with implantable programmable neurostimulator pulse generator, replacement only
L8682	Implantable neurostimulator radiofrequency receiver
L8683	Radiofrequency transmitter (external) for use with implantable neurostimulator radiofrequency receiver
L8685	Implantable neurostimulator pulse generator, single array, rechargeable, includes extension
L8686	Implantable neurostimulator pulse generator, single array, nonrechargeable, includes extension
L8687	Implantable neurostimulator pulse generator, dual array, rechargeable, includes extension
L8688	Implantable neurostimulator pulse generator, dual array, nonrechargeable, includes extension
L8689	External recharging system for battery (internal) for use with implantable neurostimulator, replacement only
L8695	External recharging system for battery (external) for use with implantable neurostimulator, replacement only
ICD-10 codes considered medically necessary if criteria are met:	
<i>Code</i>	<i>Description</i>
G20	Parkinson's disease
G21.0 - G21.9	Secondary Parkinsonism
G24.1 - G24.4, G24.8 - G24.9	Dystonia [excludes drug-induced]
G25.0, G25.2	Essential tremor [familial tremor]
G40.00 - G40.91	Epilepsy and recurrent seizures [Deep Brain Stimulation and Responsive Neurostimulation]

R56.9	Unspecified convulsions [for Responsive Neurostimulation only]
ICD-10 codes <i>not</i> considered medical necessary or considered experimental or investigational:	
<i>Code</i>	<i>Description</i>
A00.0 - B99.9	Infectious and parasitic diseases
E66.01 - E66.9	Obesity
E70.0 - E88.9	Metabolic disorders
F10.10 - F19.99	Alcohol and drug related disorders [includes abuse, dependence and use]
F31.0 - F31.9	Bipolar disorder
F32.0 - F34.9	Depression
F42.8 - F42.9	Obsessive-compulsive disorder
F44.4	Conversion disorder with motor symptom or deficit [Parkinson's disease-related camptocormia]
F50.0 - F50.9	Eating disorders
F63.81	Intermittent explosive disorder [explosive aggressive behavior]
F84.0	Autism spectrum disorder
F95.2	Tourette syndrome
G10	Huntington's chorea
G23.0 - G23.9	Other degenerative diseases of the basal ganglia
G24.01 - G24.09	Drug induced dystonia
G24.5	Blepharospasm
G30.0 - G30.9	Alzheimer's disease
G35	Multiple sclerosis
G44.021 - G44.029	Chronic cluster headache
G44.059	Short lasting unilateral neuralgiform headache with conjunctival injection and tearing (SUNCT), not intractable
G80.0 - G80.9	Cerebral palsy
G89.3	Neoplasm related pain (acute) (chronic)
G89.4	Chronic pain syndrome

G90.3	Multi-system degeneration of the autonomic nervous system
G90.50 - G90.59	Complex regional pain syndrome
I69.398	Other sequelae of cerebral infarction [when used to identify post-stroke pain]
R40.0 - R40.4	Somnolence, stupor and coma [alteration of consciousness]
S06.0x0A - S06.9x9S	Intracranial injury
X71.0xxA - X83.8xxS	Intentional self-harm

References

1. Abramowitz JS, Taylor S, McKay D. Obsessive-compulsive disorder. *Lancet*. 2009;374(9688):491-499.
2. Akbostanci MC, Slavin KV, Burchiel KJ. Stereotactic ventral intermedial thalamotomy for the treatment of essential tremor: results of a series of 37 patients. *Stereotact Funct Neurosurg* 1999; 72:174.
3. Akram H, Miller S, Lagrata S, et al. Ventral tegmental area deep brain stimulation for refractory chronic cluster headache. *Neurology*. 2016;86(18):1676-1682.
4. Alesch F, et al. Stimulation of the ventral intermediate thalamic nucleus in tremor dominated Parkinson's disease and essential tremor. *Acta Neurochir*. 1995;136:75-81.
5. Alonso P, Cuadras D, Gabriels L, et al. Deep brain stimulation for obsessive-compulsive disorder: A meta-analysis of treatment outcome and predictors of response. *PLoS One*. 2015;10(7):e0133591.
6. Alterman RL, Miravite J, Weisz D, et al. Sixty hertz pallidal deep brain stimulation for primary torsion dystonia. *Neurology* 2007; 69:681.
7. Alterman RL, Snyder BJ. Deep brain stimulation for torsion dystonia. *Acta Neurochir Suppl* 2007; 97:191.
8. Andrade DM, Hamani C, Lozano AM, Wennberg RA. Dravet syndrome and deep brain stimulation: seizure control after 10 years of treatment. *Epilepsia* 2010; 51:1314.
9. Andrade DM, Zumsteg D, Hamani C, et al. Long-term follow-up of patients with thalamic deep brain stimulation for epilepsy. *Neurology* 2006; 66:1571.
10. Anouti A, Koller WC. Tremor disorders: Diagnosis and management. *West J Med*. 1995;162:510-513.
11. Appleby BS, Duggan PS, Regenbreg A, Rabins PV. Psychiatric and neuropsychiatric adverse events associated with deep brain stimulation: A meta-analysis of ten years' experience. *Mov Disord* 2007; 22:1722.
12. Bartsch C, Kuhn J.[Deep brain stimulation for addiction, anorexia and compulsion. Rationale, clinical results and ethical implications. *Nervenarzt*. 2014;85(2):162-168.

13. Benabid AL, et al. Chronic electrical stimulation of the ventralis intermedius nucleus of the thalamus as a treatment of movement disorders. *J Neurosurg.* 1996;84:203-214.
14. Benabid AL, Koudsie A, Benazzouz A, et al. Deep brain stimulation of the corpus luyisi (subthalamic nucleus) and other targets in Parkinson's disease. Extension to new indications such as dystonia and epilepsy. *J Neurol.* 2001;248(Suppl 3):III37-III47.
15. Bergey GK, Morrell MJ, Mizrahi EM, et al. Long-term treatment with responsive brain stimulation in adults with refractory partial seizures. *Neurology* 2015; 84:810.
16. Berlim MT, McGirr A, Van den Eynde F, et al. Effectiveness and acceptability of deep brain stimulation (DBS) of the subgenual cingulate cortex for treatment-resistant depression: A systematic review and exploratory meta-analysis. *J Affect Disord.* 2014;159:31-38.
17. Berman BD, Starr PA, Marks WJ Jr, Ostrem JL. Induction of bradykinesia with pallidal deep brain stimulation in patients with cranial-cervical dystonia. *Stereotact Funct Neurosurg* 2009; 87:37.
18. Blahak C, Capelle HH, Baezner H, et al. Micrographia induced by pallidal DBS for segmental dystonia: a subtle sign of hypokinesia? *J Neural Transm (Vienna)* 2011; 118:549.
19. Blomstedt P, Sandvik U, Tisch S. Deep brain stimulation in the posterior subthalamic area in the treatment of essential tremor. *Mov Disord* 2010; 25:1350.
20. Blond S, et al. Control of tremor and involuntary movement disorders by chronic stereotactic stimulation of the ventral intermediate thalamic nucleus. *J Neurosurg.* 1992;77:62-68.
21. Boon P, Vonck K, De Herdt V, et al. Deep brain stimulation in patients with refractory temporal lobe epilepsy. *Epilepsia.* 2007;48(8):1551-1560.
22. Bronte-Stewart H. New Drugs and Devices. Deep brain stimulation. *Neurol Clin Prac.* 2012;291):67-71.
23. Burchiel KJ, Anderson VC, Favre J, et al. Comparison of pallidal and subthalamic nucleus deep brain stimulation for advanced Parkinson's disease: Results of a randomized, blinded pilot study. *Neurosurgery.* 1999;45(6):1375-1384.
24. Carpenter LL, Friehs GM, Tyrka AR, et al. Vagus nerve stimulation and deep brain stimulation for treatment resistant depression. *Med Health R I.* 2006;89(4):137, 140-141.
25. Carpenter MA, Pahwa R, Miyawaki KL, et al. Reduction in voice tremor under thalamic stimulation. *Neurology* 1998; 50:796.
26. Castelnau P, Cif L, Valente EM, et al. Pallidal stimulation improves pantothenate kinase-associated neurodegeneration. *Ann Neurol* 2005; 57:738.
27. Chabardes S, Kahane P, Minotti L, et al. Deep brain stimulation in epilepsy with particular reference to the subthalamic nucleus. *Epileptic Disord.* 2002;4 Suppl 3:S83-S93.
28. Chieng LO, Madhavan K, Wang MY. Deep brain stimulation as a treatment for Parkinson's disease related camptocormia. *J Clin Neurosci.* 2015;22(10):1555-1561.
29. Chou KL, Tarsy D. Device-assisted and lesioning procedures for Parkinson disease. *UpToDate.com.* Last updated March 4, 2022.
https://www.uptodate.com/contents/device-assisted-and-lesioning-procedures-for-parkinson-disease?sectionName=DEEP%20BRAIN%20STIMULATION&search=deep%20brain%20stimulation&topicRef=90611&anchor=H5&source=see_link#H5

30. Cif L, Vasques X, Gonzalez V, et al. Long-term follow-up of DYT1 dystonia patients treated by deep brain stimulation: an open-label study. *Mov Disord* 2010; 25:289.
31. Cleary DR, Ozpinar A, Raslan AM, Ko AL. Deep brain stimulation for psychiatric disorders: where we are now. *Neurosurgical Focus* 2015;38(6):E2
32. Cohen-Gadol AA, Britton JW, Wetjen NM, et al. Neurostimulation therapy for epilepsy: current modalities and future directions. *Mayo Clin Proc* 2003; 78:238.
33. Collins TR. Brain stimulation device found to reduce seizures in randomized trial. *Neurology Today*, October 21, 2010. pg. 14.
34. Cruccu G, Aziz TZ, Garcia-Larrea L, et al. EFNS guidelines on neurostimulation therapy for neuropathic pain. *Eur J Neurol*. 2007;14(9):952-970.
35. Cukiert A, Cukiert CM, Burattini JA, et al. Seizure outcome after hippocampal deep brain stimulation in patients with refractory temporal lobe epilepsy: A prospective, controlled, randomized, double-blind study. *Epilepsia* 2017; 58:1728.
36. Denys D, Koning P. Deep brain stimulation for treatment of obsessive-compulsive disorder. UpToDate.com. Last Updated March 8, 2022. Retrieved from https://www.uptodate.com/contents/deep-brain-stimulation-for-treatment-of-obsessive-compulsive-disorder?search=deep%20brain%20stimulation&source=search_result&selectedTitle=2~92&usage_type=default&display_rank=2
37. Deuschl G, Bain P. Deep brain stimulation for tremor [correction of trauma]: Patient selection and evaluation. *Mov Disord*. 2002;17 Suppl 3:S102-S111.
38. Deuschl G. Neurostimulation for Parkinson disease. *JAMA*. 2009;301(1):104-105.
39. Deutschländer A, Asmus F, Gasser T, et al. Sporadic rapid-onset dystonia-parkinsonism syndrome: failure of bilateral pallidal stimulation. *Mov Disord* 2005; 20:254.
40. Devinsky O, Sato S, Kufta CV, et al. Electroencephalographic studies of simple partial seizures with subdural electrical recordings. *Neurology*. 1989;39(4):527-533.
41. Diaz-Arrastia R, Agostini MA, Van Ness PC. Evolving treatment strategies for epilepsy. *JAMA*. 2002;287(22):2917-2920.
42. Edwards TC, Zrinzo L, Limousin P, Foltynie T. Deep brain stimulation in the treatment of chorea. *Movement Disorders* 2012;27(3):357-63.
43. Eitan R, Lerer B. Nonpharmacological, somatic treatments of depression: Electroconvulsive therapy and novel brain stimulation modalities. *Dialogues Clin Neurosci*. 2006;8(2):241-258.
44. Elias WJ, Huss D, Voss T, et al. A pilot study of focused ultrasound thalamotomy for essential tremor. *N Engl J Med* 2013; 369:640.
45. Elias WJ, Lipsman N, Ondo WG, et al. A Randomized Trial of Focused Ultrasound Thalamotomy for Essential Tremor. *N Engl J Med* 2016; 375:730.
46. Eltahawy HA, Saint-Cyr J, Giladi N, et al. Primary dystonia is more responsive than secondary dystonia to pallidal interventions: outcome after pallidotomy or pallidal deep brain stimulation. *Neurosurgery* 2004; 54:613.
47. Eskandar EN, Cosgrove GR, Shinobu LA. Surgical treatment of Parkinson Disease. *JAMA*. 2001;286(24):3056-3059.

48. Fayad SM, Guzick AG, Reid AM, et al. Six-Nine Year Follow-Up of Deep Brain Stimulation for Obsessive-Compulsive Disorder. *PLoS One* 2016; 11:e0167875.
49. Fields JA, Troster AI, Woods SP, et al. Neuropsychological and quality of life outcomes 12 months after unilateral thalamic stimulation for essential tremor. *J Neurol Neurosurg Psychiatry*. 2003;74(3):305-311.
50. Figueiras-Mendez R, et al. Further supporting evidence of beneficial subthalamic stimulation in Parkinson's patients. *Neurology*. 2002;58:469-470.
51. Fins JJ, Mayberg HS, Nuttin B, et al. Misuse of the FDA's humanitarian device exemption in deep brain stimulation for obsessive-compulsive disorder. *Health Aff (Millwood)*. 2011;30(2):302-311.
52. Fisher R, Salanova V, Witt T, et al. Electrical stimulation of the anterior nucleus of thalamus for treatment of refractory epilepsy. *Epilepsia* 2010; 51:899.
53. Fisher R, Salanova V, Witt T, et al; SANTE Study Group. Electrical stimulation of the anterior nucleus of thalamus for treatment of refractory epilepsy. *Epilepsia*. 2010;51(5):899-908.
54. Fisher RS. Therapeutic devices for epilepsy. *Ann Neurol* 2012; 71:157.
55. Fitzgerald PB, Brown TL, Marston NA, et al. Transcranial magnetic stimulation in the treatment of depression: a double-blind, placebo-controlled trial. *Arch Gen Psychiatry* 2003; 60:1002.
56. Flora ED, Perera CL, Cameron AL, Maddern GJ. Deep brain stimulation for essential tremor: a systematic review. *Mov Disord* 2010; 25:1550.
57. Fontaine D, Lanteri-Minet M, Ouchchane L, et al. Anatomical location of effective deep brain stimulation electrodes in chronic cluster headache. *Brain* 2010; 133:1214.
58. Fontaine D, Lazorthes Y, Mertens P, et al. Safety and efficacy of deep brain stimulation in refractory cluster headache: a randomized placebo-controlled double-blind trial followed by a 1-year open extension. *J Headache Pain* 2010; 11:23.
59. Foote KD, Seignourel P, Fernandez HH, et al. Dual electrode thalamic deep brain stimulation for the treatment of posttraumatic and multiple sclerosis tremor. *Neurosurgery*. 2006;58(4 Suppl 2):ONS-280-285; discussion ONS-285-286.
60. Forman, SF. Eating disorders: Overview of prevention and treatment. UpToDate. UpToDate.com. Waltham, MA. Accessed December 22, 2017.
61. Fountas KN, Kapsalaki E, Hadjigeorgiou G. Cerebellar stimulation in the management of medically intractable epilepsy: A systematic and critical review. *Neurosurg Focus*. 2010;29(2):E8.
62. Fox MD, Alterman RL. Brain Stimulation for Torsion Dystonia. *JAMA Neurol* 2015; 72:713.
63. Fraix V, Pollak P, Van Blercom N, et al. Effect of subthalamic nucleus stimulation on levodopa-induced dyskinesia in Parkinson's disease. *Neurology*. 2000; 55:1921-1923.
64. Gallo BV. Epilepsy, surgery, and the elderly. *Epilepsy Res*. 2006;68 Suppl 1:S83-S86.
65. Galvez-Jimenez N, Lozano A, Tasker R, et al. Pallidal stimulation in Parkinson's disease patients with a prior unilateral pallidotomy. *Can J Neurol Sci*. 1998;25(4):300-305.
66. Ge Y, Hu W, Liu C, et al. Brain stimulation for treatment of refractory epilepsy. *Chin Med J (Engl)*. 2013;126(17):3364-3370.
67. Georgiopoulos M, Katsakiori P, Kefalopoulou Z, et al. Vegetative state and minimally conscious state: A review of the therapeutic interventions. *Stereotact Funct Neurosurg*. 2010;88(4):199-207.

68. Gerschlager W, Alesch F, Cunnington R, et al. Bilateral subthalamic nucleus stimulation improves frontal cortex function in Parkinson's disease. An electrophysiological study of the contingent negative variation. *Brain*. 1999;122 (Pt 12):2365-2373.
69. Gervais-Bernard H, Xie-Brustolin J, Mertens P, et al. Bilateral subthalamic nucleus stimulation in advanced Parkinson's disease: Five year follow-up. *J Neurol*. 2009;256(2):225-233.
70. Giacino J, Fins JJ, Machado A, Schiff ND. Central thalamic deep brain stimulation to promote recovery from chronic posttraumatic minimally conscious state: Challenges and opportunities. *Neuromodulation*. 2012;15(4):339-349.
71. Giordano F, Cavallo M, Spacca B, et al. Deep brain stimulation of the anterior limb of the internal capsule may be efficacious for explosive aggressive behaviour. *Stereotact Funct Neurosurg*. 2016;94(6):371-378.
72. Gippert SM, Switala C, Bewernick BH, et al. Deep brain stimulation for bipolar disorder-review and outlook. *CNS Spectr* 2017; 22:254.
73. Gloss D, Nolan SJ, Staba R. The role of high-frequency oscillations in epilepsy surgery planning. *Cochrane Database Syst Rev*. 2014;1:CD010235.
74. Golestanirad L, Elahi B, Graham SJ, et al. Efficacy and safety of pedunculopontine nuclei (PPN) deep brain stimulation in the treatment of gait disorders: A meta-analysis of clinical studies. *Can J Neurol Sci*. 2016;43(1):120-126.
75. Gonzalez V, Cif L, Biolsi B, et al. Deep brain stimulation for Huntington's disease: long-term results of a prospective open-label study. *J Neurosurg* 2014; 121:114.
76. Goodman JH. Brain stimulation as a therapy for epilepsy. *Adv Exp Med Biol*. 2004;548:239-247.
77. Greenberg BD, Gabriels LA, Malone DA Jr, et al. Deep brain stimulation of the ventral internal capsule/ventral striatum for obsessive-compulsive disorder: worldwide experience. *Mol Psychiatry* 2010; 15:64.
78. Greenberg BD, Rezai AR. Mechanisms and the current state of deep brain stimulation in neuropsychiatry. *CNS Spectr*. 2003;8(7):522-526.
79. Hallett M, Evinger C, Jankovic J, Stacy M; BEBRF International Workshop. Update on blepharospasm: Report from the BEBRF International Workshop. *Neurology*. 2008;71(16):1275-1282.
80. Halpern C, Hurtig H, Jaggi J, et al. Deep brain stimulation in neurologic disorders. *Parkinsonism Relat Disord*. 2007;13(1):1-16.
81. Halpern CH, Torres N, Hurtig HI, et al. Expanding applications of deep brain stimulation: A potential therapeutic role in obesity and addiction management. *Acta Neurochir (Wien)*. 2011;153(12):2293-2306.
82. Halpern CH, Wolf JA, Bale TL, et al. Deep brain stimulation in the treatment of obesity. *J Neurosurg*. 2008;109(4):625-634.
83. Hamel W, Herzog J, Kopper F, et al. Deep brain stimulation in the subthalamic area is more effective than nucleus ventralis intermedius stimulation for bilateral intention tremor. *Acta Neurochir (Wien)*. 2007;149(8):749-758; discussion 758.
84. Hardenacke K, Shubina E, Bührle CP, et al. Deep brain stimulation as a tool for improving cognitive functioning in Alzheimer's dementia: A systematic review. *Front Psychiatry*. 2013;4:159.

85. Hassan A, Ahlskog JE, Matsumoto JY, et al. Orthostatic tremor: Clinical, electrophysiologic, and treatment findings in 184 patients. *Neurology*. 2016;86(5):458-464.
86. Hayes, Inc., Health Technology Assessment. *Deep Brain Stimulation for the Treatment of Refractory Obsessive-Compulsive Disorder*. Lansdale, PA: Hayes, Inc.; August 24, 2022.
87. Hayes, Inc. Health Technology Assessment. Deep Brain Stimulation of the Anterior Nucleus of the Thalamus for Treatment of Refractory Epilepsy. Lansdale, PA: Hayes, Inc.; Dec 29, 2022
88. Heck CN, King-Stephens D, Massey AD, et al. Two-year seizure reduction in adults with medically intractable partial onset epilepsy treated with responsive neurostimulation: Final results of the RNS System Pivotal trial. *Epilepsia*, **(*) :1–10, 2014.
89. Heschem S, Lim LW, Jahanshahi A, et al. Deep brain stimulation in dementia-related disorders. *Neurosci Biobehav Rev*. 2013;37(10 Pt 2):2666-2675.
90. Hodaie M, Wennberg RA, Dostrovsky JO, Lozano AM. Chronic anterior thalamus stimulation for intractable epilepsy. *Epilepsia*. 2002;43(6):603-608.
91. Holtzheimer PE 3rd, Nemeroff CB. Advances in the treatment of depression. *NeuroRx*. 2006;3(1):42-56.
92. Holtzheimer PE, Kelley ME, Gross RE, et al. Subcallosal cingulate deep brain stimulation for treatment-resistant unipolar and bipolar depression. *Arch Gen Psychiatry* 2012; 69:150.
93. Holtzheimer PE. Bipolar disorder in adults: Overview of neuromodulation procedures. UpToDate. UpToDate.com. Waltham, MA. Accessed December 22, 2017.
94. Hosomi K, Seymour B, Saitoh Y. Modulating the pain network--neurostimulation for central poststroke pain. *Nature Reviews. Neurology* 2015;11(5):290-9
95. Hubble JP, Busenbark KL, Wilkinson S, et al. Deep brain stimulation for essential tremor. *Neurology* 1996; 46:1150.
96. Isaias IU, Alterman RL, Tagliati M. Deep brain stimulation for primary generalized dystonia: long-term outcomes. *Arch Neurol* 2009; 66:465.
97. Jankovic, J. Tourette Syndrome. UpToDate. UpToDate.com. Waltham, MA. Accessed December 22, 2017.
98. Kang GA, Heath S, Rothlind J, Starr PA. Long-term follow-up of pallidal deep brain stimulation in two cases of Huntington's disease. *J Neurol Neurosurg Psychiatry*. 2011;82(3):272-277.
99. Kasoff WS, Gross RE. Deep brain stimulation: Introduction and technical aspects. In: *Neuromodulation in Psychiatry*, Hamani C, Holtzheimer PH, Lozano A, Mayberg M (Eds), Wiley & Sons, West Sussex 2016. p.245.
100. Kefalopoulou Z, Zrinzo L, Jahanshahi M, et al. Bilateral globus pallidus stimulation for severe Tourette's syndrome: a double-blind, randomised crossover trial. *Lancet Neurol* 2015; 14:595.
101. Kelly K, Theodore WH. Prognosis 30 years after temporal lobectomy. *Neurology*. 2005;64(11):1974-1976.
102. Kennedy SH, Milev R, Giacobbe P, et al; Canadian Network for Mood and Anxiety Treatments (CANMAT). Canadian Network for Mood and Anxiety Treatments (CANMAT) Clinical guidelines for the management of major depressive disorder in adults. IV. Neurostimulation therapies. *J Affect Disord*. 2009;117 Suppl 1:S44-S53.

103. Kerrigan JF, Litt B, Fisher RS, et al. Electrical stimulation of the anterior nucleus of the thalamus for the treatment of intractable epilepsy. *Epilepsia*. 2004;45(4):346-354.
104. Kisely S, Hall K, Siskind D, et al. Deep brain stimulation for obsessive-compulsive disorder: A systematic review and meta-analysis. *Psychol Med*. 2014;44(16):3533-3542.
105. Kiss ZH, Doig-Beyaert K, Eliasziw M, et al. The Canadian multicentre study of deep brain stimulation for cervical dystonia. *Brain* 2007; 130:2879.
106. Kleiner-Fisman G, Liang GS, Moberg PJ, et al. Subthalamic nucleus deep brain stimulation for severe idiopathic dystonia: impact on severity, neuropsychological status, and quality of life. *J Neurosurg* 2007; 107:29.
107. Kohl S, Schönherr DM, Luigjes J, et al. Deep brain stimulation for treatment-refractory obsessive compulsive disorder: A systematic review. *BMC Psychiatry*. 2014;14:214.
108. Koller W, Pahwa R, Busenbark K, et al. High-frequency unilateral thalamic stimulation in the treatment of essential and parkinsonian tremor. *Ann Neurol* 1997; 42:292.
109. Koller WC, Lyons KE, Wilkinson SB, Pahwa R. Efficacy of unilateral deep brain stimulation of the VIM nucleus of the thalamus for essential head tremor. *Mov Disord* 1999; 14:847.
110. Kooshkabadi A, Lunsford LD, Tonetti D, et al. Gamma Knife thalamotomy for tremor in the magnetic resonance imaging era. *J Neurosurg* 2013; 118:713.
111. Koy A, Hellmich M, Pauls KA, et al. Effects of deep brain stimulation in dyskinetic cerebral palsy: A meta-analysis. *Mov Disord*. 2013;28(5):647-654.
112. Krack P, Batir A, Van Blercom N, et al. Five-year follow-up of bilateral stimulation of the subthalamic nucleus in advanced Parkinson's disease. *N Engl J Med*. 2003;349(20):1925-1934.
113. Krishnaiah B, Ramaratnam S, Ranganathan LN. Subpial transection surgery for epilepsy. *Cochrane Database Syst Rev*. 2013;8:CD008153.
114. Kubu CS, Malone DA, Chelune G, et al. Neuropsychological outcome after deep brain stimulation in the ventral capsule/ventral striatum for highly refractory obsessive-compulsive disorder or major depression. *Stereotact Funct Neurosurg* 2013; 91:374.
115. Kumar R, Lozano AM, Kim YJ, et al. Double-blind evaluation of subthalamic nucleus deep brain stimulation in advanced Parkinson's disease. *Neurology*. 1998;51(3):850-855.
116. Kupfer DJ, Frank E, Phillips ML. Major depressive disorder: New clinical, neurobiological, and treatment perspectives. *Lancet*. 2012;379(9820):1045-1055.
117. Kupsch A, Benecke R, Müller J, et al. Pallidal deep-brain stimulation in primary generalized or segmental dystonia. *N Engl J Med* 2006; 355:1978.
118. Laxton AW, Lipsman N, Lozano AM. Deep brain stimulation for cognitive disorders. *Handbook of Clinical Neurology* 2013;116:307-11.
119. Laxton AW, Lozano AM. Deep brain stimulation for the treatment of Alzheimer disease and dementias. *World Neurosurgery* 2013;80(3-4):S28.e1-8.
120. Lemaire JJ, Sontheimer A, Nezzar H, et al. Electrical modulation of neuronal networks in brain-injured patients with disorders of consciousness: A systematic review. *Ann Fr Anesth Reanim*. 2014;33(2):88-97.
121. Leone M, Franzini A, Broggi G, Bussone G. Hypothalamic stimulation for intractable cluster headache: long-term experience. *Neurology* 2006; 67:150.

122. Leone M, Franzini A, D'Andrea G, et al. Deep brain stimulation to relieve drug-resistant SUNCT. *Ann Neurol* 2005; 57:924.
123. Leone M, May A, Franzini A, et al. Deep brain stimulation for intractable chronic cluster headache: proposals for patient selection. *Cephalalgia* 2004; 24:934.
124. Levine CB, Fahrback KR, Siderowf AD, et al. Diagnosis and treatment of Parkinson's Disease: A systematic review of the literature. Evidence Report/Technology Assessment No. 57. Rockville, MD: Agency for Healthcare Research and Quality (AHRQ); 2003.
125. Lim SY, Hodaie M, Fallis M, et al. Gamma knife thalamotomy for disabling tremor: a blinded evaluation. *Arch Neurol* 2010; 67:584.
126. Limousin P, Krack P, Pollak P, et al. Electrical stimulation of the subthalamic nucleus in advanced Parkinson's disease. *N Engl J Med*. 1998;339(16):1105-1111.
127. Limousin P, Pollak P, Benazzouz A, et al. Effect of parkinsonian signs and symptoms of bilateral subthalamic nucleus stimulation. *Lancet*. 1995;345(8942):91-95.
128. Limousin P, Speelman JD, Gielen F, Janssens M. Multicentre European study of thalamic stimulation in parkinsonian and essential tremor. *J Neurol Neurosurg Psychiatry* 1999; 66:289.
129. Limousin-Dowsey P, Pollak P, Van Blercom N, et al. Thalamic, subthalamic nucleus and internal pallidum stimulation in Parkinson's disease. *J Neurol*. 1999;246 (Suppl 2):II42-II45.
130. Lipsman N, Lam E, Volpini M, et al. Deep brain stimulation of the subcallosal cingulate for treatment-refractory anorexia nervosa: 1 year follow-up of an open-label trial. *Lancet Psychiatry* 2017; 4:285.
131. Lipsman N, Schwartz ML, Huang Y, et al. MR-guided focused ultrasound thalamotomy for essential tremor: a proof-of-concept study. *Lancet Neurol* 2013; 12:462.
132. Lipsman N, Woodside DB, Giacobbe P, et al. Subcallosal cingulate deep brain stimulation for treatment-refractory anorexia nervosa: A phase 1 pilot trial. *Lancet*. 2013;381(9875):1361-1370.
133. Litt B. Brain stimulation for epilepsy. *Epilepsy Behav*. 2001;2:S61-S67.
134. Liu C, Wen XW, Ge Y, et al. Responsive neurostimulation for the treatment of medically intractable epilepsy. *Brain Res Bull*. 2013;97:39-47.
135. Loddenkemper T, Pan A, Neme S, et al. Deep brain stimulation in epilepsy. *J Clin Neurophysiol*. 2001;18(6):514-532.
136. Louis ED. Treatment of Medically Refractory Essential Tremor. *N Engl J Med* 2016; 375:792.
137. Lozano AM, Dostrovsky J, Chen R, Ashby P. Deep brain stimulation for Parkinson's disease: Disrupting the disruption. *Lancet Neurol*. 2002;1(4):225-231.
138. Luders H, Hahn J, Lesser RP, et al. Basal temporal subdural electrodes in the evaluation with patients with intractable seizures. *Epilepsia*. 1989;30(2):131-142.
139. Luigjes J, van den Brink W, Feenstra M, et al. Deep brain stimulation in addiction: A review of potential brain targets. *Mol Psychiatry*. 2012;17(6):572-583.
140. Lyons KE, Pahwa R. Deep brain stimulation and tremor. *Neurotherapeutics*. 2008;5(2):331-338.
141. Lyons MK, Birch BD, Hillman RA, et al. Long-term follow-up of deep brain stimulation for Meige syndrome. *Neurosurg Focus*. 2010;29(2):E5.

142. Lyons MK, Dodick DW, Evidente VG. Responsiveness of short-lasting unilateral neuralgiform headache with conjunctival injection and tearing to hypothalamic deep brain stimulation. *J Neurosurg* 2009; 110:279.
143. Mallet L, Polosan M, Jaafari N, et al. Subthalamic nucleus stimulation in severe obsessive-compulsive disorder. *N Engl J Med* 2008; 359:2121.
144. Mallet L, Polosan M, Jaafari N, et al; STOC Study Group. Subthalamic nucleus stimulation in severe obsessive-compulsive disorder. *N Engl J Med*. 2008;359(20):2121-2134.
145. Mantione M, Nieman D, Figee M, et al. Cognitive effects of deep brain stimulation in patients with obsessive-compulsive disorder. *J Psychiatry Neurosci* 2015; 40:378.
146. Marks WA, Honeycutt J, Acosta F Jr, et al. Dystonia due to cerebral palsy responds to deep brain stimulation of the globus pallidus internus. *Mov Disord*. 2011;26(9):1748-1751.
147. Marson A, Ramaratnam S. Epilepsy. In: *BMJ Clinical Evidence*. London, UK: BMJ Publishing Group; updated November 2005.
148. Martin JL, Barbanoj MJ, Schlaepfer TE, et al. Transcranial magnetic stimulation for treating depression. *Cochrane Database Syst Rev* 2002; :CD003493.
149. Matharu MS, Zrinzo L. Deep brain stimulation in cluster headache: hypothalamus or midbrain tegmentum? *Curr Pain Headache Rep* 2010; 14:151.
150. Matharu, MS. Short-lasting unilateral neuralgiform headache attacks: Treatment. UpToDate. UpToDate.com. Waltham, MA. Accessed December 22, 2017.
151. May, A. Cluster headache: Treatment and prognosis. UpToDate. UpToDate.com. Waltham, MA. Accessed December 22, 2017.
152. Miller S, Akram H, Lagrata S, et al. Ventral tegmental area deep brain stimulation in refractory short-lasting unilateral neuralgiform headache attacks. *Brain* 2016; 139:2631.
153. Moro E, Lang AE, Strafella AP, et al. Bilateral globus pallidus stimulation for Huntington's disease. *Ann Neurol* 2004; 56:290.
154. Moro E, LeReun C, Krauss JK, et al. Efficacy of pallidal stimulation in isolated dystonia: a systematic review and meta-analysis. *Eur J Neurol* 2017; 24:552.
155. Morrell M. Brain stimulation for epilepsy: Can scheduled or responsive neurostimulation stop seizures? *Curr Opin Neurol*. 2006;19(2):164-168.
156. Morrell MJ, RNS System in Epilepsy Study Group. Responsive cortical stimulation for the treatment of medically intractable partial epilepsy. *Neurology* 2011; 77:1295.
157. Muller-Vahl KR, Cath DC, Cavanna AE, et al; ESSTS Guidelines Group. European clinical guidelines for Tourette syndrome and other tic disorders. Part IV: Deep brain stimulation. *Eur Child Adolesc Psychiatry*. 2011;20(4):209-217.
158. Naesstrom M, Blomstedt P, Bodlund O. A systematic review of psychiatric indications for deep brain stimulation, with focus on major depressive and obsessive-compulsive disorder. *Nord J Psychiatry*. 2016;70(7):483-491.
159. Narang P, Retzlaff A, Brar K, Lippmann S. Deep brain stimulation for treatment-refractory depression. *South Med J*. 2016;109(11):700-703.

160. Nardone R, et al. Invasive and non-invasive brain stimulation for treatment of neuropathic pain in patients with spinal cord injury: a review. *Journal of Spinal Cord Medicine* 2014;37(1):19-31.
161. Nascimento, FA. Dravet syndrome: Management and prognosis. UpToDate. UpToDate.com. Waltham, MA. Accessed December 22, 2017.
162. Nicholson T, Milne R. Pallidotomy, thalamotomy and deep brain stimulation for severe Parkinson's disease. Development and Evaluation Committee (DEC) Report No. 105. Southampton, UK: Wessex Institute for Health Research and Development (WIHRD), University of Southampton; 1999.
163. Nilsen KE, Cock HR. Focal treatment for refractory epilepsy: Hope for the future? *Brain Res Brain Res Rev.* 2004;44(2-3):141-153.
164. Niranjana A, Kondziolka D, Baser S, et al. Functional outcomes after gamma knife thalamotomy for essential tremor and MS-related tremor. *Neurology* 2000; 55:443.
165. Niranjana A, Raju SS, Kooshkabi A, et al. Stereotactic radiosurgery for essential tremor: Retrospective analysis of a 19-year experience. *Mov Disord* 2017; 32:769.
166. O'Brien M A, Wingerchuk D, Angle P, et al. Management of chronic central neuropathic pain following traumatic spinal cord injury. Evidence Report/Technology Assessment No. 45. Rockville, MD: Agency for Healthcare Research and Quality (AHRQ); 2001.
167. Ohye C, Higuchi Y, Shibasaki T, et al. Gamma knife thalamotomy for Parkinson disease and essential tremor: a prospective multicenter study. *Neurosurgery* 2012; 70:526.
168. Okun MS, Stover NP, Subramanian T, et al. Complications of gamma knife surgery for Parkinson disease. *Arch Neurol* 2001; 58:1995.
169. Olanow CW, Watts RL, Koller WC. An algorithm (decision tree) for the management of Parkinson's disease (2001): Treatment guidelines. *Neurology.* 2001;56(Suppl 5):S1-S88.
170. Ondo W, Almaguer M, Jankovic J, Simpson RK. Thalamic deep brain stimulation: comparison between unilateral and bilateral placement. *Arch Neurol* 2001; 58:218.
171. Ondo W, Jankovic J, Schwartz K, et al. Unilateral thalamic deep brain stimulation for refractory essential tremor and Parkinson's disease tremor. *Neurology* 1998; 51:1063.
172. Ooms P, Mantione M, Figee M, et al. Deep brain stimulation for obsessive-compulsive disorders: long-term analysis of quality of life. *J Neurol Neurosurg Psychiatry* 2014; 85:153.
173. Osorio I, Frei MG, Sunderam S, et al. Automated seizure abatement in humans using electrical stimulation. *Ann Neurol* 2005; 57:258.
174. Ostrem JL, Racine CA, Glass GA, et al. Subthalamic nucleus deep brain stimulation in primary cervical dystonia. *Neurology* 2011; 76:870.
175. Pahwa R, Factor SA, Lyons KE, et al. Practice Parameter: treatment of Parkinson disease with motor fluctuations and dyskinesia (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2006; 66:983.
176. Pahwa R, Factor SA, Lyons KE, et al; Quality Standards Subcommittee of the American Academy of Neurology. Practice Parameter: Treatment of Parkinson disease with motor fluctuations and dyskinesia (an evidence-based review): Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology.* 2006;66(7):983-995.

177. Pahwa R, Lyons KE, Wilkinson SB, et al. Comparison of thalamotomy to deep brain stimulation of the thalamus in essential tremor. *Mov Disord* 2001; 16:140.
178. Pahwa R, Lyons KL, Wilkinson SB, et al. Bilateral thalamic stimulation for the treatment of essential tremor. *Neurology* 1999; 53:1447.
179. Pahwa R, Wilkinson S, Smith D, et al. High-frequency stimulation of the globus pallidus for the treatment of Parkinson's disease. *Neurology*. 1997;49(1):249-253.
180. Pan I, Dendukuri N, McGregor M. Subthalamic deep brain stimulation (DBS): Clinical efficacy, safety and cost compared to medical therapy for the treatment of Parkinson's disease. Report No. 38. Montreal, QC: Technology Assessment Unit of the McGill University Health Centre (MUHC); November 27, 2009.
181. Panov F, Gologorsky Y, Connors G, et al. Deep brain stimulation in DYT1 dystonia: A 10-year experience. *Neurosurgery*. 2013;73(1):86-93; discussion 93.
182. Park HR, Kim IH, Kang H, et al. Nucleus accumbens deep brain stimulation for a patient with self-injurious behavior and autism spectrum disorder: Functional and structural changes of the brain: Report of a case and review of literature. *Acta Neurochir (Wien)*. 2017;159(1):137-143.
183. Parr JR, Green AL, Joint C, et al. Deep brain stimulation in childhood: an effective treatment for early onset idiopathic generalised dystonia. *Arch Dis Child* 2007; 92:708.
184. Pelloux Y, Baunez C. Deep brain stimulation for addiction: why the subthalamic nucleus should be favored. *Current Opinion in Neurobiology* 2013;23(4):713-20.
185. Pepper J, Hariz M, Zrinzo L, et al. Deep brain stimulation versus anterior capsulotomy for obsessive-compulsive disorder: A review of the literature. *J Neurosurg*. 2015;122(5):1028-1037.
186. Pereira EA, Aziz TZ. Neuropathic pain and deep brain stimulation. *Neurotherapeutics* 2014;11(3):496-507.
187. Pereira EA, et al. Thalamic deep brain stimulation for neuropathic pain after amputation or brachial plexus avulsion. *Neurosurgical Focus* 2013;35(3):E7.
188. Peters TE, Bhavaraju NC, Frei MG, Osorio I. Network system for automated seizure detection and contingent delivery of therapy. *J Clin Neurophysiol* 2001; 18:545.
189. Pichon Riviere A, Augustovski F, Cernadas C, et al. Epilepsy surgery [summary]. Report IRR No. 18. Buenos Aires, Argentina: Institute for Clinical Effectiveness and Health Policy (IECS); December 2003.
190. Plaha P, Khan S, Gill SS. Bilateral stimulation of the caudal zona incerta nucleus for tremor control. *J Neurol Neurosurg Psychiatry*. 2008;79(5):504-513.
191. Plow EB, Pascual-Leone A, Machado A. Brain stimulation in the treatment of chronic neuropathic and non-cancerous pain. *J Pain*. 2012;13(5):411-424.
192. Pollo C, Villemure JG. Rationale, mechanisms of efficacy, anatomical targets and future prospects of electrical deep brain stimulation for epilepsy. *Acta Neurochir Suppl*. 2007;97(Pt 2):311-320.
193. Porta M, Brambilla A, Cavanna AE, et al. Thalamic deep brain stimulation for treatment-refractory Tourette syndrome: Two-year outcome. *Neurology*. 2009;73(17):1375-1380.

194. Pretto TE, Dalvi A, Kang UJ, Penn RD. A prospective blinded evaluation of deep brain stimulation for the treatment of secondary dystonia and primary torticollis syndromes. *J Neurosurg* 2008; 109:405.
195. Rabins P, Appleby BS, Brandt J, et al. Scientific and ethical issues related to deep brain stimulation for disorders of mood, behavior, and thought. *Arch Gen Psychiatry*. 2009;66(9):931-937.
196. Reese R, Gruber D, Schoenecker T, et al. Long-term clinical outcome in meige syndrome treated with internal pallidum deep brain stimulation. *Mov Disord* 2011; 26:691.
197. Rehncrona S, Johnels B, Widner H, et al. Long-term efficacy of thalamic deep brain stimulation for tremor: Double-blind assessments. *Mov Disord*. 2003;18(2):163-170.
198. Rojas-Medina LM, Esteban-Fernandez L, Rodríguez-Berrocal V, et al. Deep brain stimulation in posttraumatic tremor: A series of cases and literature review. *Stereotact Funct Neurosurg*. 2016;94(6):379-386.
199. Salanova V, Witt T, Worth R, et al. Long-term efficacy and safety of thalamic stimulation for drug-resistant partial epilepsy. *Neurology* 2015; 84:1017.
200. Sankar T, Lipsman N, Lozano AM. Deep brain stimulation for disorders of memory and cognition. *Neurotherapeutics* 2014;11(3):527-34.
201. Schiff ND. Central thalamic deep brain stimulation to support anterior forebrain mesocircuit function in the severely injured brain. *J Neural Transm (Vienna)*. 2016;123(7):797-806.
202. Schjerling L, Hjermind LE, Jespersen B, et al. A randomized double-blind crossover trial comparing subthalamic and pallidal deep brain stimulation for dystonia. *J Neurosurg* 2013; 119:1537.
203. Schlaepfer TE, Bewernick BH, Kayser S, et al. Deep brain stimulation of the human reward system for major depression--rationale, outcomes and outlook. *Neuropsychopharmacology* 2014; 39:1303.
204. Schoenen J, Di Clemente L, Vandenheede M, et al. Hypothalamic stimulation in chronic cluster headache: a pilot study of efficacy and mode of action. *Brain* 2005; 128:940.
205. Schrader C, Benecke R, Deuschl G, et al; German Deep Brain Stimulation Association. Deep brain stimulation for dystonia. Consensus recommendations of the German Deep Brain Stimulation Association. *Nervenarzt*. 2009;80(6):656-661.
206. Schrader C, Capelle HH, Kiefe TM, et al. GPi-DBS may induce a hypokinetic gait disorder with freezing of gait in patients with dystonia. *Neurology* 2011; 77:483.
207. Schrock LE, Mink JW, Woods DW, et al. Tourette syndrome deep brain stimulation: a review and updated recommendations. *Mov Disord* 2015; 30:448.
208. Schulz-Schaeffer WJ, Margraf NG, Munser S, et al. Effect of neurostimulation on camptocormia in Parkinson's disease depends on symptom duration. *Mov Disord*. 2015;30(3):368-372.
209. Schuurman PR, Bosch DA, Bossuyt PM, et al. A comparison of continuous thalamic stimulation and thalamotomy for suppression of severe tremor. *N Engl J Med* 2000; 342:461.
210. Scottish Intercollegiate Guidelines Network (SIGN). Diagnosis and management of epilepsy in adults. SIGN No. 143. 2018. Available at: https://www.sign.ac.uk/assets/sign143_2018.pdf.

211. Servello D, Zekaj E, Saleh C, et al. Sixteen years of deep brain stimulation in Tourette's syndrome: A critical review. *J Neurosurg Sci.* 2016;60(2):218-229.
212. Sharma M, Deogaonkar M. Deep brain stimulation in Huntington's disease: Assessment of potential targets. *Journal of Clinical Neuroscience* 2015;22(5):812-7.
213. Sherry DD. Complex regional pain syndrome in children. UpToDate [serial online]. Waltham, MA: UpToDate; reviewed October 2012.
214. Siderowf A, Gollump SM, Stern MB, et al. Emergence of complex, involuntary movements after gamma knife radiosurgery for essential tremor. *Mov Disord* 2001; 16:965.
215. Sirven, JI. Evaluation and management of drug-resistant epilepsy. UpToDate. UpToDate.com. Waltham, MA. Accessed December 22, 2017.
216. Sprengers M, Vonck K, Carrette E, et al. Deep brain and cortical stimulation for epilepsy. *Cochrane Database Syst Rev* 2017; 7:CD008497.
217. Starr PA, Turner RS, Rau G, et al. Microelectrode-guided implantation of deep brain stimulators into the globus pallidus internus for dystonia: techniques, electrode locations, and outcomes. *J Neurosurg* 2006; 104:488.
218. Starr PA. Placement of deep brain stimulators into the subthalamic nucleus or Globus pallidus internus: Technical approach. *Stereotact Funct Neurosurg.* 2002;79(3-4):118-145.
219. Stein K. Deep brain stimulation for movement disorders other than Parkinsons disease. STEER: Succint and Timely Evaluated Evidence Reviews. Bazian Ltd., eds. London, UK: Wessex Institute for Health Research and Development, University of Southampton; 2001;1(2):1-10.
220. Suchowersky, O. Huntington disease: Management. UpToDate. UpToDate.com. Waltham, MA. Accessed December 22, 2017.
221. Sun B, Chen S, Zhan S, et al. Subthalamic nucleus stimulation for primary dystonia and tardive dystonia. *Acta Neurochir Suppl* 2007; 97:207.
222. Sun FT, Morrell MJ, Wharen RE Jr. Responsive cortical stimulation for the treatment of epilepsy. *Neurotherapeutics.* 2008;5(1):68-74.
223. Taha JM, Janszen MA, Favre J. Thalamic deep brain stimulation for the treatment of head, voice, and bilateral limb tremor. *J Neurosurg* 1999; 91:68.
224. Tanei T, Kajita Y, Kaneoke Y, et al. Staged bilateral deep brain stimulation of the subthalamic nucleus for the treatment of Parkinson's disease. *Acta Neurochir (Wien).* 2009;151(6):589-594.
225. Tasker RR. Deep brain stimulation is preferable to thalamotomy for tremor suppression. *Surg Neurol* 1998; 49:145.
226. Tellez-Zenteno JF, McLachlan RS, Parrent A, et al. Hippocampal electrical stimulation in mesial temporal lobe epilepsy. *Neurology.* 2006;66(10):1490-1494.
227. The Deep-Brain Stimulation for Parkinson' Disease Study Group. Deep-brain stimulation of the subthalamic nucleus or the pars interna of the globus pallidus in Parkinson's disease. *N Engl J Med.* 2001;345(13):956-963.
228. Theodore WH, Fisher RS. Brain stimulation for epilepsy. *Lancet Neurol.* 2004;3(2):111-118.
229. Timmermann L, Deuschl G, Fogel W, et al; Deep Brain Stimulation Association. Deep brain stimulation for tremor in multiple sclerosis: Consensus recommendations of the German Deep Brain Stimulation Association. *Nervenarzt.* 2009;80(6):673-677.

230. Tronnier VM, Fogel W, Kronenbueger M, et al. Is the medial globus pallidus a site for stimulation or lesioning in the treatment of Parkinson's disease? *Stereotact Funct Neurosurg.* 1997;69(1-4 Pt 2):62-68.
231. Vayssiere N, van der Gaag N, Cif L, et al. Deep brain stimulation for dystonia confirming a somatotopic organization in the globus pallidus internus. *J Neurosurg* 2004; 101:181.
232. Velasco AL, Velasco F, Velasco M, et al. Electrical stimulation of the hippocampal epileptic foci for seizure control: A double-blind, long-term follow-up study. *Epilepsia.* 2007;48(10):1895-1903.
233. Velasco F, Carrillo-Ruiz JD, Brito F, et al. Double-blind, randomized controlled pilot study of bilateral cerebellar stimulation for treatment of intractable motor seizures. *Epilepsia.* 2005;46(7):1071-1081.
234. Velasco F, Velasco M, Jiménez F, et al. Predictors in the treatment of difficult-to-control seizures by electrical stimulation of the centromedian thalamic nucleus. *Neurosurgery* 2000; 47:295.
235. Velasco M, Velasco F, Velasco AL, et al. Subacute electrical stimulation of the hippocampus blocks intractable temporal lobe seizures and paroxysmal EEG activities. *Epilepsia* 2000; 41:158.
236. Vidailhet M, Vercueil L, Houeto JL, et al. Bilateral, pallidal, deep-brain stimulation in primary generalised dystonia: a prospective 3 year follow-up study. *Lancet Neurol* 2007; 6:223.
237. Viswanathan A, Jimenez-Shahed J, Baizabal Carvallo JF, Jankovic J. Deep brain stimulation for Tourette syndrome: target selection. *Stereotact Funct Neurosurg* 2012; 90:213.
238. Volkmann J, Sturm V, Weiss P, et al. Bilateral high-frequency stimulation of the internal globus pallidus in advanced Parkinson's disease. *Ann Neurol.* 1998;44(6):953-961.
239. Volkmann J, Wolters A, Kupsch A, et al. Pallidal deep brain stimulation in patients with primary generalised or segmental dystonia: 5-year follow-up of a randomised trial. *Lancet Neurol* 2012; 11:1029.
240. Vonck K, Boon P, Achten E, et al. Long-term amygdalohippocampal stimulation for refractory temporal lobe epilepsy. *Ann Neurol* 2002; 52:556.
241. Walsh RA, Sidiropoulos C, Lozano AM, et al. Bilateral pallidal stimulation in cervical dystonia: blinded evidence of benefit beyond 5 years. *Brain* 2013; 136:761.
242. Wang J, Chang C, Geng N, et al. Treatment of intractable anorexia nervosa with inactivation of the nucleus accumbens using stereotactic surgery. *Stereotact Funct Neurosurg* 2013; 91:364.
243. Weaver FM, Follett K, Stern M, et al; CSP 468 Study Group. Bilateral deep brain stimulation vs best medical therapy for patients with advanced Parkinson disease: A randomized controlled trial. *JAMA.* 2009;301(1):63-73.
244. Weiner HL, Ferraris N, LaJoie J, et al. Epilepsy surgery for children with tuberous sclerosis complex. *J Child Neurol.* 2004;19(9):687-689.
245. Welter ML, Houeto JL, Thobois S, et al. Anterior pallidal deep brain stimulation for Tourette's syndrome: a randomised, double-blind, controlled trial. *Lancet Neurol* 2017; 16:610.
246. Witjas T, Carron R, Krack P, et al. A prospective single-blind study of Gamma Knife thalamotomy for tremor. *Neurology* 2015; 85:1562.

247. Wu H, Van Dyck-Lippens PJ, Santegoeds R, et al. Deep-brain stimulation for anorexia nervosa. *World Neurosurg.* 2013;80(3-4):S29.e1-e10.
248. Young RF, Jacques S, Mark R, et al. Gamma knife thalamotomy for treatment of tremor: long-term results. *J Neurosurg* 2000; 93 Suppl 3:128.
249. Zangiabadi N, Ladino LD, Sina F, Orozco-Hernández JP, Carter A, Téllez-Zenteno JF. Deep Brain Stimulation and Drug-Resistant Epilepsy: A Review of the Literature. *Front Neurol.* 2019;10:601. doi:10.3389/fneur.2019.00601
250. Zauber SE, Watson N, Comella CL, et al. Stimulation-induced parkinsonism after posteroventral deep brain stimulation of the globus pallidus internus for craniocervical dystonia. *J Neurosurg* 2009; 110:229.
251. Zesiewicz TA, Elble RJ, Louis ED, et al. Evidence-based guideline update: treatment of essential tremor: report of the Quality Standards subcommittee of the American Academy of Neurology. *Neurology* 2011; 77:1752.
252. Zhang JG, Ge Y, Stead M, et al. Long-term outcome of globus pallidus internus deep brain stimulation in patients with Tourette syndrome. *Mayo Clin Proc* 2014; 89:1506.
253. Zhang JG, Zhang K, Wang ZC, et al. Deep brain stimulation in the treatment of secondary dystonia. *Chin Med J (Engl)* 2006; 119:2069.
254. Zittel S, Moll CK, Gulberti A, et al. Pallidal deep brain stimulation in Huntington's disease. *Parkinsonism Relat Disord* 2015; 21:1105.
255. Carnero-Pardo, C. Should the mini-mental state examination be retired?. *Neurología (English Edition).* 2014;29(8):473-481.
256. Creavin ST, Wisniewski S, Noel-Storr AH, Trevelyan CM, Hampton T, Rayment D, Thom VM, Nash KJ E, Elhamoui H, Milligan R, Patel AS, Tsivos DV, Wing T, Phillips E, Kellman SM, Shackleton HL, Singleton GF, Neale BE, Watton ME, Cullum S. Mini-Mental State Examination (MMSE) for the detection of dementia in clinically unevaluated people aged 65 and over in community and primary care populations. *Cochrane Database of Systematic Reviews* 2016, Issue 1. Art. No.: CD011145. DOI: 10.1002/14651858.CD011145.pub2
257. Pollak, P. Deep brain stimulation for Parkinson's disease-patient selection. *Handbook of clinical neurology* Elsevier. 2013;116:97-105.
258. Tsoi, K. K., Chan, J. Y., Hirai, H. W., Wong, S. Y., & Kwok, T. C. Cognitive tests to detect dementia: a systematic review and meta-analysis. *JAMA internal medicine.* 2015;175(9): 1450-1458.
259. Jobst BC, Kapur R, Barkley GL, et al. Brain-responsive neurostimulation in patients with medically intractable seizures arising from eloquent and other neocortical areas. *Epilepsia.* 2017;58:1005-1014.
260. Geller EB, Skarpaas TL, Gross RE, et al. Brain-responsive neurostimulation in patients with medically intractable mesial temporal lobe epilepsy. *Epilepsia.* 2017;58:994-1004.
261. Matias CM, Sharan A, Wu C. Responsive Neurostimulation for the Treatment of Epilepsy. *Neurosurg Clin N Am.* 2019;30:231-242.
262. Skarpaas TL, Jarosiewicz B, Morrell MJ. Brain-responsive neurostimulation for epilepsy (RNS® System). *Epilepsy Res.* 2019;153:68-70.

263. Ma BB, Rao VR. Responsive neurostimulation: Candidates and considerations. *Epilepsy Behav.* 2018;88:388-395.
264. Geller EB. Responsive neurostimulation: Review of clinical trials and insights into focal epilepsy. *Epilepsy Behav.* 2018;88S:11-20.
265. Kogan M, McGuire M, Riley J. Deep Brain Stimulation for Parkinson Disease. *Neurosurg Clin N Am.* 2019;30:137-146.
266. Hayes MT. Parkinson's Disease and Parkinsonism. *Am J Med.* 2019 Mar 16. pii: S0002-9343(19)30235-9.
267. Limousin P, Foltynie T. Long-term outcomes of deep brain stimulation in Parkinson disease. *Nat Rev Neurol.* 2019;15:234-242.
268. Fox SH, Katzenschlager R, Lim SY, Barton B, de Bie RMA, Seppi K, Coelho M, Sampaio C; Movement Disorder Society Evidence-Based Medicine Committee. International Parkinson and movement disorder society evidence-based medicine review: Update on treatments for the motor symptoms of Parkinson's disease. *Mov Disord.* 2018;33:1248-1266.
269. Rughani A, Schwalb JM, Sidiropoulos C, Pilitsis J, Ramirez-Zamora A, Sweet JA, Mittal S, Espay AJ, Martinez JG, Abosch A, Eskandar E, Gross R, Alterman R, Hamani C. Congress of Neurological Surgeons Systematic Review and Evidence-Based Guideline on Subthalamic Nucleus and Globus Pallidus Internus Deep Brain Stimulation for the Treatment of Patients With Parkinson's Disease: Executive Summary. *Neurosurgery.* 2018;82:753-756.
270. Lee PS, Crammond DJ, Richardson RM. Deep Brain Stimulation of the Subthalamic Nucleus and Globus Pallidus for Parkinson's Disease. *Prog Neurol Surg.* 2018;33:207-221.
271. Dietz N, Neimat J. Neuromodulation: Deep Brain Stimulation for Treatment of Dystonia. *Neurosurg Clin N Am.* 2019;30:161-168.
272. Rodrigues FB, Duarte GS, Prescott D, Ferreira J, Costa J. Deep brain stimulation for dystonia. *Cochrane Database Syst Rev.* 2019 Jan 10;1:CD012405.
273. Magown P, Andrade RA, Soroceanu A, Kiss ZHT. Deep brain stimulation parameters for dystonia: A systematic review. *Parkinsonism Relat Disord.* 2018;54:9-16.
274. Lake W, Hedera P, Konrad P. Deep Brain Stimulation for Treatment of Tremor. *Neurosurg Clin N Am.* 2019;30:147-159.
275. Altinel Y, Alkhalfan F, Qiao N, Velimirovic M. Outcomes in Lesion Surgery versus Deep Brain Stimulation in Patients with Tremor: A Systematic Review and Meta-Analysis. *World Neurosurg.* 2019 ;123:443-452.e8.
276. Mendonça MD, Meira B, Fernandes M, Barbosa R, Bugalho P. Deep brain stimulation for lesion-related tremors: A systematic review and meta-analysis. *Parkinsonism Relat Disord.* 2018;47:8-14.
277. Salanova V. Deep brain stimulation for epilepsy. *Epilepsy Behav.* 2018;88S:21-24.
278. U.S. Food & Drug Administration. Medtronic DBS System for Epilepsy - P960009/S219. <https://www.fda.gov/medical-devices/recently-approved-devices/medtronic-dbs-system-epilepsy-p960009s219> Approved April 2018.

Clinical Guideline Revision / History Information

Original Date: 7/31/2018

Reviewed/Revised: 7/23/2019, 7/21/2020, 08/04/2021, 12/01/2021, 07/26/2022, 07/19/2023