

Noninvasive Positive Pressure Ventilation

Disclaimer

Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.

Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.

Noninvasive Positive Pressure Ventilation	1
Summary	2
Definitions	2
Medical Necessity Criteria for Clinical Review	4
Indication-Specific Criteria	4
Continuous Positive Airway Pressure (CPAP) / Bi-level Positive Airway Pressure (BPAP - spontaneous mode, spontaneous/timed mode) / Adaptive Servo-Ventilation (ASV) Devices	4
Home Ventilators	5
PAP Treatment Supplies	5
Masks	5
Tubing	6
Filters	6
Initial Humidifier Request	6
Humidifier Replacement Request	6
Other PAP Equipment	6
Experimental or Investigational / Not Medically Necessary	6
Applicable Billing Codes	7
References	9
Clinical Guideline Revision / History Information	12

Summary

Noninvasive positive pressure ventilation (NPPV) may be used to treat a number of conditions, including sleep-related breathing disorders and chronic lung disease. Sleep-related breathing disorders, such as obstructive sleep apnea, are conditions in which abnormal breathing events during sleep are associated with negative impacts on a person's health. Chronic lung diseases, such as chronic obstructive pulmonary disease (COPD), affect the ability of a person to breathe in oxygen and breathe out carbon dioxide. When needed to treat these conditions, NPPV is accomplished by using positive airway pressure (PAP) devices. PAP devices are noninvasive and improve oxygenation and gas exchange in the lungs by producing airflow through oral, nasal, or combination oronasal masks.

Prior to initiation of PAP therapy, a sleep study (e.g. home sleep test, attended nocturnal polysomnography) coupled with the clinically appropriate lung testing (e.g. arterial blood gas, pulmonary function testing) is required to confirm the diagnosis, level of severity, and most appropriate treatment. The device must also be prescribed by a board-certified pulmonary specialist or board-certified sleep medicine practitioner.

Definitions

"Sleep-related breathing disorders" are conditions in which abnormal breathing events occur during sleep and include:

1. Obstructive sleep apnea (OSA) - a sleep-related breathing disorder that occurs when the muscles relax during sleep, causing soft tissue in the back of the throat to collapse and block the upper airway. This causes a reduced or complete halt in airflow despite an ongoing effort to breathe.
2. Central sleep apnea (CSA) syndromes are caused by post-hyperventilation central apnea, which may be triggered by a variety of clinical conditions, and central apnea secondary to hypoventilation, which has been described with opioid use by the American Academy of Sleep Medicine. These types include:
 - a. Primary central sleep apnea
 - b. CSA due to Cheyne-Stokes breathing pattern
 - c. CSA due to a medical condition (not Cheyne-Stokes)
 - d. CSA due to a drug or substance
 - e. Primary sleep apnea of infancy
3. Complex sleep apnea (i.e., treatment-emergent central sleep apnea) - defined by the Centers for Medicare and Medicaid Services as central apnea specifically identified by the persistence or emergence of central apneas or hypopneas during the initiation of CPAP or an E0470 device even though obstructive respiratory events have been significantly reduced. Complex sleep apnea is distinct as obstructive or mixed apnea is present without treatment and can be worsened during positive airway treatment.
4. Sleep-related hypoventilation syndrome (SRHS) - a syndrome with the presence of elevated PaCO₂ levels either directly (by arterial blood gas measurement) or indirectly (by end-tidal CO₂ or transcutaneous CO₂ measurements) and includes the following disorders:
 - a. Obesity hypoventilation syndrome

- b. Congenital central alveolar hypoventilation syndrome
- c. Late-onset central hypoventilation with hypothalamic dysfunction
- d. Idiopathic central alveolar hypoventilation
- e. Sleep-related hypoventilation due to a medication or substance
- f. Sleep-related hypoventilation due to a medical disorder

“Sleep-study testing” is a diagnostic test that is used to diagnose sleep-related disorders by recording a person’s brain waves, blood oxygen levels, heart rate, and breathing during sleep. Two types of sleep-study tests are recognized in the diagnosis of sleep-disordered breathing:

- 1. “Unattended (home) polysomnography (PSG)“/ home sleep apnea test (HSAT) is a portable sleep study that can be done at home without the need for a technician on-site to monitor data.
- 2. “Attended (facility or laboratory) nocturnal polysomnography“ is a test performed overnight in a sleep lab or facility that is administered and overseen by a technician.

“Apnea-hypopnea index (AHI)” is the number of apneas and hypopneas per hour of sleep recorded during a sleep study. The AHI is commonly used to determine the severity of sleep apnea. An apnea is defined as the cessation of airflow at the nose and mouth for a minimum of 10 seconds. A hypopnea is an abnormal respiratory event lasting ≥ 10 seconds with $\geq 30\%$ reduction in thoracoabdominal movement or airflow as compared to baseline, and with at least a 4% oxygen desaturation from the pre-event baseline.

“Respiratory disturbance index (RDI)” is the number of apneas, hypopneas, and respiratory effort-related arousals (RERAs) per hour of sleep recorded during a sleep study.

“Respiratory event index (REI)” is a measurement of sleep-disordered breathing on home sleep apnea testing, defined as the number of apneas plus hypopneas during the sleep test divided by the total sleep or recording time reported in hours.

“Pulmonary function testing” is noninvasive testing that measures lung capacity, airflow, and chest wall mechanics.

“Positive airway pressure devices” are noninvasive equipment that assist in ventilation by delivering variable pressures of airflow during inspiration and expiration via an oral, nasal, or oronasal mask. They include:

- 1. Bi-level positive airway pressure (BPAP) devices
- 2. Continuous positive airway pressure (CPAP) devices
- 3. Adaptive servo-ventilation (ASV) devices

Medical Necessity Criteria for Clinical Review

Indication-Specific Criteria

Continuous Positive Airway Pressure (CPAP) / Bi-level Positive Airway Pressure (BPAP - spontaneous mode, spontaneous/timed mode) / Adaptive Servo-Ventilation (ASV) Devices

The Plan considers CPAP, BPAP, or ASV devices provided by a network durable medical equipment (DME) provider (unless the member has out-of-network benefits) medically necessary if ONE of the following criteria is met:

1. Initiation of therapy or short-term therapy (up to 90 days) when BOTH of the following criteria are met:
 - a. Prescription from treating licensed provider; *and*
 - b. Meets the diagnostic and severity criteria for a sleep-related breathing disorder or other pulmonary disease as outlined:
 - i. MCG criteria A-0431 for [Continuous Positive Airway Pressure (CPAP) Device] (HCPCS E0601); *or*
 - ii. MCG criteria A-0337 for [CPAP Titration, Home Auto-titrating PAP (APAP)] (HCPCS E0601); *or*
 - iii. MCG criteria A-0338 for [CPAP Titration, Sleep Center] (HCPCS E0601); *or*
 - iv. MCG criteria A-0994 [Bilevel Positive Airway Pressure (BPAP) Device] for
 1. BPAP without a backup respiratory rate (HCPCS E0470); *or*
 2. BPAP with a back-up respiratory rate (HCPCS E0471, E0472); *or*
 - v. Adaptive servo-ventilation (ASV) devices (e.g., BPAP Auto Servo-Ventilation) (HCPCS codes E0471) are considered medically necessary when ALL of the following criteria are met:
 1. Member is diagnosed with complex sleep apnea, i.e., treatment-emergent central sleep apnea; *and*
 2. Member does not have symptomatic chronic heart failure between Class II-IV in New York Heart Association (NYHA) Functional Classification; *and*
 3. Cardiac assessment prior to initiation of treatment shows the member does not have left ventricular ejection fraction $\leq 45\%$ and moderate to severe predominant central sleep apnea; *and*
 4. Member has diagnostic polysomnography (PSG) that shows five or more predominantly obstructive respiratory events (obstructive or mixed apneas, hypopneas or respiratory effort related arousals [RERAs]) per hour of sleep; *and*
 5. PSG during use of positive airway pressure (HCPCS code E0470) without a backup rate shows significant resolution of obstructive events and emergence or persistence of central apnea or central hypopnea with both of the following:
 - a. Central apneas and central hypopneas $\geq 5/\text{hour}$; *and*
 - b. Total number of central apneas and central hypopneas is $>50\%$ of total number of apneas and hypopneas.

2. Authorization for purchase for initial long-term therapy (more than 90 days) when ALL of the following criteria are met:
 - a. Documented symptom and/or AHI/RDI/REI improvement with PAP therapy; *and*
 - b. Documented follow up with prescribing provider within 90 days of initiation; *and*
 - c. Documented compliance with use of PAP device, defined as usage on at least 70% of nights for an average of 4 hours per 24-hour period during a period of 30 consecutive days during the first 90 days after initiation.
3. A replacement device when ALL of the following criteria are met:
 - a. Device is consistently used on at least 70% of nights for an average of 4 hours per 24-hour period; *and*
 - b. Device is not functioning appropriately; *and*
 - c. Device has been evaluated by the supplying DME provider and has been deemed unable to be repaired; *and*
 - d. The device is no longer covered under the manufacturer's warranty.

Note:

- e. Replacement due to misuse or abuse is NOT covered.
- f. Duplicate equipment for traveling is NOT medically necessary and considered a convenience item.

Home Ventilators

For home ventilators that perform noninvasive ventilation (e.g., E0466,E0467), please see MCG criteria.

PAP Treatment Supplies

The Plan considers the following PAP treatment supplies medically necessary when ALL of the following criteria are met:

1. They are provided by an in-network DME provider (unless the member has out-of-network benefits); *and*
2. Ordered via signed prescription by licensed treating practitioners dated within one year of request; *and*
3. Either a CPAP or BiPAP device has been approved based on the medical necessity criteria.

Masks

The Plan considers masks medically necessary when ONE of the following criteria is met:

1. If a member cannot tolerate the initially provided mask type, eligibility for an alternative mask type will be considered if documentation shows the member is compliant with the PAP device but unable to tolerate the mask type; *or*
2. Combination oral/nasal mask (A7027) - 1 per 3 months; *or*
 - a. Oral cushion used with combination oral/nasal mask (A7028) - 2 per month; *or*
 - b. Nasal pillows used with combination oral/nasal mask (A7029) - 2 per month; *or*
3. Full face mask (A7030) - 1 per 3 months; *or*
 - a. Full face mask interface replacement (A7031)- 2 per month; *or*

4. Nasal interface (mask or cannula type) (A7034) - 1 per 3 months; *or*
 - a. Cushion for nasal mask interface (A7032) - 2 per month; *or*
 - b. Nasal pillow for nasal cannula interface (A7033) - 2 per month; *or*
5. Oral interface (A7044) - 1 per 6 months.

Tubing

The Plan considers tubing medically necessary when ONE of the following criteria is met:

1. Tubing with integrated heating element (A4604) - 1 per 3 months; *or*
2. Tubing used with PAP devices (A7037) - 1 per 3 months.

Filters

The Plan considers filters medically necessary when ONE of the following criteria is met::

1. Disposable filter (A7038) - 2 per month; *or*
2. Non-disposable filter (A7039) - 1 per 6 months.

Initial Humidifier Request

The Plan considers an initial request for a heated (E0562) or non-heated (E0561) humidifier medically necessary when ONE of the following criteria is met:

1. No previous humidifier has been provided for initial requests within the plan year.

Humidifier Replacement Request

The Plan considers a replacement request for a heated (E0562) or non-heated (E0561) humidifier medically necessary when BOTH of the following criteria are met:

1. The device is not operating properly; *and*
2. ONE of the following:
 - a. The device has been evaluated by the supplying DME provider and has been deemed unable to be repaired; *or*
 - b. The device is no longer covered under the manufacturer's warranty.

Other PAP Equipment

The Plan considers the following PAP equipment types medically necessary when ONE of the following criteria is met:

1. Headgear (A7035) - 1 per 6 months; *or*
2. Chinstrap (A7036) - 1 per 6 months; *or*
3. Exhalation port with or without swivel (A7045, bundled with mask interfaces) - 1 per 6 months; *or*
4. Water chamber for humidifier (A7046, bundled at the time of initial supply) - 1 per 6 months.

Experimental or Investigational / Not Medically Necessary

The following items are considered comfort or convenience items, or are available over the counter, and are considered not medically necessary:

1. Batteries, SD cards, modems, or other feature components for positive airway pressure devices

2. DC adapters for positive airway pressure devices
3. Electrosleep therapy
4. Cleaning equipment for automatic or manual cleaning of interfaces, tubing, or other related supplies
5. Duplicate equipment for traveling
6. Non-PAP related home humidifiers
7. Positive airway pressure bed pillows
8. The treatment of snoring alone by any method
9. Use of the miscellaneous durable medical equipment code (HCPCS code E1399)

Applicable Billing Codes

Table 1

CPT/HCPCS codes considered medically necessary if criteria are met:

Code	Description
94002	Ventilation assist and management, initiation of pressure or volume preset ventilators for assisted or controlled breathing; hospital inpatient/observation, initial day
94003	Ventilation assist and management, initiation of pressure or volume preset ventilators for assisted or controlled breathing; hospital inpatient/observation, each subsequent day
94004	Ventilation assist and management, initiation of pressure or volume preset ventilators for assisted or controlled breathing; nursing facility, per day
94660	Continuous positive airway pressure ventilation (CPAP), initiation and management
A4604	Tubing with integrated heating element for use with positive airway pressure device
A7027	Combination oral/nasal mask, used with continuous positive airway pressure device, each
A7028	Oral cushion for combination oral/nasal mask, replacement only, each
A7029	Nasal pillows for combination oral/nasal mask, replacement only, pair
A7030	Full face mask used with positive airway pressure device, each
A7031	Face mask interface, replacement for full face mask, each
A7032	Cushion for use on nasal mask interface, replacement only, each
A7033	Pillow for use on nasal cannula type interface, replacement only, pair

Table 1

CPT/HCPCS codes considered medically necessary if criteria are met:

<i>Code</i>	<i>Description</i>
A7034	Nasal interface (mask or cannula type) used with positive airway pressure device, with or without head strap
A7035	Headgear used with positive airway pressure device
A7036	Chinstrap used with positive airway pressure device
A7037	Tubing used with positive airway pressure device
A7038	Filter, disposable, used with positive airway pressure device
A7039	Filter, non-disposable, used with positive airway pressure device
A7044	Oral interface used with positive airway pressure device, each
A7045	Exhalation port with or without swivel used with accessories for positive airway devices, replacement only
A7046	Water chamber for humidifier, used with positive airway pressure device, replacement, each
E0466	Home ventilator, any type, used with noninvasive interface, (e.g., mask, chest shell)
E0467	Home ventilator, multi-function respiratory device, also performs any or all of the additional functions of oxygen concentration, drug nebulization, aspiration, and cough stimulation, includes all accessories, components and supplies for all functions
E0470	Respiratory assist device, bi-level pressure capability, without back-up rate feature, used with noninvasive interface, e.g., nasal or facial mask
E0471	Respiratory assist device, bi-level pressure capability, with back-up rate feature, used with noninvasive interface, e.g., nasal or facial mask (intermittent assist device with continuous positive airway pressure device)
E0472	Respiratory assist device, bi-level pressure capability, with back-up rate feature, used with invasive interface, e.g., tracheostomy tube (intermittent assist device with continuous positive airway pressure device)
E0561	Humidifier, non-heated, used with positive airway pressure device
E0562	Humidifier, heated, used with positive airway pressure device
E0601	Continuous positive airway pressure (CPAP) device
K1027	Oral device/appliance used to reduce upper airway collapsibility, without fixed mechanical hinge, custom fabricated, includes fitting and adjustment

Table 2

CPT/HCPCS codes <u>not</u> considered medically necessary for indications in this guideline:	
Code	Description
A9279	Monitoring feature/device, stand-alone or integrated, any type, includes all accessories, components and electronics, not otherwise classified
E1399	Durable medical equipment, miscellaneous <ul style="list-style-type: none"> • Due to the broad nature of this code and lack of specificity in certain scenarios, clarification is provided below: • When this is billed for PAP cleaning machine, it is considered NOT medically necessary
95999	Unlisted neurological or neuromuscular diagnostic procedure <ul style="list-style-type: none"> • Due to the broad nature of this code and lack of specificity in certain scenarios, clarification is provided below: • When this is billed for PAP electrosleep therapy for OSA, it is considered NOT medically necessary

References

1. Agency for Healthcare Research and Quality (AHRQ). (2018). *Evidence-based practice center systematic review protocol: Home mechanical ventilators*. Project ID: PULT0717. <https://www.ncbi.nlm.nih.gov/books/NBK575994/bin/bm1-et1.pdf>
2. Agency for Healthcare Research and Quality (AHRQ). (2020). *Noninvasive positive pressure ventilation in the home: Final technology assessment*. Project ID: PULT0717 2/4/2020. <https://www.cms.gov/Medicare/Coverage/DeterminationProcess/downloads/id108TA.pdf>
3. American Academy of Sleep Medicine. (n.d.). *Practice guidelines*. <https://aasm.org/clinical-resources/practice-standards/practice-guidelines/>
4. Askland, K., Wright, L., Wozniak, D. R., Emmanuel, T., Caston, J., & Smith, I. (2020). Educational, supportive and behavioural interventions to improve usage of continuous positive airway pressure machines in adults with obstructive sleep apnoea. *Cochrane Database of Systematic Reviews*, 4(4), CD007736. <https://doi.org/10.1002/14651858.CD007736.pub3>
5. Aurora, R. N., Bista, S. R., Casey, K. R., Chowdhuri, S., Kristo, D. A., Mallea, J. M., ... & Heald, J. L. (2016). Updated adaptive servo-ventilation recommendations for the 2012 AASM guideline: The treatment of central sleep apnea syndromes in adults: Practice parameters with an evidence-based literature review and meta-analyses. *Journal of Clinical Sleep Medicine*, 12(5), 757-761
6. Berry, R. B., Abreua, A. R., Krishnan, V., Quan, S. F., Strollo, P. J., & Malhotra, R. K. (2022). A transition to the American Academy of Sleep Medicine-recommended hypopnea definition in

adults: Initiatives of the Hypopnea Scoring Rule Task Force. *Journal of Clinical Sleep Medicine*, 18(5), 1419–1425. <https://doi.org/10.5664/jcsm.9952>

7. Branson, R., Dichter, J. R., Feldman, H., Devereaux, A., Dries, D., Benditt, J., Hossain, T., Ghazipura, M., King, M., Baldissari, M., Christian, M. D., Dominguez-Cherit, G., Henry, K., Martland, A. M. O., Huffines, M., Ornoff, D., Persoff, J., Rodriguez, D., Jr, Maves, R. C., Kissoon, N. T., ... Rubinson, L. (2021). The US Strategic National Stockpile ventilators in coronavirus disease 2019: A comparison of functionality and analysis regarding the emergency purchase of 200,000 devices. *Chest*, 159(2), 634–652. <https://doi.org/10.1016/j.chest.2020.09.085>
8. Chasens, E. R., Korytkowski, M., Burke, L. E., Strollo, P. J., Stansbury, R., Bizhanova, Z., Atwood, C. W., & Sereika, S. M. (2022). Effect of treatment of OSA with CPAP on glycemic control in adults with type 2 diabetes: The Diabetes Sleep Treatment Trial (DSTT). *Endocrine Practice*, 28(4), 364–371. <https://doi.org/10.1016/j.eprac.2022.01.015>
9. Chowdhuri, S., Quan, S. F., Almeida, F., Ayappa, I., Batool-Anwar, S., Budhiraja, R., Cruse, P. E., Drager, L. F., Griss, B., Marshall, N., Patel, S. R., Patil, S., Knight, S. L., Rowley, J. A., Slyman, A., & ATS Ad Hoc Committee on Mild Obstructive Sleep Apnea (2016). An official American Thoracic Society research statement: Impact of mild obstructive sleep apnea in adults. *American Journal of Respiratory and Critical Care Medicine*, 193(9), e37–e54. <https://doi.org/10.1164/rccm.201602-0361ST>
10. Chowdhuri, S., & Parthasarathy, S. (2025). Treatment-emergent central sleep apnea. *UpToDate*. <https://www.uptodate.com/contents/treatment-emergent-central-sleep-apnea>
11. Gottlieb, D. J., & Punjabi, N. M. (2020). Diagnosis and management of obstructive sleep apnea. *JAMA*, 323(14), 1389. <https://doi.org/10.1001/jama.2020.3514>
12. Hayes, Inc. Evidence Analysis Research Brief. *NightBalance (Philips) for the Treatment of Positional Obstructive Sleep Apnea*. Lansdale, PA: Hayes, Inc.; January 2020.
13. Hayes, Inc. Evidence Analysis Research Brief. *Adaptive Servo-Ventilation for Treatment of Central Sleep Apnea*. Hayes, Inc.; November 2020.
14. Homer-Bouthiette, C., & Wilson, K. C. (2025). Noninvasive ventilation in acute asthma exacerbations: A systematic review. *Annals of the American Thoracic Society*, 22(5), 776–784. <https://doi.org/10.1513/AnnalsATS.202407-799OC>
15. Howard, M. E., Piper, A. J., Stevens, B., Holland, A. E., Yee, B. J., Dabscheck, E., Mortimer, D., Burge, A. T., Flunt, D., Buchan, C., Rautela, L., Sheers, N., Hillman, D., & Berlowitz, D. J. (2017). A randomised controlled trial of CPAP versus non-invasive ventilation for initial treatment of obesity hypoventilation syndrome. *Thorax*, 72(5), 437–444. <https://doi.org/10.1136/thoraxjnl-2016-208559>
16. Iftikhar, I. H., Greer, M., Wigger, G. W., & Collop, N. A. (2021). A network meta-analysis of different positive airway pressure interventions in obesity hypoventilation syndrome. *Journal of Sleep Research*, 30(3), e13158. <https://doi.org/10.1111/jsr.13158>
17. Judd, B.G., & Sateia, M.J. (2025). Classification of sleep disorders. *UpToDate*. <https://www.uptodate.com/contents/classification-of-sleep-disorders>
18. Kapur, V. K., Auckley, D. H., Chowdhuri, S., Kuhlmann, D. C., Mehra, R., Ramar, K., & Harrod, C. G. (2017). Clinical practice guideline for diagnostic testing for adult obstructive sleep apnea: An

American Academy of Sleep Medicine clinical practice guideline. *Journal of Clinical Sleep Medicine*, 13(3), 479–504. <https://doi.org/10.5664/jcsm.6506>

19. Macrea, M., Oczkowski, S., Rochwerg, B., Branson, R. D., Celli, B., Coleman, J. M., 3rd, Hess, D. R., Knight, S. L., Ohar, J. A., Orr, J. E., Piper, A. J., Punjabi, N. M., Rahangdale, S., Wijkstra, P. J., Yim-Yeh, S., Drummond, M. B., & Owens, R. L. (2020). Long-term noninvasive ventilation in chronic stable hypercapnic chronic obstructive pulmonary disease. An official American Thoracic Society clinical practice guideline. *American Journal of Respiratory and Critical Care Medicine*, 202(4), e74–e87. <https://doi.org/10.1164/rccm.202006-2382ST>
20. Masa, J. F., Pépin, J. L., Borel, J. C., Mokhlesi, B., Murphy, P. B., & Sánchez-Quiroga, M. Á. (2019). Obesity hypoventilation syndrome. *European Respiratory Review*, 28(151), 180097. <https://doi.org/10.1183/16000617.0097-2018>
21. McEvoy, C. T., MacDonald, K. D., Go, M. A., Milner, K., Harris, J., Schilling, D., Olson, M., Tiller, C., Slaven, J. E., Bjerregaard, J., Vu, A., Martin, A., Mamidi, R., Schelonka, R. L., Morris, C. D., & Tepper, R. S. (2025). Extended continuous positive airway pressure in preterm infants increases lung growth at 6 months: A randomized controlled trial. *American Journal of Respiratory and Critical Care Medicine*. <https://doi.org/10.1164/rccm.202411-2169oc>
22. Mohammadieh, A., Sutherland, K., & Cistulli, P. A. (2017). Sleep disordered breathing: Management update. *Internal Medicine Journal*, 47(11), 1241–1247. <https://doi.org/10.1111/imj.13606>
23. Mokhlesi, B., Won, C. H., Make, B. J., Selim, B. J., Sunwoo, B. Y., & ONMAP Technical Expert Panel (2021). Optimal NIV Medicare access promotion: Patients with hypoventilation syndromes: A technical expert panel report from the American College of Chest Physicians, the American Association for Respiratory Care, the American Academy of Sleep Medicine, and the American Thoracic Society. *Chest*, 160(5), e377–e387. <https://doi.org/10.1016/j.chest.2021.06.083>
24. Moran, F., Bradley, J. M., & Piper, A. J. (2017). Non-invasive ventilation for cystic fibrosis. *The Cochrane Database of Systematic Reviews*, 2(2), CD002769. <https://doi.org/10.1002/14651858.CD002769.pub5>
25. Nowalk, N. C., Neborak, J. M., & Mokhlesi, B. (2022). Is bilevel PAP more effective than CPAP in treating hypercapnic obese patients with COPD and severe OSA? *Journal of Clinical Sleep Medicine*, 18(1), 5–7. <https://doi.org/10.5664/jcsm.9710>
26. Patil, S. P., Ayappa, I. A., Caples, S. M., Kimoff, R. J., Patel, S. R., & Harrod, C. G. (2019). Treatment of adult obstructive sleep apnea with positive airway pressure: An American Academy of Sleep Medicine clinical practice guideline. *Journal of Clinical Sleep Medicine*, 15(2), 335–343. <https://doi.org/10.5664/jcsm.7640>
27. Patil, S. P., Ayappa, I. A., Caples, S. M., Kimoff, R. J., Patel, S. R., & Harrod, C. G. (2019). Treatment of adult obstructive sleep apnea with positive airway pressure: An American Academy of Sleep Medicine systematic review, meta-analysis, and GRADE assessment. *Journal of Clinical Sleep Medicine*, 15(2), 301–334. <https://doi.org/10.5664/jcsm.7638>
28. Perger, E., Inami, T., Lyons, O. D., Alshaer, H., Smith, S., Floras, J. S., Logan, A. G., Arzt, M., Duran Cantolla, J., Delgado, D., Fitzpatrick, M., Fleetham, J., Kasai, T., Kimoff, R. J., Leung, R. S., T., Lorenzi Filho, G., Mayer, P., Mielniczuk, L., Morrison, D. L., Parati, G., ... ADVENT-HF

Investigators (2017). Distinct patterns of hyperpnea during Cheyne-Stokes respiration: Implication for cardiac function in patients with heart failure. *Journal of Clinical Sleep Medicine*, 13(11), 1235–1241. <https://doi.org/10.5664/jcsm.6788>

29. Rudrappa, M., Modi, P., & Bollu, P. C. (2023, July 31). *Cheyne Stokes respirations*. StatPearls - NCBI Bookshelf. <https://www.ncbi.nlm.nih.gov/books/NBK448165>

30. Selim, B., & Ramar, K. (2021). Sleep-related breathing disorders: When CPAP is not enough. *Neurotherapeutics*, 18(1), 81–90. <https://doi.org/10.1007/s13311-020-00955-x>

31. Sheehan, D. W., Birnkrant, D. J., Benditt, J. O., Eagle, M., Finder, J. D., Kissel, J., Kravitz, R. M., Sawnani, H., Shell, R., Sussman, M. D., & Wolfe, L. F. (2018). Respiratory management of the patient with Duchenne muscular dystrophy. *Pediatrics*, 142(Suppl 2), S62–S71. <https://doi.org/10.1542/peds.2018-0333H>

32. Stefan, M. S., Hill, N. S., Raghunathan, K., Liu, X., Pekow, P. S., Memtsoudis, S. G., Ramachandran, S. K., & Lindenauer, P. K. (2016). Outcomes associated with early postoperative noninvasive ventilation in bariatric surgical patients with sleep apnea. *Journal of Clinical Sleep Medicine*, 12(11), 1507–1516. <https://doi.org/10.5664/jcsm.6278>

33. Veasey, S. C., & Rosen, I. M. (2019). Obstructive sleep apnea in adults. *New England Journal of Medicine*, 380(15), 1442–1449. <https://doi.org/10.1056/nejmcp1816152>

34. Whited, L., Hashmi, M. F., & Graham, D. D. (2023, November 5). *Abnormal respirations*. StatPearls - NCBI Bookshelf. <https://www.ncbi.nlm.nih.gov/books/NBK470309/>

35. Wilson, M. E., Dobler, C. C., Morrow, A. S., Beuschel, B., Alsawas, M., Benkhadra, R., Seisa, M., Mittal, A., Sanchez, M., Daraz, L., Holets, S., Murad, M. H., & Wang, Z. (2020). Association of home noninvasive positive pressure ventilation with clinical outcomes in chronic obstructive pulmonary disease. *JAMA*, 323(5), 455. <https://doi.org/10.1001/jama.2019.22343>

Clinical Guideline Revision / History Information

Original Date: 1/26/2017

Reviewed/Revised: 4/11/2017, 7/20/2017, 1/18/2018, 2/5/2019, 1/27/2020, 1/21/2021, 12/1/2021, 01/26/2022, 1/31/2023, 07/19/2023, 01/23/2024, 02/14/2025, 05/01/2026