

Izervay (avancinaptad pegol)

Disclaimer

Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.

Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.

Summary

Age-related macular degeneration (AMD) is a common progressive eye condition and a leading cause of vision loss in those over 65 years old. The most common form is dry AMD, affecting about 90% of AMD patients, where abnormalities in the retinal pigment epithelium and accumulation of drusen deposits in the macula lead to geographic atrophy (GA). GA represents an advanced stage of dry AMD characterized by irreversible loss of retinal cells, leading to progressive vision difficulties like blurred areas or missing spots when reading. It impacts daily functioning and can ultimately cause severe central vision loss or blindness.

Despite the significant burden of GA, effective treatment options have been lacking until recently. On August 4, 2023, the FDA granted approval to Izervay (avancinaptad pegol) as the first C5 complement inhibitor for GA secondary to AMD. As an overactive complement system contributes to AMD pathology, inhibiting C5 may reduce this damaging effect.

1. The monthly intravitreal injections of IZERVAY were studied in two phase 3 trials, GATHER1 and GATHER2, which showed significantly reduced GA lesion growth rates compared to sham injections over 12 months. However, no differences in visual acuity were observed between the treatment and placebo groups.
2. While IZERVAY represents an important advance as the first approved pharmacotherapy for GA, concerns exist regarding its benefit-risk profile. Increased adverse events like endophthalmitis and conversion to wet AMD were seen with IZERVAY versus placebo. Additionally, the lack of efficacy on visual acuity, a key goal in managing AMD, raises uncertainty about its clinical value.
3. Further research is still needed to establish the long-term impact of IZERVAY on vision outcomes and safety.

Definitions

“Age-related macular degeneration (AMD)” is a common progressive eye disease that results in damage to the macula and central vision loss. It has two main forms - dry and wet AMD.

“Drusen” is yellow extracellular deposits that accumulate under the retinal pigment epithelium characterizing early AMD.

“Dry AMD” is the most common form of AMD (~90% cases) characterized by drusen deposits and abnormalities in the retinal pigment epithelium in the macula.

“Endophthalmitis” is a serious inflammation of the interior of the eye that can lead to vision loss.

“Geographic atrophy (GA)” is an advanced form of dry AMD defined by irreversible atrophy and loss of retinal cells leading to progressive vision impairment. Also known as atrophic AMD.

“Idiopathic Polypoidal Choroidal Vasculopathy (IPCV)” is a condition characterized by the presence of polypoidal, saccular dilations of blood vessels located beneath the RPE. It can lead to serous and hemorrhagic detachments of the RPE. While IPCV shares some similarities with nAMD, it's a distinct entity.

“Intravitreal injection” refers to injection of a medication into the vitreous humor of the eye to directly deliver the drug to the retina.

“Macula” is the small central area of the retina responsible for sharp, detailed central vision.

“**Neovascular Age-Related Macular Degeneration (nAMD)**” is the wet form of AMD. It is characterized by the growth of abnormal blood vessels under the retina, which can leak fluid and blood. This leads to rapid and severe central vision loss if not treated.

“**Stargardt Disease**” is an inherited form of juvenile macular degeneration. It typically begins in childhood or adolescence and is caused by mutations in specific genes. Over time, there's a progressive loss of central vision.

Medical Necessity Criteria for Initial Authorization

The Plan considers **Izervay (avacincaptad pegol)** medically necessary when **ALL** of the following criteria are met:

1. The medication is prescribed by or in consultation with an ophthalmologist; **AND**
2. The member is 18 years of age or older; **AND**
3. The member has a diagnosis of geographic atrophy secondary to age-related macular degeneration supported by clinical documentation; **AND**
4. The member does **NOT** have **ANY** of the following:
 - a. active ocular or periocular infections; **or**
 - b. active intraocular inflammation; **AND**
5. Izervay (avacincaptad pegol) is being prescribed at a dose and frequency that is within FDA approved labeling **OR** is supported by compendia or evidence-based published dosing guidelines for the requested indication.

Medical Necessity Criteria for Reauthorization

NOTE: Izervay is approved for administration by intravitreal injection to each affected eye once monthly (approximately every 28 ± 7 days) for up to 12 months.

Reauthorization beyond 12 months will be reviewed on a case-by-case basis and may be granted based on **BOTH** of the following:

1. The medication is prescribed by or in consultation with an ophthalmologist; **AND**
2. Clinical documentation is provided showing **ALL** of the following:
 - a. continued disease stability or slowing in the growth rate of geographic atrophy lesions; **AND**
 - b. no evidence of unacceptable toxicity or adverse events, such as endophthalmitis, retinal detachment, or conversion to wet AMD; **AND**

- c. clinical rationale and supporting evidence for extended Izervay treatment beyond the 12 months initially approved by the FDA.

Experimental or Investigational / Not Medically Necessary

Izervay (avacincaptad pegol) for any other indication or use is considered not medically necessary by the Plan, as it is deemed to be experimental, investigational, or unproven. Non-covered indications include, but are not limited to, the following:

- Neovascular Age-Related Macular Degeneration (nAMD)
- Stargardt Disease
- Idiopathic Polypoidal Choroidal Vasculopathy

Applicable Billing Codes (HCPCS/CPT Codes)

Service(s) name	
CPT/HCPCS Codes considered medically necessary if criteria are met:	
<i>Code</i>	<i>Description</i>
Healthcare Common Procedure Coding System (HCPCS) code	
C9399	Unclassified drugs or biologicals
J3490	Unclassified drug - injection
J3590	Unclassified biologics
Current Procedural Terminology (CPT) codes for imaging	
92134	Optical coherence tomography
92235	Fluorescein angiography (FA)
92250	Fundus photography (Fundus autofluorescence)
92240	Indocyanine green angiography (ICG)
92242	FA and ICG

CPT code for injections	
67028	Intravitreal injection of a pharmacologic agent (separate procedure)
Site modifier	
<i>To specify which eye(s) received the injection, append the appropriate modifier to the end of the CPT code 67028. For example, if the injection was given in the left eye, the code would be 67028-LT.</i>	
LT	Indicates the procedure was performed on the left eye.
RT	Indicates the procedure was performed on the right eye.
50	Indicates the procedure was done on both eyes.
Evaluation and management (E/M) codes	
99203	New Patient Level 3 E/M
99213	Established Patient Level 3 E/M
99204	New Patient Level 4 E/M
99214	Established Patient Level 4 E/M
Office visit modifier	
25	Significant, separately identifiable evaluation and management service by the same physician or other qualified healthcare professional on the same day of the procedure or other services.
ICD-10 codes considered medically necessary if criteria are met:	
<i>Code</i>	<i>Description</i>
H35.3113	Nonexudative age-related macular degeneration, right eye, advanced atrophic without subfoveal involvement
H35.3123	Nonexudative age-related macular degeneration, left eye, advanced atrophic without subfoveal involvement
H35.3133	Nonexudative age-related macular degeneration, bilateral, advanced atrophic without subfoveal involvement
H35.3114	Nonexudative age-related macular degeneration, right eye, advanced atrophic with subfoveal involvement

H35.3124	Nonexudative age-related macular degeneration, left eye, advanced atrophic with subfoveal involvement
H35.3134	Nonexudative age-related macular degeneration, bilateral, advanced atrophic with subfoveal involvement

References

1. American Academy of Ophthalmology. Geographic atrophy. Accessed August 29, 2023. https://eyewiki.aao.org/Geographic_Atrophy
2. Apte R.S., et al.: Age related macular degeneration. N Engl J Med 2021; 385: pp. 539-547.
3. Cabral de Guimaraes TA, Daich Varela M, Georgiou M, Michaelides M: Treatments for dry age-related macular degeneration: therapeutic avenues, clinical trials and future directions. Br J Ophthalmol. 2022 Mar;106(3):297-304. doi: 10.1136/bjophthalmol-2020-318452. Epub 2021 Mar 19.
4. Colijn JM, et al. Enlargement of geographic atrophy from first diagnosis to end of life. JAMA Ophthalmol. 2021;139(7):743-750.
5. Flaxel C.J., et al.: American Academy of Ophthalmology preferred practice patterns: age related macular degeneration. Ophthalmology 2020; 127: pp. P1-P65.
6. Izervay (avacincaptad pegol) [prescribing information]. Parsippany, NJ: Iveric bio, Inc; August 2023.
7. Jaffe GJ, et al. C5 inhibitor avacincaptad pegol for geographic atrophy due to age-related macular degeneration. Ophthalmology. 2021;128(4):576-586.
8. Kim BJ, Mastellos DC, Li Y, Dunaief JL, Lambris JD. Targeting complement components C3 and C5 for the retina: key concepts and lingering questions. Prog Retin Eye Res. 2021;83:100936. doi:10.1016/j.preteyeres.2020.100936
9. Tzoumas N, Riding G, Williams MA, Steel DHW. Complement inhibitors for age-related macular degeneration. Cochrane Database of Systematic Reviews 2023, Issue 6. Art. No.: CD009300. DOI: 10.1002/14651858.CD009300.pub3.

Clinical Guideline Revision / History Information

Original Date: 9/21/2023
Reviewed/Revised: