

## Intraoperative Neuromonitoring

### Disclaimer

*Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.*

*Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.*

### Summary

The Plan members undergoing certain high-risk surgeries may benefit from specialized monitoring, known as intraoperative neuromonitoring (IONM), to help identify and/or prevent damage to critical nerve, spine, and brain structures. IONM is a broad term that includes many different monitoring techniques where a technician or physician observes the function of at risk structures while a surgeon performs the desired procedure. Examples of IONM include brainstem auditory evoked potential (BAEP), visual evoked potential (VEP), electroencephalography (EEG), electromyography (EMG), motor evoked potential (MEP), or sensory evoked potential (SEP). Each of these techniques involves the use of stimuli to a certain part of the nervous system, and then the response is recorded to determine if that specific pathway is functioning properly. Not all surgical procedures require IONM. This guideline describes the criteria and medical necessity for intraoperative neuromonitoring specifically, and does not address non-operative neurologic testing.

### Definitions

"Intraoperative neuromonitoring (IONM)" refers to the use of various electrophysiologic methods of monitoring the function of the brain, spinal cord, and associated nerves during a surgical procedure.

- "Electroencephalography (EEG)" is used to monitor electrical activity of the brain by using small electrodes placed on the scalp. This method can be used as its own form of IONM, but is also integral to detecting the brain response to stimuli in many of the other forms of IONM.

- “Brainstem auditory evoked potential (BAEP)” is a form of IONM used to monitor the function of the pathway from the auditory nerve to the brainstem. It is performed by delivering a loud, repetitive click noise inside the ear(s) and then recording the time it takes for brainstem electrical activity to change using electrodes placed on the scalp.
- “Visual evoked potential (VEP)” is a form of IONM used to monitor the function of the pathway from the eyes to the occipital lobe, the part of the brain responsible for vision. A light stimulus is shined in the eye(s) and the response is recorded with electrodes placed on the scalp.
- “Electromyography (EMG)” is a form of IONM used to monitor the function of skeletal muscles and the nerves controlling them. The nerve to a given muscle can be stimulated and the electrical activity of the target muscle is then measured.
- “Motor evoked potential (MEP)” is a form of IONM used to monitor the function of the motor cortex and outgoing motor tracts. The motor cortex is stimulated, either with direct electrical stimulation or with transcranial magnetic stimulation, and the resulting muscle activity is measured with small electrodes placed on the overlying skin.
- “Sensory/Somatosensory evoked potential (SEP/SSEP)” is a form of IONM used to monitor the function of the sensory cortex and incoming tracts. The target sensory region on the body is stimulated, and the resulting activity in the sensory cortex is recorded with electrodes placed on the scalp and/or spine.

## Clinical Indications

### General Criteria

The Plan requires ALL IONM procedures to meet ALL of the following general criteria to meet medical necessity:

1. Requested by the operating surgeon; *and*
2. The monitoring is performed by a physician (MD or DO) licensed in the state who is trained and experienced in IONM, or a technologist under the supervision of the licensed physician; *and*
3. The monitoring is interpreted by a physician (MD or DO) licensed in the state who is trained and experienced in IONM, who provides real-time supervision and recommendations while monitoring in-person or remotely; *and*
4. The physician monitoring and interpreting is NOT the surgeon; *and*
5. The physician or licensed technician monitors no more than 3 cases simultaneously; *and*
6. The period of neuromonitoring includes intra-operative time only and should not exceed the documented operative period, which does not have to be continuous (e.g., can be 30 minutes of monitoring, with a break, and then another 30 minutes, which would be 1 hour total); *and*
7. Documentation of the monitoring includes ALL of the following:
  - a. The necessity for IONM and justification for the technique(s) used; *and*
  - b. The nerve(s) being tested/monitored; *and*
  - c. The latencies at each testing point captured in software or explained narratively; *and*
  - d. Interpretation of the results as normal or abnormal; *and*
  - e. Any action or intervention taken as a result of the IONM.

### Electroencephalography (EEG)

When the general criteria outlined above are met, the Plan considers IONM using electroencephalogram (EEG) medically necessary when one of the following criteria are met:

1. Intracranial neurovascular surgeries, including but not limited to:
  - a. AV malformation surgery; *or*
  - b. Cerebral vascular aneurysm; *or*
2. Intracranial tumors:
  - a. Supratentorial tumor resection; *or*
3. The following vascular surgeries:
  - a. Arteriography requiring a temporary occlusion of the carotid artery; *or*
  - b. Circulatory arrest requiring hypothermia (*note*: circulatory bypass surgeries such as CABG do not fall under this criteria); *or*
  - c. Surgery of the thoracic aorta, aortic arch, aortic branching vessels, or carotid vessel surgeries when there is risk of cerebral ischemia.

### Brainstem auditory evoked potential (BAEP)

When the general criteria outlined above are met, the Plan considers IONM using brainstem auditory evoked potentials (BAEP) medically necessary for any one of the following indications:

1. Cerebral vascular surgery; *or*
2. Chiari malformation surgery; *or*
3. Microvascular cranial nerve decompression when performed via intracranial posterior fossa approach; *or*
4. Resection of chordoma; *or*
5. Odontoidectomy; *or*
6. Decompression of tumor from anterior brainstem or tumor above C2 in the spinal cord; *or*
7. Compressive brainstem tumor removal; *or*
8. Acoustic neuroma surgery

### Electromyography (EMG)

When the general criteria outlined above are met, the Plan considers IONM using electromyography (EMG) medically necessary for any **ONE** of the following indications:

- Procedures involving the facial nerve, including:
  - Microvascular decompression for hemifacial spasm; *or*
  - Acoustic neuroma surgery; *or*
  - Congenital auricular lesions; *or*
  - Skull base lesions; *or*
  - Surgery for cholesteatoma; *or*
  - Surgical removal of facial nerve neuroma; *or*
  - Vestibular neurectomy for Meniere's disease; *or*

- Selective dorsal rhizotomy; *or*
- Excision of neuromas involving **ONE** of the following:
  - Oculomotor nerve (CN 3); *or*
  - Trochlear nerve (CN 4); *or*
  - Abducens nerve (CN 6); *or*
  - Glossopharyngeal nerve (CN 9); *or*
  - Spinal accessory (CN 11); *or*
  - Hypoglossal nerve (CN 12); *or*
  - Recurrent laryngeal nerve; *or*
  - Superior laryngeal nerve; *or*
- Intraoperative identification of **ONE** of the following nerves during high-risk skull base, posterior fossa, or brainstem surgeries:
  - Oculomotor nerve (CN 3); *or*
  - Trochlear nerve (CN 4); *or*
  - Abducens nerve (CN 6); *or*
  - Glossopharyngeal nerve (CN 9); *or*
  - Spinal accessory (CN 11); *or*
  - Hypoglossal nerve (CN 12)
- Intraoperative monitoring of the recurrent laryngeal nerve during:
  - Anterior neck surgery; *or*
  - Thyroid or parathyroid surgery

#### Sensory/Somatosensory Evoked Potential (SEP/SSEP)

When the general criteria outlined above are met, the Plan considers IONM using SEP/SSEP with or without motor evoked potentials (MEP) medically necessary for any one of the following indications:

1. The following spinal surgeries:
  - a. Surgery for scoliosis or other significant spinal deformity requiring traction of the spinal cord; *or*
  - b. Removal of spinal cord tumors or tumors causing cord compression, including intramedullary tumors; *or*
  - c. Traumatic injury to the spinal cord requiring surgery; *or*
  - d. AV malformation of the spinal cord; *or*
  - e. Cervical decompression if myelopathy is present; *or*
  - f. Spinal corpectomy if myelopathy is present; *or*
2. The following intracranial surgeries:
  - a. Chiari malformation surgery; *or*
  - b. Cerebral vascular aneurysms; *or*
  - c. Deep brain stimulation surgery, such as for Parkinson's disease; *or*
  - d. Endolymphatic shunt for Meniere's disease; *or*

- e. Microvascular decompression of cranial nerves or removal of tumors involving cranial nerves; *or*
  - f. Skull base or cavernous sinus tumor removal; *or*
  - g. Oval or round window graft; *or*
  - h. Resection of brain tissue near the primary motor cortex; *or*
  - i. Resection of epileptogenic brain tissue or tumor; *or*
  - j. Intracranial AV malformation surgery; *or*
  - k. Surgery for movement disorders; *or*
  - l. Vestibular section; *or*
3. The following vascular surgeries:
- a. Arteriography requiring a temporary occlusion of the carotid artery; *or*
  - b. Circulatory arrest requiring hypothermia (*note*: circulatory bypass surgeries such as CABG do not fall under this criteria); *or*
  - c. Distal aortic procedures where there is risk of spinal cord ischemia; *or*
  - d. Surgery of the thoracic aorta, aortic arch, aortic branching vessels, or carotid vessel surgeries when there is risk of cerebral ischemia

*High-risk surgeries may also necessitate multiple IONM modalities. The specific IONM modalities used should be appropriate for the planned surgical intervention and are each subject to review.*

#### Experimental or Investigational / Not Medically Necessary

IONM with the following procedures or indications have limited available data and/or have not shown a benefit in the available literature and are therefore considered experimental, investigational, or unproven:

- ANY procedure or indication not meeting the above inclusion criteria.
- IONM for pain management procedures, including but not limited to:
  - Epidural steroid injections
  - Radiofrequency ablation
  - Medial branch block
  - Facet joint injections
  - Selective nerve root block
- IONM with a technician alone (e.g., no physician), or automated IONM.
- ANY intraoperative spinal monitoring for surgeries below the level of the spinal cord, which typically ends at L1-L2 in most adult patients, with the exception of patients at risk for spinal cord injury due to spinal abnormality such as tethered cord or Chiari malformation.
- Intraoperative brainstem auditory evoked potentials (BAEP) for conductive hearing loss surgeries, including stapedectomy/stapedotomy, tympanoplasty, and ossicular reconstruction.
- Visual evoked potentials (VEP) for ANY intraoperative indication.
- Intraoperative motor evoked potentials (MEP) in ANY of the following situations:
  - When performed using transcranial magnetic stimulation

- For spinal cord stimulator placement
- When used without SSEP
- Intraoperative electromyography (EMG) for ANY of the following situations:
  - When used in combination with a complete neuromuscular blockade for anesthesia, as functioning neuromuscular junction is required for EMG monitoring
  - When performed as intraoperative surface EMG
  - Facial nerve (CN 7) monitoring for any of the following indications:
    - Cochlear implant surgery
    - Parotid gland surgery
    - Tympanoplasty without mastoidectomy
    - Maxillofacial surgery
  - Trigeminal nerve (CN 5) monitoring in:
    - Decompression
    - Neurectomy
    - Radiosurgery
    - Rhizotomy
  - Monitoring of peripheral nerves
  - Aortic aneurysm repair
  - Lower extremity joint surgeries, including but not limited to hip dysplasia surgery or hip replacement, knee arthroplasty, or total knee replacement.
  - Prostatectomy
  - Rectal cancer surgery
  - Rotator cuff surgery
  - Tibial neurectomy
  - Wrist arthroscopy
  - Dorsal column stimulator placement
  - Monitoring of the recurrent laryngeal nerve during anterior cervical spine procedures
  - Adjustment of vertical expandable prosthetic titanium rib (VEPTR)
  - Spinal procedures, including decompressive surgery, discectomy, fusion or other type of invasive spinal surgery.
  - Intracranial tumor surgeries; brainstem and motor-strip mapping with EMG
- Intraoperative SSEP for the following indications:
  - Monitoring the femoral nerve during transpsoas lumbar lateral interbody fusion
  - Monitoring the facial nerve during:
    - Submandibular gland excision
    - Parotid gland surgery
  - Lower extremity joint surgeries, including but not limited to hip dysplasia surgery or hip replacement, knee arthroplasty, or total knee replacement..
  - Implantation of a spinal cord stimulator
  - Off-pump coronary artery bypass surgery
  - Thyroid surgery and parathyroid surgery

- Cochlear implantation
- Monitoring spinal injections (e.g., facet joint, interlaminar and transforaminal epidural)
- Wrist arthroscopy repair or other surgical intervention for carpal tunnel or ulnar nerve entrapment
- Prostate surgery
- Pectus excavatum surgery
- IONM for uncomplicated single level spinal procedures, including:
  - ACDF (anterior cervical discectomy and fusion)
  - Lumbar fusion
  - Lumbar discectomy
  - Spinal decompressive surgery without myelopathy
  - Spinal corpectomy without myelopathy

#### Evidence for Experimental or Investigational or Not Medically Necessary

##### *Intraoperative Brainstem auditory evoked potentials (BAEP) for conductive hearing loss surgery:*

There is limited data demonstrating a potential benefit of BAEP for this indication. The data is primarily in the form of small, single institution, retrospective studies. One study by Hsu (2011), looked at 32 consecutive patients undergoing laser stapedotomy and determined that 23% of patients undergoing the procedure had intraoperative adjustments made. Another study by Selesnick et al in 1997 found a correlation between intraoperative BAEP and post-operative hearing outcomes. However, the size and number of existing studies limit conclusions regarding the clinical utility and benefit for this indication.

##### *Intraoperative Visual evoked potentials (VEP)*

VEPs have been historically used with variable success for the monitoring of lesions near the optic chiasm. However, the literature has shown that interpreting these signals may be difficult and subjective. A 2017 meta-analysis by Metwali et al demonstrated that intraoperative VEPs have a high predictive value but low sensitivity. Other studies have shown that VEPs are only feasible in 70-80% of patients. An UpToDate review on the topic states, "Usefulness of VEP monitoring has not been established, and concerns have been raised that traditional stimulation methods do not produce responses that follow the pathway of useful vision. Also, VEPs are susceptible to effects from general anesthetics, and technical problems with stimulators make monitoring difficult." Further evidence regarding the clinical utility and potential benefit of intraoperative VEP is needed.

##### *Intraoperative Motor evoked potentials (MEP) with transcranial magnetic stimulation (TMS)*

Most procedures with MEP are performed using electrical, rather than magnetic stimulation. The existing literature for magnetic stimulation is limited to case reports and small retrospective series, and IONM using TMS has not yet received FDA approval.

##### *EMG when performed as intraoperative surface EMG*

Surface EMG has not yet been fully studied in the intraoperative setting and has not been approved for this indication, nor is it recommended by any consensus guidelines.

#### *Intraoperative EMG of the facial nerve for cochlear implant surgery*

Alzhrani et al (2016) evaluated rates of facial nerve palsy following cochlear implantation in 3403 surgeries, concluding that “Cochlear implantation entails only a minimal risk of FN palsy and that FN palsy is chiefly a transient problem”. Other studies have found rates <1% of facial nerve palsy and felt that incidence was related to heating injury or viral reactivation. Given the rarity of this finding and the lack of data using IONM with EMG for this indication, further data is needed.

#### *Intraoperative EMG or SSEP of the facial nerve for parotid gland surgery*

A 2009 study by Grosheva et al prospectively looked at patients with EMG vs. no EMG to the facial nerve receiving parotid surgery, finding no difference in immediate post-operative outcome or permanent facial nerve function. EMG significantly reduced the duration of OR time for superficial parotidectomy. Another study by Shan et al in 2014 found no difference in the incidence of post-op facial paralysis in patients monitoring with EMG vs. those who were not, but found reduced surgical time (around 60 minutes) in patients with revision surgery who underwent IONM with EMG to the facial nerve. Further evidence is needed to confirm a potential benefit in clinical outcomes for this indication.

#### *EMG for intraoperative monitoring of the trigeminal nerve*

A 2004 study by Brock et al looked at IONM with EMG and BAEP for trigeminal neuralgia surgical decompression, and concluded “There were no correlations between the entity of the intraoperative EMG discharges and the postoperative facial and trigeminal function.” Further evidence is needed to confirm if this technique should be used for this indication.

#### *EMG for intraoperative monitoring of peripheral nerves*

There has been no prospective or randomized data suggesting that IONM with EMG for surgical procedures involving peripheral nerves has any benefit or improvement in clinical outcomes. Further research is needed to define the role of EMG in this setting.

*EMG for intraoperative monitoring in any of the following settings has not been studied or there is not a clear, documented benefit in the literature:*

- *Aortic aneurysm repair*
- *Hip dysplasia surgery or hip replacement:* Data is limited to a small, 12 patient study showing a single patient where persistent postoperative muscle weakness may have been prevented.<sup>113</sup>
- *Prostatectomy:* There are current clinical trials recruiting patients for intraoperative EMG with robot assisted radical prostatectomy, however, the current evidence is limited for this indication.<sup>55</sup>
- *Rectal cancer surgery:* Recent studies have begun examining intraoperative EMG for low anterior resection in rectal cancer (both robotic and non-robotic surgeries). Kauff et al conducted a randomized trial in 2016 (NEUROS) looked at “Continuous intraoperative monitoring of pelvic



autonomic nerves during TME to prevent urogenital and anorectal dysfunction in rectal cancer patients”, however, the results have yet to be published.

- *Rotator cuff surgery*
- *Tibial neurectomy*
- *Wrist arthroscopy*

#### *EMG monitoring of the recurrent laryngeal nerve anterior cervical spine surgery:*

A study by Jellish et al (1999) looked at rates of adverse events involving the recurrent laryngeal nerve after anterior cervical spine surgeries, finding 38% experienced hoarseness and 15% had “severe” side effects. They found that a greater number of EMG activations was associated with adverse events; however, this study was limited by its non-randomized nature and lack of a control group to determine statistical differences. In 2009, Dimopoulos et al conducted a prospective study on 298 patients undergoing anterior cervical discectomy and fusion (ACDF) to evaluate the role of EMG in predicting post-operative outcomes. They found “Postoperative RLN injury occurred in 2.3% of our patients. The sensitivity of IEMG was 100%, the specificity 87%, the positive predictive value 16%, and its negative predictive value 97%.” The low PPV and high NPV indicate that EMG may be a good tool at excluding injury to the RLN but is not a good predictor of injury in patients with positive EMG findings, limiting clinical utility. Another study by Chen et al (2014) looked at almost 2000 patients and found that only 0.16% of all patients undergoing anterior c-spine surgery experience long-term (>6 months) symptoms as a result of recurrent laryngeal injury, indicating that monitoring in this population may not be necessary. Further randomized, clinical evidence is required to determine the utility of this technique.

#### *EMG monitoring for spinal procedures*

A 2017 study by Ajiboye et al looked at almost 10,000 pedicle screw placements, finding no significant difference in outcomes between surgeries with EMG and those without. Another study by the same group looked at >26,000 patients undergoing anterior cervical spine surgeries, and concluded that “for ACDFs, there is no difference in the risk of neurological injury with or without ION use.” At the present time, there is insufficient evidence for EMG use in spinal procedures.

#### *IONM for adjustment of vertical expandable prosthetic titanium rib (VEPTR):*

A 2009 study by Skaggs et al examined 1736 VEPTR procedures for the efficacy of IONM. of the 1736 procedures, only 8 (0.5%) demonstrated any perioperative neurologic injury and only 1 of these 8 was permanent. The use of IONM in this setting should be further examined to determine whether there is a true clinical benefit to the procedure as the prevalence of injury during this surgery may be exceptionally low.

#### *SSEP for intraoperative monitoring of the femoral nerve during transposas lumbar lateral interbody fusion*

A study by Silverstein et al in 2014 looked at 41 consecutive procedures and found 3/41 patients with post-operative symptoms after 5/41 had intraoperative SSEP findings, and no patients with normal SSEP

findings experienced a deficit. Despite these findings, further data is needed to confirm any potential benefit for this indication.

*SSEP for intraoperative spinal cord monitoring for cervical, lumbar, or thoracic laminectomy or fusion*

As per 2019 systematic review by Martino et al., there were limitations in this study on the outcomes, sensitivity, and specificity of SSEP for degenerative cervical spinal stenosis. Furthermore, there were no RCTs available and the risk of bias were moderate to severe among prospective case studies, case studies, and retrospective case studies.

Overall, for lumbar, thoracic or thoracolumbar spinal procedures, there are limited studies available or outdated more than 10 years. Furthermore, as per 2020 narrative review, there are no prospective studies validating the efficacy of IONM. There are retrospective reviews supporting SEP/SSEP for cervical spinal procedures and cervical pathologies. As per 2023 Basu & Gohil, there were no prospective randomized control trials conducted for intraoperative neuromonitoring in spine surgery. Cervical spinal procedures without myelopathy is considered experimental, investigational, or unproven. The 2018 AANS/CNS Joint Section on Disorders of the Spine and Peripheral Nerves Updated Position Statement: Intraoperative Electrophysiological Monitoring states there is Level I evidence that IOM is a reliable diagnostic tool for assessment of spinal cord integrity during surgery. There is insufficient evidence (Level III) of a therapeutic benefit of IOM during spinal surgery. While IOM is generally regarded as integral to lateral spine surgery, there is insufficient evidence to support a therapeutic benefit.

As per 2017 American Association of Neurological Surgeons (AANS)/Congress of Neurological Surgeons (CNS), Multimodality intraoperative monitoring (MIOM), including somatosensory evoked potentials (SSEPs) and motor evoked potentials (MEPs) recording, during spinal cord/spinal column surgery is a reliable and valid diagnostic adjunct to assess spinal cord integrity and is recommended if utilized for this purpose. However, intraoperative monitoring for preventive or therapeutic benefit on improving postoperative neurological outcomes does not have sufficient evidence.

According American Association of Neuromuscular & Electrodiagnostic Medicine practice guidelines there is no proven benefit and American Clinical Neurophysiology Society does not have guidance on SSEP. In 2017, a retrospective case-control study of 121 patients who underwent spinal cord procedures with the combination of intraoperative neuromonitoring was found to have low sensitivity, false negatives, and poor positive predictive value of neurological injury.

*IONM for uncomplicated single level spinal procedures:*

Hayes, Inc., (2023) provides a D2 rating for intraoperative neuromonitoring (IONM) for detection and prevention of nerve damage due to lumbar spinal discectomy or fusion surgery in adult patients based on the available body of evidence reflecting very low quality and insufficient to draw evidence. Cole et al (2014) conducted a retrospective review of 85,640 patients undergoing single-level spinal procedures. 12.66% of the reviewed procedures utilized intraoperative neuromonitoring.

“Lumbar laminectomies had reduced 30-day neurological complication rate with neuromonitoring (0.0% vs. 1.18%, P=0.002). Neuromonitoring did not correlate with reduced intraoperative neurological complications in ACDFs (0.09% vs. 0.13%), lumbar fusions (0.32% vs. 0.58%), or lumbar discectomy (1.24% vs. 0.91%)”. The authors concluded that “with intraoperative neurological monitoring in single-level procedures, neurological complications were decreased only among lumbar laminectomies. No difference was observed in ACDFs, lumbar fusions, or lumbar discectomies”. Another retrospective study by Ajiboye et al (2017) looked at 15,395 patients undergoing an ACDF, where patients received IONM with “SSEPs only (48.7%), MMEPs only (5.3%), and combined SSEPs and MMEPs (46.1%). Neurological injuries occurred in 0.23% and 0.27% of patients with and without ION, respectively (P=0.84).” The authors concluded that “Use of ION does not further prevent the rate of postoperative neurological complications for ACDFs as compared with the cases without ION. The utility of routine ION for ACDFs is questionable.”

#### Applicable Billing Codes (HCPCS/CPT Codes)

<i>Intraoperative Neuromonitoring</i>	
CPT/HCPCS Codes considered medically necessary if criteria are met:	
<i>Code</i>	<i>Description</i>
92652	Auditory evoked potentials; for threshold estimation at multiple frequencies, with interpretation and report
92653	Auditory evoked potentials; neurodiagnostic, with interpretation and report
95829	Electrocorticogram at surgery (separate procedure)
95865	Needle measurement and recording of electrical activity of muscles of voice box
95867	Needle electromyography; cranial nerve supplied muscle(s), unilateral
95868	Needle electromyography; cranial nerve supplied muscles, bilateral
95925	Short-latency somatosensory evoked potential study, stimulation of any/all peripheral nerves or skin sites, recording from the central nervous system; in upper limbs
95926	Short-latency somatosensory evoked potential study, stimulation of any/all peripheral nerves or skin sites, recording from the central nervous system; in lower limbs
95927	Short-latency somatosensory evoked potential study, stimulation of any/all peripheral nerves or skin sites, recording from the central nervous system; in upper and lower limbs
95938	Short-latency somatosensory evoked potential study, stimulation of any/all peripheral nerves or skin sites, recording from the central nervous system; in upper and lower limbs

95940	Continuous intraoperative neurophysiology monitoring in the operating room, one on one monitoring requiring personal attendance, each 15 minutes (List separately in addition to code for primary procedure)
95941	Continuous intraoperative neurophysiology monitoring, from outside the operating room (remote or nearby) or for monitoring of more than one case while in the operating room, per hour (List separately in addition to code for primary procedure)
95955	Electroencephalogram (EEG) during nonintracranial surgery (eg, carotid surgery)
95999	Unlisted neurological or neuromuscular diagnostic procedure <ul style="list-style-type: none"> <li>• <u>Due to the broad nature of this code and lack of specificity in certain scenarios, clarification is provided below:</u></li> <li>• When this code is billed for Train of four testing which allows measurement of the extent of the NMB, it is medically necessary. The measurement of the degree of neuromuscular blockade, which may be integral to the procedure, is not separately reimbursable.</li> </ul>
G0453	Continuous intraoperative neurophysiology monitoring, from outside the operating room (remote or nearby), per patient, (attention directed exclusively to one patient) each 15 minutes (list in addition to primary procedure)
ICD-10 codes considered medically necessary if criteria are met:	
<i>Code</i>	<i>Description</i>
C70.1	Malignant neoplasm of spinal meninges
C71.7	Malignant neoplasm of brain stem
C71.9	Malignant neoplasm of brain, unspecified
C72.0	Malignant neoplasm of spinal cord
C72.20 - C72.22	Malignant neoplasm of olfactory nerve
C72.30 - C72.32	Malignant neoplasm of optic nerve
C72.40 - C72.42	Malignant neoplasm of acoustic nerve
C72.50 - C72.59	Malignant neoplasm of other and unspecified cranial nerves
C73	Malignant neoplasm of thyroid gland
C79.31 - C79.32	Secondary malignant neoplasm of brain and cerebral meninges
D32.1	Benign neoplasm of spinal meninges
D33.0	Benign neoplasm of brain, supratentorial
D33.1	Benign neoplasm of brain, infratentorial
D33.2	Benign neoplasm of brain, unspecified

D33.3	Benign neoplasm of cranial nerves
D33.4	Benign neoplasm of spinal cord
D42.0 - D42.9	Neoplasm of uncertain behavior of meninges
D43.0 - D43.9	Neoplasm of uncertain behavior of brain and central nervous system
D49.6	Neoplasm of unspecified behavior of brain
D49.7	Neoplasm of unspecified behavior of endocrine glands and other parts of nervous system
E04.2	Nontoxic multinodular goiter
I60.00 - I60.02	Nontraumatic subarachnoid hemorrhage from carotid siphon and bifurcation
I60.10 - I60.12	Nontraumatic subarachnoid hemorrhage from middle cerebral artery
I60.30 - I60.32	Nontraumatic subarachnoid hemorrhage from posterior communicating artery
I60.4	Nontraumatic subarachnoid hemorrhage from basilar artery
I60.50 - I60.52	Nontraumatic subarachnoid hemorrhage from vertebral artery
I60.6	Nontraumatic subarachnoid hemorrhage from other intracranial arteries
I60.7	Nontraumatic subarachnoid hemorrhage from unspecified intracranial artery
I60.8	Other nontraumatic subarachnoid hemorrhage
I60.9	Nontraumatic subarachnoid hemorrhage, unspecified
I65.21 - I65.29	Occlusion and stenosis of carotid artery
I67.1	Cerebral aneurysm, nonruptured
M41.00 - M41.08	Infantile idiopathic scoliosis
M41.112 - M41.119	Juvenile idiopathic scoliosis
M41.122 - M41.129	Adolescent idiopathic scoliosis
M41.20 - M41.27	Other idiopathic scoliosis
M41.30 - M41.35	Thoracogenic scoliosis
M41.40 - M41.47	Neuromuscular scoliosis

M41.50 - M41.57	Other secondary scoliosis
M41.80 - M41.87	Other forms of scoliosis
M41.9	Scoliosis, unspecified
M43.21	Fusion of spine, occipito-atlanto-axial region
M43.22	Fusion of spine, cervical region
M43.23	Fusion of spine, cervicothoracic region
M43.8X1	Other specified deforming dorsopathies, occipito-atlanto-axial region
M43.8X2	Other specified deforming dorsopathies, cervical region
M43.8X3	Other specified deforming dorsopathies, cervicothoracic region
M47.11	Other spondylosis with myelopathy, occipito-atlanto-axial region
M47.12	Other spondylosis with myelopathy, cervical region
M47.13	Other spondylosis with myelopathy, cervicothoracic region
M47.22	Other spondylosis with radiculopathy, cervical region
M47.811	Other spondylosis with myelopathy, occipito-atlanto-axial region
M47.812	Other spondylosis with myelopathy, cervical region
M47.813	Other spondylosis with myelopathy, cervicothoracic region
M47.892	Other spondylosis, cervical region
M48.02	Spinal stenosis, cervical region
M50.00	Cervical disc disorder with myelopathy, unspecified cervical region
M50.02	Cervical disc disorder with myelopathy, mid-cervical region
M50.022	Cervical disc disorder at C5-C6 level with myelopathy
M50.023	Cervical disc disorder at C6-C7 level with myelopathy
M50.10	Cervical disc disorder with radiculopathy, unspecified cervical region
M50.122	Cervical disc disorder at C5-C6 level with radiculopathy
M50.123	Cervical disc disorder at C6-C7 level with radiculopathy
M50.20	Other cervical disc displacement, unspecified cervical region
M50.22	Other cervical disc displacement, mid-cervical region

M50.221	Other cervical disc displacement at C4-C5 level
M50.222	Other cervical disc displacement at C5-C6 level
M50.223	Other cervical disc displacement at C6-C7 level
M50.30	Other cervical disc degeneration, unspecified cervical region
M50.322	Other cervical disc degeneration at C5-C6 level
M50.323	Other cervical disc degeneration at C6-C7 level
M50.820	Other cervical disc disorders, mid-cervical region, unspecified level
M50.821	Other cervical disc disorders at C4-C5 level
M50.822	Other cervical disc disorders at C5-C6 level
M50.823	Other cervical disc disorders at C6-C7 level
M50.83	Other cervical disc disorders, cervicothoracic region
M54.12	Radiculopathy, cervical region
Q04.4	Septo-optic dysplasia of brain
Q04.5	Megalencephaly
Q04.6	Congenital cerebral cysts
Q04.8	Other specified congenital malformations of brain
Q04.9	Congenital malformation of brain, unspecified
Q05.5	Cervical spina bifida without hydrocephalus
Q05.6	Thoracic spina bifida without hydrocephalus
Q05.7	Lumbar spina bifida without hydrocephalus
Q05.8	Sacral spina bifida without hydrocephalus
Q05.9	Spina bifida, unspecified
Q06.1	Hypoplasia and dysplasia of spinal cord
Q06.3	Other congenital cauda equina malformations
Q06.8	Other specified congenital malformations of spinal cord
Q06.9	Congenital malformation of spinal cord, unspecified
Q28.0	Arteriovenous malformation of precerebral vessels

Q28.2	Arteriovenous malformation of cerebral vessels
Q28.3	Other malformations of cerebral vessels
Q67.5	Congenital deformity of spine
Q76.3	Congenital scoliosis due to congenital bony malformation
Q76.411	Congenital kyphosis, occipito-atlanto-axial region
Q76.412	Congenital kyphosis, cervical region
Q76.413	Congenital kyphosis, cervicothoracic region
Q76.49	Other congenital malformations of spine, not associated with scoliosis
S14.111A - S14.111S	Complete lesion at C1 level of cervical spinal cord
S14.121A - S14.129S	Central cord syndrome of cervical spinal cord
S14.131A - S14.139S	Anterior cord syndrome of cervical spinal cord
S14.141A - S14.149S	Brown-Sequard syndrome of cervical spinal cord
S14.151A - S14.159S	Other incomplete lesions of cervical spinal cord
S14.2xxA - S14.2xxS	Injury of nerve root of cervical spine
S24.111A - S24.119S	Complete lesion of thoracic spinal cord
S24.131A - S24.139S	Anterior cord syndrome of thoracic spinal cord
S24.141A - S24.149S	Brown-Sequard syndrome of thoracic spinal cord
S24.151A - S24.159S	Other incomplete lesions of thoracic spinal cord
S24.2xxA S24.2xxS	Injury of nerve root of thoracic spine
S34.111A - S34.119S	Complete lesion of lumbar spinal cord
S34.121A - S34.129S	Incomplete lesion of lumbar spinal cord



S34.21xA - S34.21xS	Injury of nerve root of lumbar spine
------------------------	--------------------------------------

CPT/HCPCS codes <i>not</i> considered medically necessary or considered experimental or investigational:	
<i>Code</i>	<i>Description</i>
51784	Electromyography studies (EMG) of anal or urethral sphincter, other than needle, any technique
51785	Needle electromyography studies (EMG) of anal or urethral sphincter, any technique
95860	Needle electromyography; 1 extremity with or without related paraspinal areas
95861	Needle measurement and recording of electrical activity of muscles of arms or legs
95863	Needle electromyography; 3 extremities with or without related paraspinal areas
95864	Needle electromyography; 4 extremities with or without related paraspinal areas
95869	Needle electromyography; thoracic paraspinal muscles (excluding T1 or T12)
95870	Needle electromyography; limited study of muscles in 1 extremity or non-limb (axial) muscles (unilateral or bilateral), other than thoracic paraspinal, cranial nerve supplied muscles, or sphincters
95928	Central motor evoked potential study (transcranial motor stimulation); upper limbs
95929	Central motor evoked potential study (transcranial motor stimulation); lower limbs
95930	Visual evoked potential (VEP) checkerboard or flash testing, central nervous system except glaucoma, with interpretation and report
95939	Central motor evoked potential study (transcranial motor stimulation); in upper and lower limbs
0333T	Visual evoked potential, screening of visual acuity, automated with report
S3900	Surface electromyography (EMG)

## References

1. Agarwal, N., Shabani, S., Huang, J., Ben-Natan, A. R., & Mummaneni, P. V. (2022). Intraoperative Monitoring for Spinal Surgery. *Neurologic clinics*, 40(2), 269–281. <https://doi.org/10.1016/j.ncl.2021.11.006>

2. Ajiboye RM, D'oro A, Ashana AO, et al. Routine Use of Intraoperative Neuromonitoring During ACDs for the Treatment of Spondylotic Myelopathy and Radiculopathy Is Questionable: A Review of 15,395 Cases. *Spine*. 2017;42(1):14-19.
3. Alzhrani F, Lenarz T, Teschner M. Facial palsy following cochlear implantation. *Eur Arch Otorhinolaryngol*. 2016;273(12):4199-4207.
4. American Academy of Audiology. Intraoperative neurophysiological monitoring. Scope of Practice. Reston, VA: American Academy of Audiology; updated January 2004.
5. American Academy of Otolaryngology - Head and Neck Surgery (AAO-HNS). Facial nerve monitoring. Position Statements. Alexandria, VA: AAO-HNS; submitted September 12, 1998. Available at: [http://www.entlink.net/practice/rules/facial\\_nerve\\_monitoring.cfm](http://www.entlink.net/practice/rules/facial_nerve_monitoring.cfm).
6. American Association of Neuromuscular & Electrodiagnostic Medicine. (n.d.). *Somatosensory Evoked Potentials: Clinical Uses*. Retrieved February, 17, 2021 from <https://www.aanem.org/Practice/Guidelines/Somatosensory-Evoked-Potentials-Clinical-Uses>
7. American Association of Neurological Surgeons/Congress of Neurological Surgeons (AANS/CNS) Joint Section on Disorders of the Spine and Peripheral Nerves (DSPN), Updated Position Statement: Intraoperative Electrophysiological Monitoring. January 2018. Available at: <https://spinesection.org/about/position-statements/interoperative-electrophysiological-monitoring/>.
8. American Association of Neurological Surgeons (AANS)/Congress of Neurological Surgeons (CNS). (2017). Guidelines for the Use of Electrophysiological Monitoring for Surgery of the Human Spinal Column and Spinal Cord. *Neurosurgery* 81(5):p 713-732, DOI: 10.1093/neuros/nyx466.
9. Bajwa ZH, Ho CC, Khan SA. Trigeminal neuralgia. UpToDate Inc., Waltham, MA. Last reviewed December 2017.
10. Barczyński M, Konturek A, Cichoń S. Randomized clinical trial of visualization versus neuromonitoring of recurrent laryngeal nerves during thyroidectomy. *Br J Surg*. 2009;96(3):240-246.
11. Basu, S., & Gohil, K. (2023). Intraoperative neuromonitoring in spine surgery: Does it reduce neural complications? Current evidence. *Indian Spine Journal*, 6(1), 15-26.
12. Beatty RM, McGuire P, Moroney JM, Holladay FP. Continuous intraoperative electromyographic recording during spinal surgery. *J Neurosurg*. 1995;82(3):401-405.
13. Bejjani GK, Nora PC, Vera PL, et al. The predictive value of intraoperative somatosensory evoked potential monitoring: Review of 244 procedures. *Neurosurgery*. 1998;43(3):491-500.
14. Beldi G, Kinsbergen T, Schlumpf R. Evaluation of intraoperative recurrent nerve monitoring in thyroid surgery. *World J Surg*. 2004; 28(6):589-591.
15. Bernhardt G, Awiszus F, Meister U, et al. The effect of intraoperative screw monitoring (root monitoring) with the INS-1 System (NUVASIVE) on the radiological outcome of dorsal instrumentation of the lumbar spine. *Z Orthop Unfall*. 2016;154(3):269-274.
16. Brauckhoff M, Gimm O, Thanh PN, et al. First experiences in intraoperative neurostimulation of the recurrent laryngeal nerve during thyroid surgery of children and adolescents. *J Pediatr Surg*. 2002;37(10):1414-1418.
17. Brennan J, Moore EJ, Shuler KJ. Prospective analysis of the efficacy of continuous intraoperative nerve monitoring during thyroidectomy, parathyroidectomy, and parotidectomy. *Otolaryngol Head Neck Surg*. 2001;124(5):537-543.
18. Brock S, Scaioli V, Ferroli P, Broggi G. Neurovascular decompression in trigeminal neuralgia: Role of intraoperative neurophysiological monitoring in the learning period. *Stereotact Funct Neurosurg*. 2004;82(5-6):199-206.

19. Brown MJ. Anesthesia for elective spine surgery in adults. UpToDate.com. Last Update: Jan 12, 2022. Retrieved from:  
[https://www.uptodate.com/contents/anesthesia-for-elective-spine-surgery-in-adults?search=anesthesia%20for%20elective%20spine%20surgery%20in%20adults&source=search\\_result&selectedTitle=1~150&usage\\_type=default&display\\_rank=1](https://www.uptodate.com/contents/anesthesia-for-elective-spine-surgery-in-adults?search=anesthesia%20for%20elective%20spine%20surgery%20in%20adults&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1)
20. Buhl, L. K., Bastos, A. B., Pollard, R. J., Arle, J. E., Thomas, G. P., Song, Y., & Boone, M. D. (2021). Neurophysiologic Intraoperative Monitoring for Spine Surgery: A Practical Guide From Past to Present. *Journal of intensive care medicine*, 36(11), 1237–1249.  
<https://doi.org/10.1177/0885066620962453>
21. Cavicchi O, Caliceti U, Fernandez IJ, et al. The value of neurostimulation and intraoperative nerve monitoring of inferior laryngeal nerve in thyroid surgery. *Otolaryngol Head Neck Surg*. 2009;140(6):866-870.
22. Charalampidis, A., Jiang, F., Wilson, J. R., Badhiwala, J. H., Brodke, D. S., & Fehlings, M. G. (2020). The use of intraoperative neurophysiological monitoring in spine surgery. *Global spine journal*, 10(1\_suppl), 104S-114S.
23. Chiang FY, Lu IC, Kuo WR, et al. The mechanism of recurrent laryngeal nerve injury during thyroid surgery--the application of intraoperative neuromonitoring. *Surgery*. 2008;143(6):743-749.
24. Cole T, Veeravagu A, Zhang M, Li A, Ratliff JK. Intraoperative neuromonitoring in single-level spinal procedures: a retrospective propensity score-matched analysis in a national longitudinal database. *Spine*. 2014;39(23):1950-9.
25. Dackiw AP, Rotstein LE, Clark OH. Computer-assisted evoked electromyography with stimulating surgical instruments for recurrent/external laryngeal nerve identification and preservation in thyroid and parathyroid operation. *Surgery*. 2002;132(6):1100-1106; discussion 1107-1108.
26. de Blas G, Barrios C, Regidor I, et al. Safe pedicle screw placement in thoracic scoliotic curves using t-EMG: Stimulation threshold variability at concavity and convexity in apex segments. *Spine (Phila Pa 1976)*. 2012;37(6):E387-E395.
27. Di Martino, A., Papalia, R., Caldaria, A., Torre, G., Denaro, L., & Denaro, V. (2019). Should evoked potential monitoring be used in degenerative cervical spine surgery? A systematic review. *Journal of Orthopaedics and Traumatology*, 20, 1-8.
28. Di Martino E, Sellhaus B, Haensel J, et al. Fallopian canal dehiscences: A survey of clinical and anatomical findings. *Eur Arch Otorhinolaryngol*. 2005;262(2):120-126.
29. Dimov RS, Doikov IJ, Mitov FS, et al. Intraoperative identification of recurrent laryngeal nerves in thyroid surgery by electrical stimulation. *Folia Med (Plovdiv)*. 2001;43(4):10-13.
30. Dionigi G, Donatini G, Boni L, et al. Continuous monitoring of the recurrent laryngeal nerve in thyroid surgery: A critical appraisal. *Int J Surg*. 2013;11 Suppl 1:S44-S46.
31. Djohan RS, Rodriguez HE, Connolly MM, et al. Intraoperative monitoring of recurrent laryngeal nerve function. *Am Surg*. 2000;66(6):595-597.
32. Dodo, Y., Okano, I., Zelenty, W. D., Paek, S., Sarin, M., Haffer, H., ... & Hughes, A. P. (2023). The Utilization of Intraoperative Neurophysiological Monitoring for Lumbar Decompression and Fusion Surgery in New York State. *Spine*, 10-1097.
33. Dulguerov P, Marchal F, Lehmann W. Postparotidectomy facial nerve paralysis: Possible etiologic factors and results with routine facial nerve monitoring. *Laryngoscope*. 1999;109(5):754-762.
34. Erickson L, Costa V, and McGregor M. Use of intraoperative neurophysiological monitoring during spinal surgery. McGill University Health Center. Report Number 20, July 7, 2005. Available at: [http://upload.mcgill.ca/tau/SPINAL\\_MONITORING\\_Final.pdf](http://upload.mcgill.ca/tau/SPINAL_MONITORING_Final.pdf)

35. Erickson L, Costa V, McGregor M. Intraoperative neurophysiological monitoring during spinal surgery. Montreal, QC: Technology Assessment Unit of the McGill University Health Centre (MUHC); 2005:1-39.
36. Fabregas N, Gomar C. Monitoring in neuroanaesthesia: Update of clinical usefulness. *Eur J Anaesthesiol.* 2001;18(7):423-439.
37. Fayad JN, Wanna GB, Micheletto JN, Parisier SC. Facial nerve paralysis following cochlear implant surgery. *Laryngoscope.* 2003;113(8):1344-6.
38. Fehlings MG, Brodke DS, Norvell DC, Dettori JR. The evidence for intraoperative neurophysiological monitoring in spine surgery: Does it make a difference? *Spine.* 2010;35(9 Suppl):S37-S46.
39. Finkelstein JA. Can triggered electromyograph thresholds predict safe thoracic pedicle screw placement. *Spine.* 2003;28(9):960.
40. Glassman SD, Dimar JR, Puno RM, et al. A prospective analysis of intraoperative electromyographic monitoring of pedicle screw placement with computed tomographic scan confirmation. *Spine (Phila Pa 1976).* 1995;20:1375-1379.
41. Grabb PA, Albright AL, Scwabassi RJ, Pollack IF. Continuous intraoperative electromyographic monitoring of cranial nerves during resection of fourth ventricular tumors in children. *Neurosurg.* 1997;86(1):1-4.
42. Grade M, Beham AW, Schüler P, Kneist W, Ghadimi BM. Pelvic intraoperative neuromonitoring during robotic-assisted low anterior resection for rectal cancer. *J Robot Surg.* 2016;10(2):157-60.
43. Grosheva M, Klussmann JP, Grimminger C, et al. Electromyographic facial nerve monitoring during parotidectomy for benign lesions does not improve the outcome of postoperative facial nerve function: A prospective two-center trial. *Laryngoscope.* 2009;119(12):2299-2305.
44. Grover HJ, Thornton R, Lutchman LN, Blake JC. Using Transcranial Magnetic Stimulation to Evaluate the Motor Pathways After an Intraoperative Spinal Cord Injury and to Predict the Recovery of Intraoperative Transcranial Electrical Motor Evoked Potentials: A Case Report. *J Clin Neurophysiol.* 2016;33(3):e8-e11.
45. Gunnarsson T, Krassioukov AV, Sarjeant R, Fehlings MG. Real-time continuous intraoperative electromyographic and somatosensory evoked potential recordings in spinal surgery: Correlation of clinical and electrophysiologic findings in a prospective, consecutive series of 213 cases. *Spine.* 2004;29(6):677-684.
46. Harner SG, Daube JR, Ebersold MJ, Beatty CW. Improved preservation of facial nerve function with use of electrical monitoring during removal of acoustic neuromas. *Mayo Clin Proc.* 1987;62(2):92-102.
47. Harper C. Intraoperative cranial nerve monitoring. *Muscle Nerve.* 2004;29(3):339-351.
48. Harper CM, Daube JR. Facial nerve electromyography and other cranial nerve monitoring. *J Clin Neurophysiol.* 1998;15(3):206-216.
49. Harrison BJ, Triponez F. Intraoperative adjuncts in surgery for primary hyperparathyroidism. *Langenbecks Arch Surg.* 2009;394(5):799-809.
50. Hausmann ON, Boni T, Pfirrmann CW, et al. Preoperative radiological and electrophysiological evaluation in 100 adolescent idiopathic scoliosis patients. *Eur Spine J.* 2003;12(5):501-506.
51. Hayashi H, Kawaguchi M. Intraoperative monitoring of flash visual evoked potential under general anesthesia. *Korean J Anesthesiol.* 2017;70(2):127-135.
52. Hayes, Inc. Health Technology Assessment. *Intraoperative Neuromonitoring for Lumbar Spinal Fusion and Discectomy.* Lansdale, PA: Hayes, Inc. Annual Review: Dec 28, 2023.
53. Hemmerling TM, Schmidt J, Bosert C, et al. Intraoperative monitoring of the recurrent laryngeal nerve in 151 consecutive patients undergoing thyroid surgery. *Anesth Analg.* 2001;93(2):396-399.

54. Hermann M, Hellebart C, Freissmuth M. Neuromonitoring in thyroid surgery: Prospective evaluation of intraoperative electrophysiological responses for the prediction of recurrent laryngeal nerve injury. *Ann Surg.* 2004;240(1):9-17.
55. Hillermann CL, Tarpey J, Phillips DE. Laryngeal nerve identification during thyroid surgery -- feasibility of a novel approach. *Can J Anaesth.* 2003;50(2):189-192.
56. Holland NR. Intraoperative electromyography during thoracolumbar spinal surgery. *Spine.* 1998;23(17):1915-1922.
57. Holland NR. Intraoperative electromyography. *J Clin Neurophysiol.* 2002;19(5):444-453.
58. Horn D, Rotzsch VM. Intraoperative electromyogram monitoring of the recurrent laryngeal nerve: Experience with an intralaryngeal surface electrode. A method to reduce the risk of recurrent laryngeal nerve injury during thyroid surgery. *Langenbecks Arch Surg.* 1999;384(4):392-395.
59. Houlden, D. A., Turgeon, C. A., Amyot, N. S., Edem, I., Sinclair, J., Agbi, C., ... & Alkherayf, F. (2019). Intraoperative flash visual evoked potential recording and relationship to visual outcome. *Canadian Journal of Neurological Sciences*, 46(3), 295-302.
60. Hsieh HS, Wu CM, Zhuo MY, Yang CH, Hwang CF. Intraoperative facial nerve monitoring during cochlear implant surgery: an observational study. *Medicine (Baltimore).* 2015;94(4):e456.
61. Hsu GS. Improving hearing in stapedectomy with intraoperative auditory brainstem response. *Oto Head Neck Surg.* 2011 Jan;(144)1):60-3
62. Sobottka SB, Schackert G, May SA, Wiegleb M, Reiss G. Intraoperative facial nerve monitoring (IFNM) predicts facial nerve outcome after resection of vestibular schwannoma. *Acta Neurochir (Wien).* 1998;140(3):235-42.
63. Hu J, Fleck TR, Xu J, et al. Contemporary changes with the use of facial nerve monitoring in chronic ear surgery. *Otolaryngol Head Neck Surg.* 2014;151(3):473-477.
64. Huang BR, Chang CN, Hsu JC. Intraoperative electrophysiological monitoring in microvascular decompression for hemifacial spasm. *J Clin Neurosci.* 2009;16(2):209-213.
65. Hussain NS. Analysis of 1014 consecutive operative cases to determine the utility of intraoperative neurophysiological data. *Asian J Neurosurg.* 2015;10(3):166-172.
66. Ibrahim T, Mrowczynski O, Zalatimo O, et al. The Impact of Neurophysiological Intraoperative Monitoring during Spinal Cord and Spine Surgery: A Critical Analysis of 121 Cases. *Cureus.* 2017;9(11):e1861. Published 2017 Nov 19. doi:10.7759/cureus.1861
67. Ingelmo I, Trapero JG, Puig A, et al. Intraoperative monitoring of the facial nerve: Anesthesia and neurophysiology considerations. *Rev Esp Anesthesiol Reanim.* 2003;50(9):460-471.
68. NCT02446808: Intraoperative Nerve Monitoring During Robot-assisted Laparoscopic Prostatectomy. Accessed December 9, 2017. <https://clinicaltrials.gov/ct2/show/NCT02446808>
69. Jiang L, Ma Y, Jiao Y. Decompression and monitor of facial nerve during cholesteatoma surgery in petrous part of temporal bone. *Lin Chuang Er Bi Yan Hou Ke Za Zhi.* 2006;20(16):741-743.
70. Jimenez JC, Sani S, Braverman B, et al. Palsies of the fifth cervical nerve root after cervical decompression: Prevention using continuous intraoperative electromyography monitoring. *J Neurosurg Spine.* 2005;3(2):92-97.
71. Jonas J, Bahr R. Neuromonitoring of the external branch of the superior laryngeal nerve during thyroid surgery. *Am J Surg.* 2000;179(3):234-236.
72. Kaliya-Perumal AK, Charng JR, Niu CC, et al. Intraoperative electromyographic monitoring to optimize safe lumbar pedicle screw placement - a retrospective analysis. *BMC Musculoskelet Disord.* 2017;18(1):229.
73. Kauff DW, Kronfeld K, Gorbulev S, Wachtlin D, Lang H, Kneist W. Continuous intraoperative monitoring of pelvic autonomic nerves during TME to prevent urogenital and anorectal

- dysfunction in rectal cancer patients (NEUROS): a randomized controlled trial. *BMC Cancer*. 2016;16:323.
74. Kiviniemi H, Vornanen T, Mäkelä J. Prevention of complications of thyroid and parathyroid surgery. *Duodecim*. 2010;126(3):269-275.
  75. Koht A, Sloan TB, Hemmer LB. Neuromonitoring in surgery and anesthesia. UpToDate.com. Last Updated Oct 21, 2022. Retrieved from: [https://www.uptodate.com/contents/neuromonitoring-in-surgery-and-anesthesia?search=Neuro monitoring%20in%20surgery%20and%20anesthesia&source=search\\_result&selectedTitle=1~150 &usage\\_type=default&display\\_rank=1](https://www.uptodate.com/contents/neuromonitoring-in-surgery-and-anesthesia?search=Neuro%20monitoring%20in%20surgery%20and%20anesthesia&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1)
  76. Kombos T, Suess O, Kern BC, et al. Can continuous intraoperative facial electromyography predict facial nerve function following cerebellopontine angle surgery? *Neurol Med Chir (Tokyo)*. 2000;40(10):501-505; discussion 506-507.
  77. Krassioukov AV, Sarjeant R, Arkia H, Fehlings MG. Multimodality intraoperative monitoring during complex lumbosacral procedures: Indications, techniques, and long-term follow-up review of 61 consecutive cases. *J Neurosurg Spine*. 2004;1(3):243-253.
  78. Krause, K. L., Cheaney II, B., Obayashi, J. T., Kawamoto, A., & Than, K. D. (2020). Intraoperative neuromonitoring for one-level lumbar discectomies is low yield and cost-ineffective. *Journal of Clinical Neuroscience*, 71, 97-100.
  79. Kundnani VK, Zhu L, Tak H, Wong H. Multimodal intraoperative neuromonitoring in corrective surgery for adolescent idiopathic scoliosis: Evaluation of 354 consecutive cases. *Indian J Orthop*. 2010;44(1):64-72.
  80. Kwartler JA, Luxford WM, Atkins J, Shelton C. Facial nerve monitoring in acoustic tumor surgery. *Otolaryngol Head Neck Surg*. 1991;104(6):814-817.
  81. Lee CH, Kim HW, Kim HR, et al. Can triggered electromyography thresholds assure accurate pedicle screw placements? A systematic review and meta-analysis of diagnostic test accuracy. *Clin Neurophysiol*. 2015;126(10):2019-2025.
  82. Lefaucheur JP, Neves DO, Vial C. Electrophysiological monitoring of cranial motor nerves (V, VII, IX, X, XI, XII). *Neurochirurgie*. 2009;55(2):136-141.
  83. Legatt AD, Schroeder CE, Gill B, Goodrich JT. Electrical stimulation and multichannel EMG recording for identification of functional neural tissue during cauda equina surgery. *Childs Nerv Syst*. 1992;8(4):185-189.
  84. Legatt AD, Emerson RG, Epstein CM. ACNS Guideline: Transcranial Electrical Stimulation Motor Evoked Potential Monitoring. *Journal of Clinical Neurophysiology*. 2016; 33(1): 42–50. [https://www.acns.org/UserFiles/file/ACNS\\_Guideline\\_\\_\\_Transcranial\\_Electrical.9\\_v1.pdf](https://www.acns.org/UserFiles/file/ACNS_Guideline___Transcranial_Electrical.9_v1.pdf)
  85. Leonetti JP, Matz GJ, Smith PG, Beck DL. Facial nerve monitoring in otologic surgery: Clinical indications and intraoperative technique. *Ann Otol Rhinol Laryngol*. 1990;99(11):911-918.
  86. Limbrick DD Jr, Wright NM. Verification of nerve root decompression during minimally-invasive lumbar microdiscectomy: A practical application of surgeon-driven evoked EMG. *Minim Invasive Neurosurg*. 2005;48(5):273-277.
  87. López JR. Neurophysiologic intraoperative monitoring of the oculomotor, trochlear, and abducens nerves. *J Clin Neurophysiol*. 2011;28(6):543-50.
  88. Lopez JR. The use of evoked potentials in intraoperative neurophysiologic monitoring. *Phys Med Rehabil Clin N Am*. 2004;15(1):63-84.
  89. Luo Y, Regli L, Bozinov O, Sarnthein J. Clinical utility and limitations of intraoperative monitoring of visual evoked potentials. *PLoS ONE*. 2015;10(3):e0120525.
  90. Maguire J, Wallace S, Madiga R, et al. Evaluation of intrapedicular screw position using intraoperative evoked electromyography. *Spine*. 1995;20(9):1068-1074.

91. Mammis A, Mogilner AY. The use of intraoperative electrophysiology for the placement of spinal cord stimulator paddle leads under general anesthesia. *Neurosurgery*. 2012;70(2 Suppl Operative):230-236.
92. Mao, G., Elhamdani, S., Gigliotti, M. J., Mace, Z., Sciabassi, R., Oh, M., & Whiting, D. (2021). Neurologic Complications in Monitored versus Unmonitored Image-Guidance Assisted Posterior Lumbar Instrumentation. *World neurosurgery*, 152, e155-e160.
93. Marcus B, Edwards B, Yoo S, et al. Recurrent laryngeal nerve monitoring in thyroid and parathyroid surgery: The University of Michigan experience. *Laryngoscope*. 2003;113(2):356-361.
94. Marusch F, Hussock J, Haring G, et al. Influence of muscle relaxation on neuromonitoring of the recurrent laryngeal nerve during thyroid surgery. *Br J Anaesth*. 2005;94(5):596-600.
95. Metwali H, Kniese K, Fahlbusch R. Intraoperative monitoring of the integrity of the anterior visual pathways: a methodological review and meta-analysis. *World Neurosurg*. 2017;
96. Mikula AL, Williams SK, Anderson PA. The use of intraoperative triggered electromyography to detect misplaced pedicle screws: A systematic review and meta-analysis. *J Neurosurg Spine*. 2016;24(4):624-638.
97. Minahan RE, Mandir AS. Neurophysiologic intraoperative monitoring of trigeminal and facial nerves. *J Clin Neurophysiol*. 2011;28(6):551-565.
98. National Institute for Health and Clinical Excellence (NICE). Intraoperative nerve monitoring during thyroid surgery. *Interventional Procedure Guidance 255*. London, UK: NICE; March 2008. Available at: <http://www.nice.org.uk/nicemedia/pdf/IPG255Guidance.pdf>.
99. Neff BA, Ting J, Dickinson SL, Welling DB. Facial nerve monitoring parameters as a predictor of postoperative facial nerve outcomes after vestibular schwannoma resection. *Otol Neurotol*. 2005;26(4):728-732.
100. Nuwer, M. R., Husain, A. M., & Soto, F. (2022). Overview of intraoperative neuromonitoring. *Handbook of clinical neurology*, 186, 3–9. <https://doi.org/10.1016/B978-0-12-819826-1.00011-9>
101. Assessment: Intraoperative neurophysiology. Report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology*. 1990;40(11):1644-1646.
102. Intraoperative monitoring of the facial nerve - evoked potentials. *Tech Brief*. Chicago, IL: American Medical Association; March 1994.
103. Norcross-Nechay K, Mathew T, Simmons JW, et al. Intraoperative somatosensory evoked potential findings in acute and chronic spinal canal compromise. *Spine*. 1999;24(10):1029-1033.
104. Nuwer MR, Emerson RG, Galloway G, et al. Evidence-based guideline update: Intraoperative spinal monitoring with somatosensory and transcranial electrical motor evoked potentials: Report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology and the American Clinical Neurophysiology Society. *Neurology* 2012;78;585-589.
105. Olds MJ, Rowan PT, Isaacson JE, Silverstein H. Facial nerve monitoring among graduates of the Ear Research Foundation. *Am J Otol*. 1997;18(4):507-511.
106. Ota T, Kawai K, Kamada K, Kin T, Saito N. Intraoperative monitoring of cortically recorded visual response for posterior visual pathway. *J Neurosurg*. 2010;112(2):285-94.
107. Otto RA, Cochran CS. Sensitivity and specificity of intraoperative recurrent laryngeal nerve stimulation in predicting postoperative nerve paralysis. *Ann Otol Rhinol Laryngol*. 2002;111(11):1005-1007.
108. Ovadia D, Korn A, Fishkin M, et al. The contribution of an electronic conductivity device to the safety of pedicle screw insertion in scoliosis surgery. *Spine (Phila Pa 1976)*. 2011;36:E1314-E1321.
109. Owen JH, Kostuik JP, Gornet M, et al. The use of mechanically elicited electromyograms to protect nerve roots during surgery for spinal degeneration. *Spine*. 1994;19(15):1704-1710.

110. Paradiso G, Lee GY, Sarjeant R, et al. Multimodality intraoperative neurophysiologic monitoring findings during surgery for adult tethered cord syndrome: Analysis of a series of 44 patients with long-term follow-up. *Spine*. 2006;31(18):2095-2102.
111. Prell J, Strauss C, Plontke SK, Rampf S. Intraoperative monitoring of the facial nerve : Vestibular schwannoma surgery. *HNO*. 2017;65(5):404-412.
112. Raynor BL, Lenke LG, Kim Y, et al. Can triggered electromyograph thresholds predict safe thoracic pedicle screw placement? *Spine*. 2002;27(18):2030-2035.
113. Raynor BL, Padberg AM, Lenke LG, et al. Failure of intraoperative monitoring to detect postoperative neurologic deficits: A 25-year experience in 12,375 spinal surgeries. *Spine (Phila Pa 1976)*. 2016;41(17):1387-1393.
114. Reidy DP, Houlden D, Nolan PC, et al. Evaluation of electromyographic monitoring during insertion of thoracic pedicle screws. *J Bone Joint Surg Br*. 2001;83(7):1009-1014.
115. Resnick D, Choudhri T, Dailey A et al. Guidelines for the performance of fusion procedures for degenerative disease of the lumbar spine. Part 15: Electrophysiological monitoring and lumbar fusion. *J Neurosurg Spine*. 2005;2(6):725-732.
116. Robertson ML, Steward DL, Gluckman JL, and Welge J. Continuous laryngeal nerve integrity monitoring during thyroidectomy: Does it reduce risk or injury? *Otolaryngol Head Neck Surg*. 2004;131(5):596-600.
117. Roland PS, Meyerhoff WL. Intraoperative electrophysiological monitoring of the facial nerve: Is it standard of practice? *Am J Otolaryngol*. 1994;15(4):267-270.
118. Roland PS. Monitors, facial nerve. *eMedicine Otolaryngology*. Omaha, NE: eMedicine.com; updated May 6, 2003. Available at: <http://www.emedicine.com/ent/topic447.htm>.
119. Romaniszyn M, Walega P, Nowakowski M, Nowak W. Can surface electromyography improve surgery planning? Electromyographic assessment and intraoperative verification of the nerve bundle entry point location of the gracilis muscle. *J Electromyogr Kinesiol*. 2016;28:1-6.
120. Sasaki T. Intraoperative monitoring of visual evoked potential: introduction of a clinically useful method. *J Neurosurg*. 2010 Feb;112(2):273-84. doi:10.3171/2008.9.JNS08451.
121. Scherl SA. Treatment and prognosis of adolescent idiopathic scoliosis. Last reviewed December 2017. UpToDate Inc. Waltham, MA.
122. Schlake HP, Goldbrunner R, Siebert M, Behr R, Roosen K. Intra-Operative electromyographic monitoring of extra-ocular motor nerves (Nn. III, VI) in skull base surgery. *Acta Neurochir (Wien)*. 2001;143(3):251-61.
123. Schlake HP, Goldbrunner RH, Milewski C, et al. Intra-operative electromyographic monitoring of the lower cranial motor nerves (LCN IX-XII) in skull base surgery. *Clin Neurol Neurosurg*. 2001;103(2):72-82.
124. Sekiya T, Hatayama T, Iwabuchi T, Maeda S. Intraoperative recordings of evoked extraocular muscle activities to monitor ocular motor nerve function. *Neurosurgery*. 1993;32(2):227-35.
125. Selesnick SH, Lynn-Macrae AG. The incidence of facial nerve dehiscence at surgery for cholesteatoma. *Otol Neurotol*. 2001;22(2):129-132.
126. Selesnick SH, Victor JD, Tikoo RK, Eisenman DJ. Predictive value of intraoperative brainstem auditory evoked responses in surgery for conductive hearing losses. *Am J Otol*. 1997;18(1):2-9.
127. Shan XF, Lin B, Lu XG, et al. Electromyographic monitoring of facial nerve during parotid surgery. *Beijing Da Xue Xue Bao*. 2014;46(1):48-52.
128. Sheshadri V, Bharadwaj S, Chandramouli BA. Intra-operative electrooculographic monitoring to prevent post-operative extraocular motor nerve dysfunction during skull base surgeries. *Indian J Anaesth*. 2016;60(8):560-5.
129. Shils JL, Arle JE. Intraoperative neurophysiologic methods for spinal cord stimulator placement under general anesthesia. *Neuromodulation*. 2012;15(6):560-571; discussion 571-572.



130. Shkarubo AN, Chernov IV, Ogurtsova AA, et al. Neurophysiological Identification of Cranial Nerves During Endoscopic Endonasal Surgery of Skull Base Tumors: Pilot Study Technical Report. *World Neurosurg.* 2017;98:230-238.
131. Siebenrock KA, Rösler KM, Gonzalez E, Ganz R. Intraoperative electromyography of the superior gluteal nerve during lateral approach to the hip for arthroplasty: a prospective study of 12 patients. *J Arthroplasty.* 2000;15(7):867-70.
132. Sitthinamsuwan B, Chanvanitkulchai K, Phonwijit L, et al. Utilization of intraoperative electromyography for selecting targeted fascicles and determining the degree of fascicular resection in selective tibial neurotomy for ankle spasticity. *Acta Neurochir (Wien).* 2013;155(6):1143-1149.
133. Skaggs DL, Choi PD, Rice C, et al. Efficacy of intraoperative neurologic monitoring in surgery involving a vertical expandable prosthetic titanium rib for early-onset spinal deformity. *J Bone Joint Surg Am.* 2009;91(7):1657-1663.
134. Slimp JC. Electrophysiologic intraoperative monitoring for spine procedures. *Phys Med Rehabil Clin N Am.* 2004;15(1):85-105.
135. Smith PN, Balzer JR, Khan MH, et al. Intraoperative somatosensory evoked potential monitoring during anterior cervical discectomy and fusion in nonmyelopathic patients--a review of 1,039 cases. *Spine J.* 2007;7(1):83-7.
136. Spitz S, Felbaum D, Aghdam N, Sandhu F. Delayed postoperative C5 root palsy and the use of neurophysiologic monitoring. *Eur Spine J.* 2015;24(12):2866-2871.
137. Swanson BT, Holst B, Infante J, Poenitzsch J, Ortiz A. EMG activity of selected rotator cuff musculature during grade III distraction and posterior glide glenohumeral mobilization: results of a pilot trial comparing painful and non-painful shoulders. *J Man Manip Ther.* 2016;24(1):7-13.
138. Terrell JE, Kileny PR, Yian C, et al. Clinical outcome of continuous facial nerve monitoring during primary parotidectomy. *Arch Otolaryngol Head Neck Surg.* 1997;123(10):1081-1087.
139. Thirumala PD, Mohanraj SK, Habeych M, et al. Value of free-run electromyographic monitoring of lower cranial nerves in endoscopic endonasal approach to skull base surgeries. *J Neurol Surg B Skull Base.* 2012;73(4):236-44.
140. Thirumala PD, Muralidharan A, Loke YK, et al. Value of intraoperative neurophysiological monitoring to reduce neurological complications in patients undergoing anterior cervical spine procedures for cervical spondylotic myelopathy. *J Clin Neurosci.* 2016;25:27-35.
141. Thom JJ, Carlson ML, Olson MD, et al. The prevalence and clinical course of facial nerve paresis following cochlear implant surgery. *Laryngoscope.* 2013;123(4):1000-4.
142. Thomusch O, Sekulla C, Walls G, et al. Intra-operative neuromonitoring of surgery for benign goiter. *Am J Surg.* 2002;183(6):673-678.
143. Timon CI, Rafferty M. Nerve monitoring in thyroid surgery: Is it worthwhile? *Clin Otolaryngol.* 1999;24(6):487-490.
144. Tomoda C, Hirokawa Y, Uruno T, et al. Sensitivity and specificity of intraoperative recurrent laryngeal nerve stimulation test for predicting vocal cord palsy after thyroid surgery. *World J Surg.* 2006;30(7):1230-1233.
145. Tschopp KP, Gottardo C. Comparison of various methods of electromyographic monitoring of the recurrent laryngeal nerve in thyroid surgery. *Ann Otol Rhinol Laryngol.* 2002;111(9):811-816.
146. Uribe, J. S., Isaacs, R. E., Youssef, J. A., Khajavi, K., Balzer, J. R., Kanter, A. S., ... & SOLAS Degenerative Study Group. (2015). Can triggered electromyography monitoring throughout retraction predict postoperative symptomatic neuropraxia after XLIF? Results from a prospective multicenter trial. *European Spine Journal*, 24, 378-385.

147. Vial Ch, Bouhour F. Electrophysiological examination of the cranial nerves: Technical aspects and practical applications. *Rev Med Liege*. 2004;59 Suppl 1:82-90.
148. Weiss DS. Spinal cord and nerve root monitoring during surgical treatment of lumbar stenosis. *Clin Orthop Relat Res*. 2001;(384):82-100.
149. Wi, S. M., Lee, H. J., Kang, T., Chang, S. Y., Kim, S. M., Chang, B. S., ... & Kim, H. (2020). Clinical significance of improved intraoperative neurophysiological monitoring signal during spine surgery: a retrospective study of a single-institution prospective cohort. *Asian spine journal*, 14(1), 79.
150. Witt RL. Facial nerve monitoring in parotid surgery: The standard of care? *Otolaryngol Head Neck Surg*. 1998;119(5):468-470.
151. Silverstein J, Mermelstein L, DeWal H, Basra S. Saphenous nerve somatosensory evoked potentials: A novel technique to monitor the femoral nerve during transpsoas lumbar lateral interbody fusion. *Spine (Phila Pa 1976)*. 2014;39(15):1254-1260.
152. Kneist, W. et al. Is intraoperative neuromonitoring associated with better functional outcome in patients undergoing open TME? *European Journal of Surgical Oncology*, Volume 39, Issue 9, 994 - 999.
153. Jellish WS, Jensen RL, Anderson DE, Shea JF. Intraoperative electromyographic assessment of recurrent laryngeal nerve stress and pharyngeal injury during anterior cervical spine surgery with Caspar instrumentation. *J Neurosurg*. 1999;91(2 Suppl):170-4.
154. Chen CC, Huang YC, Lee ST, Chen JF, Wu CT, Tu PH. Long-term result of vocal cord paralysis after anterior cervical disectomy. *Eur Spine J*. 2014;23(3):622-6.
155. Dimopoulos VG, Chung I, Lee GP, et al. Quantitative estimation of the recurrent laryngeal nerve irritation by employing spontaneous intraoperative electromyographic monitoring during anterior cervical disectomy and fusion. *J Spinal Disord Tech*. 2009;22(1):1-7.
156. Macdonald DB, Skinner S, Shils J, Yingling C; American Society of Neurophysiological Monitoring. Intraoperative motor evoked potential monitoring - a position statement by the American Society of Neurophysiological Monitoring. *Clin Neurophysiol*. 2013;124:2291-2316.
157. Cofano F, Zenga F, Mammi M, Altieri R, Marengo N, Ajello M, Pacca P, Melcarne A, Junemann C, Ducati A, Garbossa D. Intraoperative neurophysiological monitoring during spinal surgery: technical review in open and minimally invasive approaches. *Neurosurg Rev*. 2018 Jan 8. doi: 10.1007/s10143-017-0939-4. [Epub ahead of print]
158. Daniel JW, Botelho RV, Milano JB, Dantas FR, Onishi FJ, Neto ER, Bertolini EF, Borgheresi MAD, Joaquim AF. Intraoperative Neurophysiological Monitoring in Spine Surgery: A Systematic Review and Meta-Analysis. *Spine (Phila Pa 1976)*. 2018;43:1154-1160.
159. Pastorelli F, Di Silvestre M, Plasmati R, Michelucci R, Greggi T, Morigi A, Bacchin MR, Bonarelli S, Cioni A, Vommaro F, Fini N, Lolli F, Parisini P. The prevention of neural complications in the surgical treatment of scoliosis: the role of the neurophysiological intraoperative monitoring. *Eur Spine J*. 2011;20 Suppl 1:S105-114.
160. Heman-Ackah, SE, Gupta S, Lalwani AK. Is facial nerve integrity monitoring of value in chronic ear surgery? *The Laryngoscope*. 2012:2-3. Doi:10.1002/lary.23363
161. Hu, J, Fleck TR, Xu J, et al. Contemporary changes with the use of facial nerve monitoring in chronic ear surgery. *Otolaryngology--Head and Neck Surgery*. 2014;151(3): 473-477. Doi:10.1177/0194599814537223
162. American Academy of Otolaryngology - Head and Neck Surgery (AAO-HNS). Position Statement: Intraoperative Nerve Monitoring in Otologic Surgery. Alexandria, VA: AAO-HNS; adopted March 12, 2017. Available at: <https://www.entnet.org/intraoperative-nerve-monitoring>. Accessed March 5, 2018.

163. Wilson L, Lin E, Lalwani A. Cost-effectiveness of intraoperative facial nerve monitoring in middle ear or mastoid surgery. *The Laryngoscope*. 2003;113(10):1736-1745.
164. Guideline 11 C: RECOMMENDED STANDARDS FOR INTRAOPERATIVE MONITORING OF AUDITORY EVOKED POTENTIALS. Acns.org.  
<https://www.acns.org/pdf/guidelines/Guideline-11C.pdf> Published Oct, 2009.
165. Guideline 11A: RECOMMENDED STANDARDS FOR NEUROPHYSIOLOGIC INTRAOPERATIVE MONITORING – PRINCIPLES. Acns.org.  
<https://www.acns.org/pdf/guidelines/Guideline-11A.pdf> Published Oct, 2009.
166. Sinha SR, Sullivan LR, Sabauet D, et al. American Clinical Neurophysiology Society Guideline 1: Minimum technical requirements for performing clinical electroencephalography. *The Neurodiagnostic Journal*. 2016; 56(4): 235-244.

#### Clinical Guideline Revision / History Information

Original Date: 4/13/2018

Reviewed/Revised: 4/15/2019, 7/21/2020, 04/21/2021, 12/01/2021, 04/25/2022, 05/02/2023,  
04/16/2024