

Outpatient Physical Therapy (PT) and Occupational Therapy (OT)

Disclaimer

Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.

Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.

Summary

The Plan members may require physical therapy (PT) when it is prescribed by a qualified health professional to significantly restore or improve functioning, relieve disease symptoms, and prevent disability in individuals with acute and/or chronic disease. A PT treatment plan of care may consist of heat and cold therapy, electric stimulation, manual therapy, a variety of exercise regimens, functional training for ambulatory activities, and the development of a home exercise program. PT may only be performed by a qualified, licensed physical therapist or by a physical therapist assistant (PTA) under the supervision of a qualified, licensed physical therapist.

The Plan members may require occupational therapy (OT) when it is prescribed by a qualified health professional to significantly regain skills of daily living that have been lost or impaired through disease or injury. OT treatments are used for both rehabilitation and habilitation, and are designed with purposeful activities. OT may only be performed by a qualified, licensed occupational therapist, or by an occupational therapist assistant (OTA) under the supervision of a qualified, licensed occupational therapist.

PT and OT are often coordinated by a multidisciplinary team of licensed therapists, nurses, and prescribing clinicians. PT or OT is usually one of numerous components that are part of a multidisciplinary treatment plan of care following injury or the diagnosis of a chronic disease.

For home physical therapy, please refer to the Plan Clinical Guideline: Home Care - Physical Therapy (PT) and Occupational Therapy (OT) (CG021).

Definitions

“Physical Therapy (PT)” refers to supervised therapeutic procedures performed by licensed healthcare professionals which are intended to relieve disease symptoms, prevent disability and restore clinical function. PT is often one of many components in a multidisciplinary treatment plan following injury or in chronic disease. Physical therapy may include, but is not limited to:

- Ambulation and mobility training
- Gait and balance training
- Strength training
- Joint mobilization
- Neuromuscular reeducation
- Therapeutic exercises
- Assistive device and adaptive equipment training
- Orthotic or prosthetic training
- Manual therapy

“Occupational Therapy (OT)” refers to a therapeutic intervention program designed and supervised by a team of physicians and occupational therapists to assist members in regaining skills of daily living that have been lost or impaired. Such programs are individualized to each member to help improve quality of life by restoring independence. Occupational therapy may include, but is not limited to:

- Activities of Daily Living (ADL) training
- Muscle re-education
- Cognitive or neurodevelopmental training
- Perceptual motor training
- Fine motor coordination/strength training
- Assistive device and adaptive equipment training
- Environment modification recommendations and training
- Transfer training
- Functional mobility training

“Activities of Daily Living (ADLs)” are defined as routine activities that most healthy persons perform daily without requiring assistance. These include, but are not limited to: bathing, communication, transferring from bed to standing, wheelchair, or walker, dressing, feeding, grooming, mobility (ambulating), personal hygiene, self-maintenance, skin management, and toileting.

“Instrumental Activities of Daily Living (IADLs)” are defined as activities that may be performed daily but are not fundamental for daily functioning. These include, but are not limited to: the use of public

transportation, balancing a checkbook, community living activities, meal preparation, laundry, leisure activities and sports, and motor vehicle operation.

“Rehabilitative Treatments” are OT or PT treatments provided with the goal of restoring or improving upon functions that have been lost or impaired due to injury, disease, or congenital abnormality. Rehabilitative treatments are restorative and differentiated from “habilitative treatments” in that the individual has previously met these functional milestones but has lost them due to some process.

“Habilitative Treatments” are OT or PT treatments provided with the primary goal of developing skills needed to perform ADLs or IADLs which, as a result of injury, disease, or congenital abnormality, are not developed to the normal level of functioning. It is differentiated from rehabilitative treatment in that habilitative treatments are for individuals that have never met the initial development milestone.

“Custodial Care” or “Long-term Care” are non-skilled, personal care to maintain the member’s ADLs or IADLS over a long-term duration and do not require oversight or skilled services by trained health professionals or technical personnel. These services are not part of a medical treatment plan for recovery, rehabilitation, habilitation, or improvement in sickness or injury. Custodial services may be provided in the home, assisted living facilities, or nursing homes, etc. This type of custodial or long-term care typically does not apply for plan benefits, please see the member’s plan benefit.

Clinical Indications

General Criteria

Outpatient Physical Therapy and Occupational Therapy is considered for initial requests when ALL of the following criteria are met:

1. The treatment plan is prescribed and monitored by a licensed provider (MD, DO, PA, or NP) as per individual state law and must be provided by a licensed physical therapist or occupational therapist; *and*
2. Medical necessity criteria in the appropriate MCG Ambulatory Care > Rehabilitation > Physical or Occupational Therapy Services guideline is met; *and*
3. When relevant, medical necessity criteria in the appropriate MCG Ambulatory Care > Rehabilitation > Therapeutic Modalities guideline is met (e.g., aquatic therapy, transcutaneous electrical nerve stimulation, functional and neuromuscular electrical stimulation); *and*
4. Therapy is aimed at establishing or restoring function that was lost or impaired *as a result* of disease, injury, or procedure; *and*
5. Rehab potential is evident based on a review of the member’s condition, and the member’s function is not expected to improve in the absence of therapy; *and*
6. Improvement can be expected within 1 month for initial requests (appropriate to stage of injury, illness or disease) of beginning therapy and with sustainable benefit in range of motion, strength, function, reduced pain level, and independence of ADLs; *and*
7. The written plan of care includes an initial evaluation and is sufficient to determine the necessity of therapy, including ALL of the following elements:

- a. The diagnosis, the date of onset or exacerbation of the disorder/diagnosis, the duration, the severity, the anticipated course (stable, progressive or, improving), and the prognosis; *and*
 - b. Prior functioning level; *and*
 - c. Long-term (3 months) and short-term goals that are specific, quantitative, objective, and attainable in no more than 3 months; *and*
 - d. The frequency and duration of proposed treatment; *and*
 - e. The specific treatment techniques and/or exercises to be used; *and*
 - f. Education to help the member to self-manage and continue exercises and pain management program (e.g., TENS) without supervision; *and*
 - g. Discharge plan; *and*
 - h. Re-evaluation (documentation by a qualified provider/practitioner/physical or occupational therapist practicing within the scope of state specific licensure) performed at least monthly and the results as well as any proposed changes to address progress or lack thereof; *and*
8. Therapy is no more than 60 minutes per day, unless specifically justified and approved as part of the initial evaluation and treatment plan.

Duration of Therapy

The duration of therapy is dependent on the treatment plan of care and the severity of the member's condition. The Plan utilizes MCG rehabilitation criteria for the recommended visits per episode.

Extension Requests

A Plan member who requires continued PT or OT, beyond the original treatment plan of care, may receive extended treatment when ALL of the following criteria are met:

1. A completed re-evaluation documented by a qualified provider/practitioner/physical or occupational therapist practicing within the scope of state specific licensure, which has been conducted within 30 days of the service dates; *and*
2. The member has shown progress and improvement upon successful completion of the original treatment plan of care; *and*
3. Further significant improvement can be expected and continuation of PT or OT services must require the supervision of a licensed physical therapist; *and*
4. The written plan of continued care includes a complete history and documentation of progress from the original written plan of care, the member continues to have goals, and the elements noted above in the Clinical Indications section above.

Members may receive continued PT or OT equivalent to a maximum of 50% of the original treatment plan of care, with documentation and justification from the provider and may be subject to further review. An exception to the maximum amount of PT or OT can be made if medically necessary and determined by a qualified health professional(s) managing the Plan member's treatment plan.

Experimental or Investigational / Not Medically Necessary

Physical and occupational therapy should be discontinued when any ONE of the following is present:

- The member reaches the predetermined goals or skilled treatment is no longer required; *or*
- The member has reached maximum rehab potential; *or*
- The goals will not be met and there is no expectation of meeting them in reasonable time; *or*
- The member can safely and effectively continue their rehabilitation in a home exercise program or self-management program (maintenance); *or*
- The member's medical condition prevents further therapy; *or*
- The member refuses treatment.

Physical/Occupational Therapy is NOT considered medically necessary for the following:

- Asymptomatic members or those without an identifiable clinical condition; *or*
- Improvement in functioning is not expected over a reasonable and predictable period of time (i.e., a "stable deficit"); *or*
- Cases of transient or easily reversible loss or reduction in function which could be reasonably expected to improve spontaneously as the member gradually resumes normal activities; *or*
- Long-term maintenance therapy, as it is aimed to preserve the present level of function or to prevent regression below an acceptable level of functioning; *or*
- Custodial or long-term care services; *or*
- General exercises to promote fitness or flexibility are not medically necessary; *or*
- Duplicative therapy services or programs; *or*
- Occupational or recreational programs aiming to augment or improve upon normal human functioning; this includes services considered as routine, educational, for employment or job training, or as part of a fitness or sports program; *or*
- Sports Rehabilitation where treatment is extended what is needed to improve above and beyond the normal ability to perform activities of daily living
- Vertebral axial decompression, as they are considered experimental or investigational, including but not limited to the following devices:
 - Decompression Reduction Stabilization (DRS) System
 - DRX 9000
 - DX2 Decompression System
 - IDD Therapy (Intervertebral Differential Dynamics Therapy)
 - Integrity Spinal Care System
 - Lordex Lumbar Spine System
 - MTD 4000 Mettler Traction Decompression System
 - SpineRx-LDM
 - VAX-D Spinal Decompression System
- Treatment modalities where the benefits of PT/OT are not adequately supported by peer literature or accepted standards of practice to be safe and effective and therefore, considered experimental or investigational include, but are not limited to:
 - Applied Functional Science

- Augmented soft tissue mobilization is considered investigational due to limited evidence of improved outcomes over standard techniques for soft tissue mobilization
- Biofeedback for anal/perianal/urethral sphincters due to lack of high grade quality of evidence to support at this time.
- Driver/safety training
- Equestrian therapy (hippotherapy)
- Group therapy (criteria require *individualized* plans)
- Hands-Free Ultrasound and Low-Frequency Sound (Infrasound)
- Hivamat therapy (deep oscillation therapy)
- Interferential stimulation or interferential current therapy is the superficial application of a medium-frequency alternating current, modulated to produce low frequencies up to 150 Hz and is considered experimental and investigational due to the lack of high grade quality literature. Devices such as neoGEN-Series System (RST-Sanexas) as a form of interferential current therapy for neuropathic pain provide ultra-high digital frequency to produce pulsed electronic signal energy waves that are delivered transcutaneously via contact electrodes. As per Hayes 2023, there are no relevant clinical studies, systematic reviews, or guidance documents supporting use of RST-Sanexas.
- Kinesio taping for back pain or radicular pain is considered investigational and not clearly established in the literature
- Low level laser therapy (LLLT)
- Microcurrent electrical nerve stimulation (MENS)
- Percutaneous electrical nerve stimulation (PENS) involves electroacupuncture (needle electrodes connected to a power source) and inserted into the skin near the site of pain. These needles are not permanently implanted, but tunneled percutaneously under ultrasound guidance into the subcutaneous tissue. Due to a lack of peer-reviewed reviewed literature supporting the effectiveness of PENS and pain reduction, ongoing clinical trials, and clinical practice guidelines stating there is insufficient evidence to recommend PENS for lower back pain, this type of procedure is considered experimental or investigational.
- Pilates, Tai Chi and Qi Gong
- Sensory integrative techniques
- The Interactive Metronome Program
- MEDEK Therapy
- McKenzie Method of Mechanical Diagnosis and Therapy
- Conditions where the benefits of PT/OT are not adequately supported by peer literature include, but are not limited to:
 - Constipation
 - Vaginismus
 - Social functioning
 - Sexual dysfunction (erectile dysfunction, premature ejaculation), extracorporeal shockwave therapy for erectile dysfunction
 - Scoliosis

- TMJ pain

Applicable Billing Codes (HCPCS/CPT Codes)

CPT/HCPCS Codes considered medically necessary if criteria are met:	
<i>Code</i>	<i>Description</i>
94667	Manipulation chest wall, such as cupping, percussing, and vibration to facilitate lung function; initial demonstration and/or evaluation
94668	Manipulation chest wall, such as cupping, percussing, and vibration to facilitate lung function; subsequent
97039	Unlisted modality (specify type and time if constant attendance)
97010	Application of a modality to one or more areas; hot or cold packs
97012	Application of a modality to one or more areas; traction, mechanical
97014	Application of a modality to one or more areas; electrical stimulation (unattended)
97016	Application of a modality to one or more areas; vasopneumatic devices
97018	Application of a modality to one or more areas; paraffin bath
97022	Application of a modality to one or more areas; whirlpool
97024	Application of a modality to one or more areas; diathermy (e.g., microwave)
97026	Application of a modality to one or more areas; infrared
97028	Application of a modality to one or more areas; ultraviolet
97032	Application of a modality to one or more areas; electrical stimulation (manual), each 15 minutes
97033	Application of a modality to one or more areas; iontophoresis, each 15 minutes
97034	Application of a modality to one or more areas; contrast baths, each 15 minutes
97035	Application of a modality to one or more areas; ultrasound, each 15 minutes
97036	Application of a modality to one or more areas; Hubbard tank, each 15 minutes

97110	Therapeutic procedure, one or more areas, each 15 minutes; therapeutic exercises to develop strength and endurance, range of motion and flexibility
97112	Therapeutic procedure, one or more areas, each 15 minutes; neuromuscular reeducation of movement, balance, coordination, kinesthetic sense, posture, and/or proprioception for sitting and/or standing activities
97113	Therapeutic procedure, one or more areas, each 15 minutes; aquatic therapy with therapeutic exercise
97116	Therapeutic procedure, one or more areas, each 15 minutes; Gait training (includes stair climbing)
97124	Therapeutic procedure, one or more areas, each 15 minutes; massage, including effleurage, petrissage and/or tapotement (stroking compression, percussion)
97129	Therapeutic interventions that focus on cognitive function (eg, attention, memory, reasoning, executive function, problem solving, and/or pragmatic functioning) and compensatory strategies to manage the performance of an activity (eg, managing time or schedules, initiating, organizing, and sequencing tasks), direct (one-on-one) patient contact; initial 15 minutes
97130	Therapeutic interventions that focus on cognitive function (eg, attention, memory, reasoning, executive function, problem solving, and/or pragmatic functioning) and compensatory strategies to manage the performance of an activity (eg, managing time or schedules, initiating, organizing, and sequencing tasks), direct (one-on-one) patient contact; each additional 15 minutes (List separately in addition to code for primary procedure)
97140	Manual therapy techniques (e.g., mobilization/manipulation, manual lymphatic drainage, manual traction), one or more regions, each 15 minutes
97161	Physical therapy evaluation: low complexity, requiring these components: A history with no personal factors and/or comorbidities that impact the plan of care; An examination of body system(s) using standardized tests and measures addressing 1-2 elements from any of the following: body structures and functions, activity limitations, and/or participation restrictions; A clinical presentation with stable and/or uncomplicated characteristics; and Clinical decision making of low complexity using standardized patient assessment instrument and/or measurable assessment of functional outcome. Typically, 20 minutes are spent face-to-face with the patient and/or family.

97162	Physical therapy evaluation: moderate complexity, requiring these components: A history of present problem with 1-2 personal factors and/or comorbidities that impact the plan of care; An examination of body systems using standardized tests and measures in addressing a total of 3 or more elements from any of the following: body structures and functions, activity limitations, and/or participation restrictions; An evolving clinical presentation with changing characteristics; and Clinical decision making of moderate complexity using standardized patient assessment instrument and/or measurable assessment of functional outcome. Typically, 30 minutes are spent face-to-face with the patient and/or family.
97163	Physical therapy evaluation: high complexity, requiring these components: A history of present problem with 3 or more personal factors and/or comorbidities that impact the plan of care; An examination of body systems using standardized tests and measures addressing a total of 4 or more elements from any of the following: body structures and functions, activity limitations, and/or participation restrictions; A clinical presentation with unstable and unpredictable characteristics; and Clinical decision making of high complexity using standardized patient assessment instrument and/or measurable assessment of functional outcome. Typically, 45 minutes are spent face-to-face with the patient and/or family.
97164	Re-evaluation of physical therapy established in plan of care
97165	Occupational therapy evaluation, low complexity, requiring these components: An occupational profile and medical and therapy history, which includes a brief history including review of medical and/or therapy records relating to the presenting problem; An assessment(s) that identifies 1-3 performance deficits (ie, relating to physical, cognitive, or psychosocial skills) that result in activity limitations and/or participation restrictions; and Clinical decision making of low complexity, which includes an analysis of the occupational profile, analysis of data from problem-focused assessment(s), and consideration of a limited number of treatment options. Patient presents with no comorbidities that affect occupational performance. Modification of tasks or assistance (eg, physical or verbal) with assessment(s) is not necessary to enable completion of evaluation component. Typically, 30 minutes are spent face-to-face with the patient and/or family.
97166	Occupational therapy evaluation, moderate complexity, requiring these components: An occupational profile and medical and therapy history, which includes an expanded review of medical and/or therapy records and additional review of physical, cognitive, or psychosocial history related to current functional

	<p>performance; An assessment(s) that identifies 3-5 performance deficits (ie, relating to physical, cognitive, or psychosocial skills) that result in activity limitations and/or participation restrictions; and Clinical decision making of moderate analytic complexity, which includes an analysis of the occupational profile, analysis of data from detailed assessment(s), and consideration of several treatment options. Patient may present with comorbidities that affect occupational performance. Minimal to moderate modification of tasks or assistance (eg, physical or verbal) with assessment(s) is necessary to enable patient to complete evaluation component. Typically, 45 minutes are spent face-to-face with the patient and/or family.</p>
97167	<p>Occupational therapy evaluation, high complexity, requiring these components: An occupational profile and medical and therapy history, which includes review of medical and/or therapy records and extensive additional review of physical, cognitive, or psychosocial history related to current functional performance; An assessment(s) that identifies 5 or more performance deficits (ie, relating to physical, cognitive, or psychosocial skills) that result in activity limitations and/or participation restrictions; and Clinical decision making of high analytic complexity, which includes an analysis of the patient profile, analysis of data from comprehensive assessment(s), and consideration of multiple treatment options. Patient presents with comorbidities that affect occupational performance. Significant modification of tasks or assistance (eg, physical or verbal) with assessment(s) is necessary to enable patient to complete evaluation component. Typically, 60 minutes are spent face-to-face with the patient and/or family.</p>
97168	<p>Re-evaluation of occupational therapy established for plan of care</p>
97530	<p>Therapeutic activities, direct (one-on-one) patient contact (use of dynamic activities to improve functional performance), each 15 minutes</p>
97533	<p>Sensory integrative techniques to enhance sensory processing and promote adaptive responses to environmental demands, direct (one-on-one) patient contact, each 15 minutes</p>
97535	<p>Self care/home management training (e.g., activities of daily living [ADL] and compensatory training, meal preparation, safety procedures, and instructions in use of assistive technology devices/adaptive equipment) direct one-on-one contact by provider, each 15 minutes</p>
97542	<p>Wheelchair management (e.g., assessment, fitting, training), each 15 minutes</p>

97750	Physical performance test or measurement (e.g., musculoskeletal functional capacity), with written report, each 15 minutes
97760	Orthotic(s) management and training (including assessment and fitting when not otherwise reported), upper extremity(ies), lower extremity(ies) and/or trunk, initial orthotic(s) encounter, initial orthotic(s) encounter, each 15 minutes
97761	Prosthetic(s) training, upper and/or lower extremity(ies), initial prosthetic(s) encounter, each 15 minutes
97763	Orthotic(s)/prosthetic(s) management and/or training, upper extremity(ies), lower extremity(ies), and/or trunk, subsequent orthotic(s)/prosthetic(s) encounter, each 15 minutes
97799	Unlisted physical medicine/rehabilitation service or procedure
A4558	Conductive gel or paste, for use with electrical device (e.g., TENS, NMES), per oz
A4595	Electrical stimulator supplies, 2 lead, per month, (e.g., TENS, NMES)
E0720	Transcutaneous electrical nerve stimulation (TENS) device, two-lead, localized stimulation
E0730	Transcutaneous electrical nerve stimulation (TENS) device, four or more leads, for multiple nerve stimulation
E0731	Form-fitting conductive garment for delivery of TENS or NMES (with conductive fibers separated from the patient's skin by layers of fabric)
E0745	Neuromuscular stimulator, electronic shock unit
G0129	Occupational therapy requiring the skills of a qualified occupational therapist, furnished as a component of a partial hospitalization treatment program, per day
G0281	Electrical stimulation (unattended), to one or more areas, for chronic stage III and stage IV pressure ulcers, arterial ulcers, diabetic ulcers, and venous stasis ulcers not demonstrating measurable signs of healing after 30 days of conventional care, as part of a therapy plan of care
G0282	Electrical stimulation (unattended), to one or more areas, for wound care, other than described in G0281

G0283	Electrical stimulation (unattended), to one or more areas for indication(s) other than wound care, as part of a therapy plan of care
-------	--

CPT/HCPCS codes <i>not</i> considered medically necessary:	
<i>Code</i>	<i>Description</i>
90912	Biofeedback training, perineal muscles, anorectal or urethral sphincter, including EMG and/or manometry, when performed; initial 15 minutes of one-on-one physician or other qualified health care professional contact with the patient
90913	Biofeedback training, perineal muscles, anorectal or urethral sphincter, including EMG and/or manometry, when performed; each additional 15 minutes of one-on-one physician or other qualified health care professional contact with the patient (List separately in addition to code for primary procedure)
97169	Athletic training evaluation, low complexity, requiring these components: A history and physical activity profile with no comorbidities that affect physical activity; An examination of affected body area and other symptomatic or related systems addressing 1-2 elements from any of the following: body structures, physical activity, and/or participation deficiencies; and Clinical decision making of low complexity using standardized patient assessment instrument and/or measurable assessment of functional outcome. Typically, 15 minutes are spent face-to-face with the patient and/or family
97170	Athletic training evaluation, moderate complexity, requiring these components: A medical history and physical activity profile with 1-2 comorbidities that affect physical activity; An examination of affected body area and other symptomatic or related systems addressing a total of 3 or more elements from any of the following: body structures, physical activity, and/or participation deficiencies; and Clinical decision making of moderate complexity using standardized patient assessment instrument and/or measurable assessment of functional outcome. Typically, 30 minutes are spent face-to-face with the patient and/or family.
97171	Athletic training evaluation, high complexity, requiring these components: A medical history and physical activity profile, with 3 or more comorbidities that affect physical activity; A comprehensive examination of body systems using standardized tests and measures addressing a total of 4 or more elements from any of the following: body structures, physical activity, and/or participation deficiencies; Clinical presentation with unstable and unpredictable characteristics; and Clinical decision making of high complexity using

	standardized patient assessment instrument and/or measurable assessment of functional outcome. Typically, 45 minutes are spent face-to-face with the patient and/or family.
97172	Re-evaluation of athletic training established plan of care requiring these components: An assessment of patient's current functional status when there is a documented change; and A revised plan of care using a standardized patient assessment instrument and/or measurable assessment of functional outcome with an update in management options, goals, and interventions. Typically, 20 minutes are spent face-to-face with the patient and/or family.
97537	Community/work reintegration training (e.g., shopping, transportation, money management, avocational activities and/or work environment/modification analysis, work task analysis, use of assistive technology device/adaptive equipment), direct one-on-one contact by provider, each 15 minutes
97545	Work hardening/conditioning; initial 2 hours
97546	Work hardening/conditioning; each additional hour (List separately in addition to code for primary procedure)
E0746	Electromyography (EMG), biofeedback device
E0769	Electrical stimulation or electromagnetic wound treatment device, not otherwise classified
S8990	Physical or manipulative therapy performed for maintenance rather than restoration
S9117	Back school, per visit
S9970	Health club membership, annual

CPT/HCPCS codes considered experimental or investigational:	
<i>Code</i>	<i>Description</i>
97150	Therapeutic procedure(s), group (2 or more individuals)
S8940	Equestrian/hippotherapy, per session

S8948	Application of a modality (requiring constant provider attendance) to one or more areas, low-level laser; each 15 minutes
S9090	Vertebral axial decompression, per session

CPT/HCPCS codes for electrical stimulation considered experimental or investigational (Interferential therapy, MENS, PENS):	
<i>Code</i>	<i>Description</i>
64999	Unlisted procedure, nervous system <ul style="list-style-type: none"> • <u>Due to the broad nature of this code and lack of specificity in certain scenarios, clarification is provided below:</u> • When this code is billed for percutaneous electrical nerve stimulation (PENS), it is considered experimental or investigational
E1399	Durable medical equipment, miscellaneous <ul style="list-style-type: none"> • <u>Due to the broad nature of this code and lack of specificity in certain scenarios, clarification is provided below:</u> • When this code is billed for microcurrent electrical nerve stimulation (MENS), it is considered experimental or investigational
S8130	Interferential current stimulator, 2 channel
S8131	Interferential current stimulator, 4 channel

References

1. The American College of Obstetrics and Gynecologists. (July 2019). Female Sexual Dysfunction. Practice Bulletin Number 213.
<https://www.acog.org/clinical/clinical-guidance/practice-bulletin/articles/2019/07/female-sexual-dysfunction>
2. American Occupational Therapy Association. (2021). Standards of practice for occupational therapy. The American Journal of Occupational Therapy, 75(Supplement_3).
3. American Physical Therapy Association (APTA). Guide to Physical Therapist Practice, 3rd Revised Edition. Updated November 2016.
4. American Physical Therapy Association. Standards of practice for physical therapy. Last updated Aug 12,2020. Available at:
<https://www.apta.org/apta-and-you/leadership-and-governance/policies/standards-of-practice-p>
t
5. Bavarian R, Khawaja SN, Ajisafe AH, Sultan AS. The efficacy of microcurrent electrical nerve stimulation in treating masticatory myofascial pain: A systematic review and meta-analysis. *The*

Journal of Craniomandibular & Sleep Practice. 2021;
doi: <https://doi.org/10.1080/08869634.2021.2020006>

6. Bizzarri P, Buzzatti L, Cattysse E, Scafoglieri A. Thoracic manual therapy is not more effective than placebo thoracic manual therapy in patients with shoulder dysfunctions: A systematic review with meta-analysis. *Musculoskeletal Science and Practice*. 2018;33:1-10.
doi:10.1016/j.msksp.2017.10.006
7. Beltran-Alacreu H, Serrano-Muñoz D, Martín-Caro Álvarez D, Fernández-Pérez JJ, Gómez-Soriano J, Avendaño-Coy J. Percutaneous Versus Transcutaneous Electrical Nerve Stimulation for the Treatment of Musculoskeletal Pain. A Systematic Review and Meta-Analysis. *Pain Med*. 2022 Aug 1;23(8):1387-1400. doi: 10.1093/pm/pnac027. PMID: 35167691.
8. Burnett AL, Nehra A, Breau RH et al: Erectile dysfunction: AUA guideline. *J Urol* 2018; 200: 633.
9. Collins N, Barton C, van Middelkoop M et al. 2018 Consensus statement on exercise therapy and physical interventions (orthoses, taping and manual therapy) to treat patellofemoral pain: recommendations from the 5th International Patellofemoral Pain Research Retreat, Gold Coast, Australia, 2017. *Br J Sports Med*. 2018;52(18):1170-1178. doi:10.1136/bjsports-2018-099397
10. Deer, T., Shah, A., Slavin, K., Vorenkamp, K. E., Shah, S., Leong, M., & McRoberts, W. P. (2023). Birds of a Feather Redux: Defining Ways to Stimulate the Peripheral Nervous System. *Journal of Pain Research*, 1219-1224.
11. de Freitas, G. R., Szpoganicz, C., & Ilha, J. (2018). Does neuromuscular electrical stimulation therapy increase voluntary muscle strength after spinal cord injury? A systematic review. *Topics in spinal cord injury rehabilitation, Top Spinal Cord Inj Rehabil*. 24(1), 6-17.
12. De Oliveira, G.R., & Vidal MF. (2021). Developmental outcomes in a child with corpus callosum abnormalities and congenital heart disease after Cuevas Medek Exercises: A case report. *Clinical Case Reports*. Doi: <https://doi.org/10.1002/ccr3.4637>
13. Department of Veterans Affairs. (2022). VA/DoD CLINICAL PRACTICE GUIDELINE FOR DIAGNOSIS AND TREATMENT OF LOW BACK PAIN. Version 3.0. Retrieved from <https://www.healthquality.va.gov/guidelines/Pain/lbp/VADoDLBPCPGFinal508.pdf>
14. Scherl SA. Adolescent idiopathic scoliosis: Management and prognosis. UpToDate. <https://www.uptodate.com/contents/adolescent-idiopathic-scoliosis-management-and-prognosis> Accessed September 23, 2017.
15. McNeely ML, Armijo Olivo S, Magee DJ. A systematic review of the effectiveness of physical therapy interventions for temporomandibular disorders. *Phys Ther* 2006; 86:710.
16. Craane B, Dijkstra PU, Stappaerts K, De Laat A. Randomized controlled trial on physical therapy for TMJ closed lock. *J Dent Res* 2012; 91:364.
17. Scrivani SJ. Temporomandibular disorders in adults. UpToDate. <https://www.uptodate.com/contents/temporomandibular-disorders-in-adults>. Accessed September 23, 2017.
18. King, PM, Myers, CA, Ling, FW, Rosenthal, RH. Musculoskeletal factors in chronic pelvic pain. *J Psychosom Obstet Gynaecol* 1991; 12:87.
19. Rampazo ÉP, Liebano RE. Analgesic Effects of Interferential Current Therapy: A Narrative Review. *Medicina*. 2022; 58(1):141. <https://doi.org/10.3390/medicina58010141>

20. Romano M, Minozzi S, Bettany-Saltikov J, et al. Exercises for adolescent idiopathic scoliosis. *Cochrane Database Syst Rev* 2012; :CD007837.
21. Bylund P, Aaro S, Gottfries B, Jansson E. Is lateral electric surface stimulation an effective treatment for scoliosis? *J Pediatr Orthop* 1987; 7:298.
22. Wong MS, Mak AF, Luk KD, et al. Effectiveness of audio-biofeedback in postural training for adolescent idiopathic scoliosis patients. *Prosthet Orthot Int* 2001; 25:60.
23. Melham TJ, Sevier TL, Malnofski MJ, et al. Chronic ankle pain and fibrosis successfully treated with a new noninvasive augmented soft tissue mobilization technique (ASTM): A case report. *Med Sci Sports Exerc.* 1998;30(6):801-804.
24. Cheatham SW, Lee M, Cain M, Baker R. The efficacy of instrument assisted soft tissue mobilization: a systematic review. *J Can Chiropr Assoc.* 2016;60(3):200-211.
25. Halseth T, McChesney JW, DeBeliso M, et al. The Effects of Kinesio Taping on proprioception at the ankle. *J Sports Sci Med.* 2004;3:1-7.
26. Yasukawa A, Patel P, Sisung C. Pilot study: Investigating the effects of Kinesio Taping in an acute pediatric rehabilitation setting. *Am J Occup Ther.* 2006;60(1):104-110.
27. Fu TC, Wong AM, Pei YC, et al. Effect of Kinesio taping on muscle strength in athletes-a pilot study. *J Sci Med Sport.* 2008;11(2):198-201.
28. Thelen MD, Dauber JA, Stoneman PD. The clinical efficacy of kinesio tape for shoulder pain: A randomized, double-blinded, clinical trial. *J Orthop Sports Phys Ther.* 2008;38(7):389-395.
29. Clarke CE, Patel S, Ives N, et al. Clinical effectiveness and cost-effectiveness of physiotherapy and occupational therapy versus no therapy in mild to moderate Parkinson's disease: A large pragmatic randomised controlled trial (PD REHAB). *Health Technol Assess.* 2016;20(63):1-96.
30. García-Collado A, Valera-Calero JA, Fernández-de-Las-Peñas C, Arias-Burúa JL. Effects of ultrasound-guided nerve stimulation targeting peripheral nerve tissue on pain and function: a scoping review. *J Clin Med.* 2022;11(13):3753. doi:10.3390/jcm11133753
31. Gibson, W., Wand, B. M., Meads, C., Catley, M. J., & O'Connell, N. E. (2019). Transcutaneous electrical nerve stimulation (TENS) for chronic pain-an overview of Cochrane Reviews. *Cochrane Database of Systematic Reviews.* doi: 10.1002/14651858.CD011890.pub3
32. Hayes, Inc. Evidence Analysis Research Brief. *Percutaneous Electrical Nerve Stimulation for Treatment of Low Back Pain.* Lansdale, PA: Hayes, Inc.; Dec 22, 2022.
33. Hidalgo B, Hall T, Bossert J, Dugeny A, Cagnie B, Pitance L. The efficacy of manual therapy and exercise for treating non-specific neck pain: A systematic review. *J Back Musculoskelet Rehabil.* 2018;30(6):1149-1169. doi:10.3233/bmr-169615
34. LaBan MM, Martin T, Pechur J, et al. Physical and occupational therapy in the treatment of patients with multiple sclerosis. *Phys Med Rehabil Clin N Am.* 1998;9(3):603-614, vii.
35. Nordin M, Campello M. Physical therapy: exercises and the modalities: when, what, and why? *Neurol Clin.* 1999;17(1):75-89.
36. Hicks JE. Role of rehabilitation in the management of myopathies. *Curr Opin Rheumatol.* 1998;10(6):548-555.
37. Medicare Claims Processing Manual, Chapter 5 – Part B Outpatient Rehabilitation and CORF/OPT Services. CMS.gov.

- <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/downloads/clm104c05.pdf>. Revised December 2, 2016. Accessed September 10, 2017.
38. Rosenquist, EWK. Overview of the treatment of chronic non-cancer pain. UpToDate, Inc. Waltham, MA. Updated August 2017. Accessed September 15, 2017.
 39. Hoenig, H. Overview of geriatric rehabilitation: Program components and settings for rehabilitation. UpToDate, Inc. Waltham, MA. Updated August 2017. Accessed September 15, 2017.
 40. Minor MA, Sanford MK. The role of physical therapy and physical modalities in pain management. *Rheum Dis Clin North Am*. 1998;25(1):233-248, viii.
 41. McKivigan, J., Krisak, B., Vogel, B., Harrington, A., & Nguyen, V. (2019). Evaluating the Efficacy of Vertebral Axial Decompression Therapy in Treatment of Patients with Chronic Lower Back Pain: A Systematic Review. *Rehabilitation*, 4(2), 25-34.
 42. Zhou K, Krug K, Brogan MS. Physical therapy in wound care: A cost-effectiveness analysis. *Medicine (Baltimore)*. 2015;94(49):e2202.
 43. Centers for Medicare and Medicaid. Medicare Benefit Policy Manual Chapter 15 – Covered Medical and Other Health Services (Rev. 10880, 08-06-21). Retrieved from: <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c15.pdf>
 44. CMS LCD Guidelines. <http://www.cms.gov/mcd/index>. Accessed on September 10, 2016.
 - a. Local Coverage Determination (LCD) for PHYSICAL THERAPY - Home Health (L33942)
 - b. Local Coverage Determination (LCD) for PHYSICAL THERAPY - Home Health (L34564)
 - c. Local Coverage Determination (LCD) for Outpatient PHYSICAL and Occupational THERAPY Services Health (L34049)
 - d. Local Coverage Determination (LCD) for Medicine: PHYSICAL THERAPY - Outpatient (L34310)
 45. Frontera W, Silver J, Rizzo TD editors. *Essentials of Physical Medicine and Rehabilitation*. 4th ed. Philadelphia, PA: Elsevier, 2018.
 46. Gulick DT. Comparison of tissue heating between manual and hands-free ultrasound techniques. *Physiother Theory Pract*. 2010;26(2):100-106.
 47. Katalinic OM, Harvey LA, Herbert RD. Effectiveness of stretch for the treatment and prevention of contractures in people with neurological conditions: A systematic review. *Phys Ther*. 2011;91(1):11-24.
 48. Atamaz FC, Durmaz B, Baydar M, et al. Comparison of the efficacy of transcutaneous electrical nerve stimulation, interferential currents, and shortwave diathermy in knee osteoarthritis: a double-blind, randomized, controlled, multicenter study. *Arch Phys Med Rehabil*. 2012 May;93(5):748-56.
 49. Clare HA, Adams R, Maher CG. A systematic review of efficacy of McKenzie therapy for spinal pain. *Aust J Physiother*. 2004;50(4):209-216.
 50. Garcia AN, Costa Lda C, Hancock MJ, et al. Efficacy of the McKenzie method in patients with chronic nonspecific low back pain: A protocol of randomized placebo-controlled trial. *Phys Ther*. 2015;95(2):267-273.

51. Moss M, Nordon-Craft A, Malone D, et al. A randomized trial of an intensive physical therapy program for acute respiratory failure patients. *Am J Respir Crit Care Med*. 2016;193(10):1101-1110.
52. Klugarova J, Klugar M, Mareckova J, et al. The effectiveness of inpatient physical therapy compared to outpatient physical therapy in older adults after total hip replacement in the post-discharge period: A systematic review. *JBIC Database System Rev Implement Rep*. 2016;14(1):174-209.
53. Legg LA, Drummond AE, Langhorne P. Occupational therapy for patients with problems in activities of daily living after stroke. *Cochrane Database Syst Rev*. 2006;(4):CD003585.
54. National Institute for Health and Care Excellence. (2016). Low back pain and sciatica in over 16s: assessment and management [NG59]. Retrieved from <https://www.nice.org.uk/guidance/ng59>
55. Sakzewski L, Ziviani J, Boyd R. Systematic review and meta-analysis of therapeutic management of upper-limb dysfunction in children with congenital hemiplegia. *Pediatrics*. 2009 Jun;123(6):e1111-22.
56. Shank TM, & Harron W. A retrospective outcomes study examining the effect of interactive metronome on hand function. *Journal of Hand Therapy*. 2015; 28 (4): 396-402. Doi: <https://doi.org/10.1016/j.jht.2015.06.003>
57. Rajpurohit B, Khatri SM, Metgud D, Bagewadi A. Effectiveness of transcutaneous 38 electrical nerve stimulation and microcurrent electrical nerve stimulation in bruxism 39 associated with masticatory muscle pain--a comparative study. *Indian J Dent Res*. 2010 40 Jan-Mar;21(1):104-6.
58. Reitz SM, Austin DJ, Brandt LC, et al. Guidelines to the Occupational Therapy Code of Ethics. *American Journal Occupational Therapy*. 2006; 60(6):652-668.
59. Steultjens EM, Dekker JJ, Bouter LM, et al. Evidence of the efficacy of occupational therapy in different conditions: an overview of systematic reviews. *Clin Rehabil*. 2005; 19(3):247-254.
60. Steultjens EM, Dekker JJ, Bouter LM, et al. Occupational therapy for rheumatoid arthritis. *Cochrane Database of Syst Rev*. 2004;(1):CD003114.
61. STECKO, Monika Joanna, WAWRYKÓW, Agata, KORABIUSZ, Katarzyna, KORDEK, Agnieszka, FABIAN - DANIELEWSKA, Anna, WILCZYŃSKA, Agnieszka, JANIK, Inga & MACIEJEWSKA, Martyna. The methods of therapy of children suffering from cerebral palsy. *Journal of Education, Health and Sport* [online]. 9 September 2018, T. 8, nr 9, s. 562–569. [accessed March 9, 2022].
62. Thomaz, S. R., Cipriano Jr, G., Formiga, M. F., Fachin-Martins, E., Cipriano, G. F. B., Martins, W. R., & Cahalin, L. P. (2019). Effect of electrical stimulation on muscle atrophy and spasticity in patients with spinal cord injury—a systematic review with meta-analysis. *Spinal cord*, 57(4), 258-266.
63. Toth MJ, Tourville TW, Voight TB, et al. Utility of Neuromuscular Electrical Stimulation to Preserve Quadriceps Muscle Fiber Size and Contractility After Anterior Cruciate Ligament Injuries and Reconstruction: A Randomized, Sham-Controlled, Blinded Trial. *The American Journal of Sports Medicine*. Sage Journals. 2020; 48 (10), p: 2429-2437. Doi:<https://doi.org/10.1177/0363546520933622>

64. Trombly CA, Ma HI. A synthesis of the effects of occupational therapy for persons with stroke, Part I: Restoration of roles, tasks, and activities. *American Journal Occupational Therapy*. 2002;56(3):250-9.
65. Ma HI, Trombly CA. A synthesis of the effects of occupational therapy for persons with stroke. Part II: Remediation of impairments. *American Journal Occupational Therapy*. 2002;56(3):260-274.
66. Namnaqani, F. I., Mashabi, A. S., Yaseen, K. M., & Alshehri, M. A. (2019). The effectiveness of McKenzie method compared to manual therapy for treating chronic low back pain: a systematic review. *Journal of Musculoskeletal & Neuronal Interactions*, 19(4), 492.
67. Nair HKR. Microcurrent as an adjunct therapy to accelerate chronic wound healing and 2 reduce patient pain. *J Wound Care*. 2018 May 2;27(5):296-306.
68. Walker MF, Leonardi-Bee J, Bath P, Langhorne P, Dewey M, Corr S, et al. Individual patient data metaanalysis of randomized controlled trials of community occupational therapy for stroke patients. *Stroke*. 2004 Sep;35(9):2226-32.
69. Hesselstrand M, Samuelsson K, Liedberg G. Occupational therapy interventions in chronic pain - A systematic review. *Occupational Therapy International*. 2015;22(4):183-194.
70. Kuhaneck HM, Madonna S, Novak A, Pearson E. Effectiveness of interventions for children with autism spectrum disorder and their parents: A systematic review of family outcomes. *American Journal Occupational Therapy*. 2015;69(5):6905180040p1-14.
71. Clarke CE, Patel S, Ives N, et al. Clinical effectiveness and cost-effectiveness of physiotherapy and occupational therapy versus no therapy in mild to moderate Parkinson's disease: A large pragmatic randomised controlled trial (PD REHAB). *Health Technol Assess*. 2016;20(63):1-96.
72. Qaseem A, et al. Noninvasive Treatments for Acute, Subacute, and Chronic Low Back Pain: A Clinical Practice Guideline From the American College of Physicians. 2017 Apr.
73. Szulc P, Wendt M, Waszak M, Tomczak M, Cieřlik K, Trzaska T. Impact of McKenzie Method Therapy Enriched by Muscular Energy Techniques on Subjective and Objective Parameters Related to Spine Function in Patients with Chronic Low Back Pain. *Med Sci Monit*. 2015;21:2918–2932. doi:10.12659/MSM.894261
74. Lam OT, Strenger DM, Chan-Fee M, et al. Effectiveness of the McKenzie Method of Mechanical Diagnosis and Therapy for Treating Low Back Pain: Literature Review With Meta-analysis *Journal of Orthopaedic & Sports Physical Therapy*. 2018;48(6):476-490.
75. Laver K, Dyer S, Whitehead C, et al. Interventions to delay functional decline in people with dementia: a systematic review of systematic reviews. *BMJ Open*. 2016;6(4):e010767. doi: 10.1136/bmjopen-2015-010767.
76. Lee HS, Park SW, Park YJ. Effects of physical activity programs on the improvement of dementia symptom: a meta-analysis. *Biomed Res Int*. 2016;2016:2920146. Epub 2016 Oct 13.
77. Forbes D, Forbes SC, Blake CM, et al. Exercise programs for people with dementia (review). *Cochrane Database Syst Rev*. 2015;(4). doi: 10.1002/14651858.CD006489.pub4.
78. Winstein CJ, Stein J, Arena R, et al. Guidelines for Adult Stroke Rehabilitation and Recovery A Guideline for Healthcare Professionals From the American Heart Association/American Stroke Association. *Stroke*. 2016;47:e98–e169. doi: 10.1161/STR.0000000000000098

79. Giggins OM, Persson UM, Caulfield B. Biofeedback in rehabilitation. *Journal of NeuroEngineering and Rehabilitation*. 2013;10(60). doi: 10.1186/1743-0003-10-60
80. Young, C., & Argáez, C. (2020). *Manual Therapy for Chronic Non-Cancer Back and Neck Pain: A Review of Clinical Effectiveness*. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK562937/>
81. Yu GH, Lee JS, Kim SK, & Cha TH. (2017). Effects of interactive metronome training on upper extremity function, ADL and QOL in stroke patients. *NeuroRehabilitation*. 2017; 41(1), pp. 161-168. DOI: 10.3233/NRE-171468
82. Hayes, Inc. Evolving Evidence Review. NeoGEN-Series System (RST-Sanexas) for Treatment of Neuropathic Pain. Lansdale, PA: Hayes, Inc.; Annual review: Jan 2023.
83. Chou R. Subacute and chronic low back pain: Nonpharmacologic and pharmacologic treatment. UpToDate.com. Accessed from https://www.uptodate.com/contents/subacute-and-chronic-low-back-pain-nonpharmacologic-and-pharmacologic-treatment?search=interferential%20current%20therapy&source=search_result&selectedTitle=2~150&usage_type=default&display_rank=2 Last updated Aug 2, 2022.
84. Nazligul T, Akpınar P, Aktas I, Unlu Ozkan F, Cagliyan Hartevioglu H. The effect of interferential current therapy on patients with subacromial impingement syndrome: a randomized, double-blind, sham-controlled study. *Eur J Phys Rehabil Med*. 2018 Jun;54(3):351-357. doi: 10.23736/S1973-9087.17.04743-8. Epub 2017 Sep 11. PMID: 28895673.
85. van Reijn-Baggen DA, Han-Geurts IJM, Voorham-van der Zalm PJ, Pelger RCM, Hagenaars-van Miert CHAC, Laan ETM. Pelvic Floor Physical Therapy for Pelvic Floor Hypertonicity: A Systematic Review of Treatment Efficacy. *Sex Med Rev*. 2022 Apr;10(2):209-230. doi: 10.1016/j.sxmr.2021.03.002. Epub 2021 Jun 12. PMID: 34127429.

Clinical Guideline Revision / History Information

Original Date: 4/13/2018

Reviewed/Revised: 4/15/2019, 7/21/2020, 04/21/2021, 12/01/2021, 04/25/2022, 05/02/2023, 04/16/2024