

Optical Coherence Tomography (OCT)

Disclaimer

Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.

Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.

Summary

Optical Coherence Tomography, or "OCT", is a medical imaging test that uses light waves to capture live 3-dimensional images. It is similar in principle to ultrasound (which uses sound echoes, rather than light wave reflections), however OCT provides much higher spatial resolution. OCT has been used to image different structures of the body, including the eye, the heart, the gastrointestinal (GI) system, the breast, and the upper airway. In the eyes, OCT allows for imaging of the retina and optic nerve. This imaging is helpful in monitoring and treating retinal disorders or optic nerve disorders such as glaucoma. It does not require any contact with the target surfaces and does not produce any ionizing radiation. In some cases, OCT can be used with other instruments such as an endoscope in the GI system or as an intravascular device in the arteries of the heart. OCT is a relatively novel technology and is rapidly evolving in both technique and clinical utility. This guideline provides the clinical criteria and exclusions for the currently supported clinical applications of Optical Coherence Tomography.

Definitions

"Retinopathy" refers to diseases of the retina that may impair vision, and is most often due to diabetes or hypertension. Diabetic retinopathy can be "proliferative" or "nonproliferative", depending on the severity of the disease.

"Macular Degeneration" is a condition that affects the macula, which is the central portion of the retina responsible for fine detail vision. It can be further categorized as "wet" or "dry" depending on the

underlying process, and stratified by stage (early, intermediate, late). Macular degeneration is a leading cause of vision loss.

“Macular Edema” occurs when fluid builds up behind the macula of the eye, leading to swelling and distortion of central vision. Macular edema can occur in a number of diseases, including macular degeneration, diabetic retinopathy, and retinal vein occlusions.

“Optical Coherence Tomography (OCT)” is an imaging technique that uses the reflections of light particles to create live 3-D images. Because OCT is based on light, it has significantly higher spatial resolution than comparable imaging techniques such as ultrasound and MRI. However, given the poor penetration of light into tissue, OCT is limited to relatively superficial surfaces. OCT does not generate ionizing radiation. OCT can be used to evaluate the retinal, optic nerve, and anterior chamber.

- “Spectral Domain OCT” is a newer version of OCT technology which can obtain images up to 50 times faster than traditional “time domain” and may be more appropriate in the diagnosis and assessment of certain conditions.
- “OCT Angiography” or “Intravascular OCT” refers to OCT imaging conducted with specialized, miniature devices from within a blood vessel.

“Posterior Segment” refers to the back two-thirds of the eye, and includes the vitreous humor, retina, choroid, macula, and optic nerve

“Anterior Segment” refers to the anterior one-third of the eye, and includes the cornea, iris, ciliary body, and lens.

“Glaucoma Suspect” refers to an individual with clinical findings and/or risk factors that indicate an elevated risk of developing primary open angle glaucoma. Risk factors include age older than 50 years, family history of glaucoma, and black race. Clinical findings include optic nerve or nerve fiber layer defect suggestive of glaucoma, visual field abnormality consistent with glaucoma, and/or elevated intraocular pressure (IOP) >21 mm Hg.

Clinical Indications

The Plan considers posterior Optical Coherence Tomography medically necessary when any ONE of the following criteria is met:

1. Spectral Domain OCT is indicated when a member is taking chloroquine, hydroxychloroquine, ezogabine, or vigabatrin and ONE of the following criteria are met:
 - a. Baseline exam within the first year of medication use; *or*
 - b. As a once yearly exam for patients with 1 or more of the following:
 - i. 5 years or more of use; *or*
 - ii. Documentation of elevated risk for developing retinopathy, defined by:
 1. Concurrent macular disease; *or*
 2. Concurrent renal disease; *or*

3. Concomitant use of tamoxifen; *or*
 4. High-dose chloroquine (>2.3mg/kg) or hydroxychloroquine (>5mg/kg).
2. Macular edema when at least ONE of the following criteria is met:
 - a. Needed to establish the diagnosis of macular edema; *or*
 - b. Cystoid macular edema; *or*
 - c. When the results may impact the treatment plan (e.g., the need for antiangiogenic treatment).
 3. Macular degeneration when at least ONE of the following criteria is met:
 - a. Neovascular age-related macular degeneration and OCT needed to establish the diagnosis when fluorescein angiography is contraindicated or unavailable; *or*
 - b. Cystoid macular degeneration; *or*
 - c. When the results may impact the treatment plan (e.g. the need for antiangiogenic treatment).
 4. OCT may be indicated to document the appearance of posterior segment structures in members who have a diagnosis of at least ONE of the following:
 - a. Age-related macular degeneration; *or*
 - b. Benign neoplasms of the retina and choroid; *or*
 - c. Central Serous Retinopathy (CSR); *or*
 - d. Diabetic retinopathy; *or*
 - e. Glaucoma or glaucoma suspect, no more than once per year; *or*
 - f. Inherited retinal dystrophy (e.g., RPE65 gene mutations); *or*
 - g. Macular hole; *or*
 - h. Macular edema; *or*
 - i. Malignant neoplasms of the retina and choroid; *or*
 - j. Retinal vein occlusion; *or*
 - k. Other retinal disorders and retinopathy; *or*
 - l. Panuveitis; *or*
 - m. Papilledema associated with increased intracranial pressure; *or*
 - n. Pseudotumor cerebri; *or*
 - o. Posterior vitreous detachment (vitreous degeneration); *or*
 - p. Vitreomacular adhesion or vitreomacular traction; *or*
 - q. Vogt-Koyanagi-Haradas.

Experimental or Investigational / Not Medically Necessary

The Plan considers Optical Coherence Tomography experimental and investigational for the following indications, as the current evidence is insufficient to demonstrate clear clinical benefit:

- Gastrointestinal usage, including but not limited to assessment or diagnosis of:
 - Esophageal mucosal diseases (e.g. Barrett's esophagus or squamous cell carcinoma)
 - Gastric mucosa
 - Diseases of the colon and small bowel (e.g. inflammatory bowel disorders, polyps)

- Biliary and pancreatic duct measurements
- Upper airway OCT for obstructive sleep apnea
- Any intraoperative OCT, including OCT for the purpose of lymph node or tumor margin assessment
- Ocular indications other than those defined above, including but not limited to:
 - Anterior segment OCT (AS-OCT) imaging (e.g., cornea, iris, ciliary body, and lens) is considered experimental/investigational as evidence is still insufficient for anterior OCT to be used as a substitute or as a stand-alone method for guiding treatment of anterior segment conditions. Overall, there is a lack of recommendations from society practice guidelines and peer-reviewed literature strongly supporting AS-OCT, and it has not been shown to improve net health outcomes
 - *Gonioscopy is the gold-standard for evaluating the anterior segment of the eye, per the American Academy of Ophthalmology (AAO).*
 - Upon review of recent literature, guidelines and Hayes, the role of anterior segment OCT is still evolving in the utility of AS-OCT for a variety of conditions or with different methodologies such as swept source anterior segment optical coherence tomography (SS-ASOCT). Although there may be some literature and cross-sectional studies on clinical validity for AS-OCT indications, overall, there is a lack of peer-reviewed literature supporting clinical utility with AS-OCT guided treatment in RCT trials or comparative trials with AS-OCT and another technology, placebo, or sham (Hayes, Inc., 2023; Fernández-Vigo et al., 2022; Marin et al., 2022; Preetam Peraka et al., 2023; Pujari et al., 2021; Triolo et al., 2021)
 - For the 2023 Glaucoma Summary Benchmarks by AAO, optical coherence tomography is not included in the preferred practice pattern guidelines for evaluation, diagnosis, or management of primary open-angle glaucoma or primary angle closure.
 - The American Academy of Ophthalmology (AAO) Preferred Practice Patterns for Primary Angle Closure (2015) state that AS-OCT is “limited to evaluating the iridocorneal angle” and that it “may prove useful in evaluating secondary causes of angle closure”. Gonioscopy is discussed as the gold standard to be performed in all patients with suspected angle closure. There have been no large, prospective, randomized clinical trials looking at the clinical effectiveness of AS-OCT for primary angle closure.
 - For the 2023 Cornea/External Disease Summary Benchmarks, anterior segment optical coherence tomography is listed in the preferred practice pattern guidelines as an applicable diagnostic test, but not an explicit recommendation for Corneal Edema and Opacification (Initial Evaluation).
 - For the 2023 Retina Summary Benchmarks, the preferred practice pattern guidelines mention for macular hole post-operative follow-up visit that an OCT

evaluation to document post-op macular anatomy may be indicated, but not an explicit recommendation for the anterior chamber and central retina.

- A 2022 systematic review by Marin and colleagues assessed anterior segment-OCT and image analysis methods as this field is less developed compared to posterior segment-OCT; however, comparisons for AS-OCT were difficult due to limited number of images from studies and additional research is needed.
 - For lens-to-cornea fit of rigid gas-permeable (RGP) lenses: Piotrowiak et al (2014) showed AS-OCT was inferior to fluorescein pattern assessment, with lower sensitivity for apical clearance detection.
 - Anterior chamber angle (ACA) measurement - Maram et al (2015) looked at 20 eyes and found low reproducibility among experienced clinicians. Further literature is limited for this indication.
 - Other potential indications for AS-OCT, including but not limited to intraoperative OCT, graft versus host disease (GVHD), anterior segment vascular imaging, and assessment of Haab striae, have inadequate clinical evidence in the currently available literature.
- Routine *screening*, including but not limited to the following:
 - Glaucoma, pre-glaucoma, or ocular hypertension
 - *Rationale:* The USPSTF (2022) concludes that for asymptomatic adults 40 years or older, the current evidence is insufficient to assess the balance of benefits and harms of screening (e.g., OCT, visual field assessment) for primary open-angle glaucoma in adults.
Bussel et al (2014) summarized the findings of 7 studies on glaucoma screening and monitoring of progression using OCT. They found that “in summary, OCT currently lacks the necessary diagnostic performance for general population glaucoma screening.” While there is some evidence of the ability of OCT to differentiate normal and glaucomatous eyes, the current clinical evidence has not been fully validated.
 - Cataracts
 - *Rationale:* OCT is not used for the diagnosis or screening of cataracts. OCT has been used in the pre-operative planning or for monitoring of post-operative complications following cataract surgery; however, the clinical evidence is limited for these indications. Furthermore, the presence of cataracts may impact OCT image quality and retinal thickness measurements (Van Velthoven 2006).
 - Corneal conditions, including but not limited to keratitis, Thygeson's disease
 - Keratoconjunctivitis sicca (i.e., dry eyes)
 - *Rationale:* A single center, prospective study by Ibrahim et al (2010) looked at OCT for diagnosing keratoconjunctivitis sicca in 24 patients and 27 control subjects. Sensitivity and specificity were 67% and 81%, respectively. Further research is needed to identify the clinical outcomes using OCT for this indication.

- Posterior capsule opacification
- Neurodegenerative disorders affecting the optic nerve (e.g., multiple sclerosis and optic neuritis)
 - *Rationale:* While the role for optic nerve measurements using OCT has been outlined above, routine screening using OCT for neurodegenerative disorders that may affect the optic nerve is not indicated. OCT has not been adequately studied for this purpose.
- Papilledema, Unexplained vision loss, (not caused by diabetic retinopathy or pseudotumor cerebri)
 - *Rationale:* Extensive literature review by the AAO states that there is not currently enough randomized evidence to use OCT for routine evaluation of unexplained vision loss, in routine screening for diabetic retinopathy, or for “other causes” of macular swelling. OCT is not mentioned as indicated or not indicated for other disease processes in the AAO guidelines.
- Identification of fungal endophthalmitis after cataract surgery
 - *Rationale:* The evidence for the use of AS-OCT in the identification of fungal infections after cataract surgery is limited to case reports (Kitahata 2016) and has not been validated in a randomized, prospective clinical trial.
- Imaging of extra- or intra-ocular musculature
 - *Rationale:* Several studies (Pihlblad 2016, Ngo 2015, Park 2014) have looked at AS-OCT for imaging of the ocular musculature. While the results on the ability to accurately and reproducibly measure the muscle insertion distances for pre-operative planning have been promising, the current evidence has not been validated in clinical studies nor has it demonstrated any improved clinical outcomes.
- Any other procedure or indication not meeting the above medical necessity criteria

The Plan considers “OCT Angiography” or “Intravascular OCT” experimental or investigational, as the current evidence is insufficient to demonstrate clear clinical benefit. This includes, but is not limited to:

- Diagnosis of spontaneous coronary artery dissection (SCAD)
- Diagnosis or assessment of coronary artery plaques
- Treatment of coronary disease (as an adjunct to percutaneous coronary intervention (PCI))
- Assessment or guidance of coronary artery stent placement (including evaluation of arterial bifurcations)
- Assessment of coronary artery stent failure (malposition)
- Identification of angiographically unclear lesions
- Assessment of acute coronary syndromes
- Diagnosis or assessment of intracranial aneurysms, ruptured or intact
- Assessment of carotid artery stenosis and/or stroke risk
- Assessment of pulmonary arterial wall fibrosis

Clinical Evidence on Intravascular OCT

- A systematic review of 15 studies was published in 2015 by D'Ascenzo et al. to evaluate the accuracy of intravascular OCT and intravascular ultrasound (IVUS) in identifying functional coronary stenosis. The group found that both modalities had only a moderate diagnostic accuracy for hemodynamically significant lesions. The authors concluded that both the sensitivity and specificity were inadequate to guide revascularization. (*D'Ascenzo F, Barbero U, Cerrato E, et al. Accuracy of intravascular ultrasound and optical coherence tomography in identifying functionally significant coronary stenosis according to vessel diameter: a meta-analysis of 2,581 patients and 2,807 lesions. Am Heart J. 2015; 169(5):663-673*)
- The Society of Cardiovascular Angiography and Interventions released a consensus statement in 2014 evaluating IVUS and intravascular OCT, concluding that "the appropriate role for optical coherence tomography in routine clinical decision making has not been established". (*Lotfi A, Jeremias A, Fearon WF, et al. Society of Cardiovascular Angiography and Interventions. Expert consensus statement on the use of fractional flow reserve, intravascular ultrasound, and optical coherence tomography: a consensus statement of the Society of Cardiovascular Angiography and Interventions. Catheter Cardiovasc Interv. 2014; 83(4):509-418*)
- The ILUMIEN IV study (NCT NCT03507777) is underway for a prospective, single-blind clinical investigation randomizing subjects to OCT-guided coronary stent implantation vs. angiography-guided coronary stent implantation in a 1:1 ratio. The clinical investigation will be conducted at approximately 125 centers in North America (US and Canada), Europe, Middle East and Asia-Pacific. After hospital discharge, all patients will have clinical follow-up at 30 days, 1 year, and 2 years. As of April, 2024, the publication for this ILUMIEN IV study has not been completed. The ILUMIEN III: OPTIMIZE PCI trial was performed to compare IVUS, OCT, and coronary angiography (CA) in guiding coronary stent placement. The randomized study demonstrated that IVUS and OCT were non-inferior, however CA was superior to both modalities. (*Ali ZA, Maehara A, Généreux P, et al. Optical coherence tomography compared with intravascular ultrasound and angiography to guide coronary stent implantation (ILUMIEN III: OPTIMIZE PCI): a randomised controlled trial. Lancet 2016; 388:2618.*)
- For coronary plaque characterization and stent implantation for coronary artery revascularization, the FDA has 510(k) clearance for Tigereye Cto-Crossing Catheter, Pantheris System, Optis Mobile Next Imaging System, Optis Integrated Next Imaging System, Otis 2.1 Optical Coherence Tomography System, Thia Optical Coherence Tomography System, Apollovue S100 Image System. The 2021 ACC/AHA/SCA Guideline states that, "In patients undergoing coronary stent implantation, OCT is a reasonable alternative to IVUS for procedural guidance, except in ostial left main disease"(2a) and "In patients with stent failure, IVUS or OCT is reasonable to determine the mechanism of stent failure. (2a)" However, because OCT requires blood clearance, its effectiveness for imaging ostial left main disease is limited. The results of the ILUMIEN IV study will be pivotal for practice guidance.

Applicable Billing Codes (HCPCS & CPT Codes)

CPT/HCPCS Codes considered medically necessary if criteria are met:	
<i>Code</i>	<i>Description</i>
92133	Scanning computerized ophthalmic diagnostic imaging, posterior segment, with interpretation and report, unilateral or bilateral; optic nerve
92134	Scanning computerized ophthalmic diagnostic imaging, posterior segment, with interpretation and report, unilateral or bilateral; retina
ICD-10 codes considered medically necessary if criteria are met:	
C69.20 - C69.32	Malignant neoplasm of the retina and choroid
D18.09	Malignant neoplasm of the retina and choroid
D31.20 - D31.32	Benign neoplasm of the retina and choroid
E08.311 - E08.3599	Diabetes mellitus due to underlying condition with ophthalmic complications
E09.311 - E09.3599	Drug or chemical induced diabetes mellitus with ophthalmic complications
E10.311 - E10.3599	Type 1 diabetes mellitus with ophthalmic complications
E11.311 - E11.3599	Type 2 diabetes mellitus with ophthalmic complications
E13.311 - E13.3599	Other specified diabetes mellitus with ophthalmic complications
G40.201 - G40.219	Localization-related (focal)(partial) symptomatic epilepsy and epileptic syndromes with complex partial seizures, not intractable and intractable [screening for vigabatrin (Sabril) toxicity]
G40.401 - G40.419	Other generalized epilepsy and epileptic syndromes, not intractable and intractable, with and without status epilepticus [screening for vigabatrin (Sabril) toxicity]
G40.821 - G40.824	Epileptic spasms [screening for vigabatrin (Sabril) toxicity]
G93.2	Benign intracranial hypertension
H20.821 - H20.823	Vogt-Koyanagi syndrome
H34.8111 - H34.9	Retinal vascular occlusions

H35.00 - H35.23	Other retinal disorders and retinopathy
H35.30	Unspecified macular degeneration Age-related macular degeneration
H35.3110 - H35.3194	Nonexudative age-related macular degeneration
H35.3210 - H35.3293	Exudative age-related macular degeneration
H35.341 - H35.343	Macular cyst, hole or pseudohole
H35.351 - H35.359	Cystoid macular degeneration
H35.711	Central serous chorioretinopathy, right eye
H35.712	Central serous chorioretinopathy, left eye
H35.713	Central serous chorioretinopathy, bilateral
H35.81	Retinal edema
H40.001 - H40.9	Glaucoma
H43.811 - H43.813	Vitreous degeneration
H43.821 - H43.823	Vitreomacular adhesion or traction
H44.11 - H44.113	Panuveitis
H47.11	Papilledema associated with increased intracranial pressure
H59.031- H59.039	Cystoid macular edema following cataract surgery
Q15.0	Congenital glaucoma
T37.2X1A - T37.2X4S	Poisoning by antimalarials and drugs acting on other blood protozoa
Z79.899	Other long term (current) drug therapy
ICD-10 codes <i>NOT</i> considered medically necessary, or experimental or investigational:	
C69.10 - C69.12	Malignant neoplasm of cornea
C69.40 - C69.42	Malignant neoplasm of ciliary body
D31.10 - D31.12	Benign neoplasm of cornea

D31.40 - D31.42	Benign neoplasm of ciliary body
D89.810 - D89.813	Graft-versus-host disease
G35	Multiple sclerosis
G47.33	Obstructive sleep apnea
H04.121 - H04.129	Dry eye syndrome
H16.001 - H22	Disorders of the cornea, iris and ciliary body
H25.011 - H25.9	Age-related cataract
H26.001 - H26.9	Other cataract
H27.00 - H27.9	Other disorders of lens
H28	Cataract in diseases classified elsewhere
H46.00 - H46.9	Optic neuritis
H47.10	Unspecified papilledema
H47.12	Papilledema associated with decreased ocular pressure
H47.13	Papilledema associated with retinal disorder
H49.00 - H52.7	Disorders of ocular muscles, binocular movement, accommodation and refraction
H53.121 - H53.129	Transient visual loss
H53.131 - H53.139	Sudden visual loss
K20.0 - K31.A29	Diseases of esophagus, stomach and duodenum
K50.00 - K52.9	Noninfective enteritis and colitis
K55.011 - K64.9	Other diseases of intestines
M35.00 - M35.09	Sjogren syndrome

	Sicca syndrome
Q12.0 - Q12.9	Congenital lens malformations
Z13.5	Encounter for screening for eye and ear disorders
Z46.0	Encounter for fitting and adjustment of spectacles and contact lenses

CPT/HCPCS considered experimental or investigational:	
<i>Code</i>	<i>Description</i>
92132	Scanning computerized ophthalmic diagnostic imaging, anterior segment, with interpretation and report, unilateral or bilateral
92978	Endoluminal imaging of coronary vessel or graft using intravascular ultrasound (IVUS) or optical coherence tomography during diagnostic evaluation and/or therapeutic intervention including imaging supervision, interpretation and report; initial vessel
92979	Endoluminal imaging of coronary vessel or graft using intravascular ultrasound (IVUS) or optical coherence tomography during diagnostic evaluation and/or therapeutic intervention including imaging supervision, interpretation and report; each additional vessel
0351T	Optical coherence tomography of breast or axillary lymph node, excised tissue, each specimen; real time intraoperative
0352T	Optical coherence tomography of breast or axillary lymph node, excised tissue, each specimen; interpretation and report, real-time or referred
0353T	Optical coherence tomography of breast, surgical cavity; real-time intraoperative
0354T	Optical coherence tomography of breast, surgical cavity; interpretation and report, real-time or referred

References

1. Ali ZA, Maehara A, Généreux P, et al. Optical coherence tomography compared with intravascular ultrasound and with angiography to guide coronary stent implantation (ILUMIEN III: OPTIMIZE PCI): a randomised controlled trial. *Lancet* 2016; 388:2618.
2. Allam RS, AbdElmohsen MN, Khafagy MM, Raafat KA, Sheta SM. Spectral domain optical coherence tomography of preclinical chloroquine maculopathy in Egyptian rheumatoid arthritis patients. *Journal of Ophthalmology* 2015;2015:292357

3. American Academy of Ophthalmology. (Dec 2023). Cornea/External Disease Summary Benchmarks for Preferred Practice Pattern® Guidelines. Retrieved from <https://www.aao.org/education/summary-benchmark-detail/cornea-external-disease-summary-benchmarks-2020>
4. American Academy of Ophthalmology. (Dec 2023). Glaucoma Summary Benchmarks for Preferred Practice Pattern® Guidelines. Retrieved from <https://www.aao.org/education/summary-benchmark-detail/glaucoma-summary-benchmarks-2020>
5. American Academy of Ophthalmology. (Dec 2023). Retina Summary Benchmarks for Preferred Practice Pattern® Guidelines. Retrieved from <https://www.aao.org/education/summary-benchmark-detail/retina-summary-benchmarks-2020>
6. Arf, S., Sayman Muslubas, I., Hocaoglu, M., Ersoz, M. G., Ozdemir, H., & Karacorlu, M. (2020). Spectral domain optical coherence tomography classification of diabetic macular edema: a new proposal to clinical practice. *Graefe's archive for clinical and experimental ophthalmology*, 258, 1165-1172.
7. Armstrong JJ, Leigh MS, Sampson DD, et al. Quantitative upper airway imaging with anatomic optical coherence tomography. *Am J Respir Crit Care Med* 2006; 173:226.
8. Arroyo JG. Age-related macular degeneration: Clinical presentation, etiology, and diagnosis. In: *UpToDate*, Trobe J (Ed), *UpToDate*, Waltham, MA (Accessed March 13, 2017)
9. Arvanitakis M, Hookey L, Tessier G, et al. Intraductal optical coherence tomography during endoscopic retrograde cholangiopancreatography for investigation of biliary strictures. *Endoscopy* 2009; 41:696.
10. Baskin DE. Optical coherence tomography in diabetic macular edema. *Current Opinion in Ophthalmology* 2010;21(3):1727.
11. Bezerra H. Intravascular OCT in PCI. *JACC*. June 13, 2016.
12. Boppart SA, Herrmann J, Pitris C, et al. High-resolution optical coherence tomography-guided laser ablation of surgical tissue. *J Surg Res* 1999; 82:275.
13. Boyd, K. (March 27, 2020). Eye Cancer. Retrieved 6/10/2024 from <https://www.aao.org/eye-health/diseases/eye-cancer#:~:text=The%20most%20common%20cancer%20that, get%20it%20in%20your%20eye.>
14. Boyd, K. (April 23m 2023). CME (Cystoid Macular Edema). Retrieved 6/10/20204 from <https://www.aao.org/eye-health/diseases/cme-cystoid-macular-edema>
15. Chou, R., Selph, S., Blazina, I., Bougatsos, C., Jungbauer, R., Fu, R., ... & Tehrani, S. (2022). Screening for glaucoma in adults: updated evidence report and systematic review for the US Preventive Services Task Force. *Jama*, 327(20), 1998-2012.
16. Chou, R., Selph, S., Blazina, I., Bougatsos, C., Jungbauer, R., Fu, R., ... & Tehrani, S. (2022). Screening for Glaucoma in Adults: A Systematic Review for the US Preventive Services Task Force.
17. Clayton LM, Devile M, Punte T, et al. Retinal nerve fiber layer thickness in vigabatrin- exposed patients. *Ann Neurol* 2011;69:845-854.

18. ClinicalTrials.gov. (last update posted June 9, 2023) ILUMIEN IV: OPTIMAL PCI. Retrieved from <https://clinicaltrials.gov/study/NCT03507777?tab=results> Accessed 5/21/24.
19. Costello F, Coupland S, Hodge W, et al. Quantifying axonal loss after optic neuritis with optical coherence tomography. *Ann Neurol* 2006; 59:963.
20. Consugar, M. B., Navarro-Gomez, D., Place, E. M., Bujakowska, K. M., Sousa, M. E., Fonseca-Kelly, Z. D.,... & Pierce, E. A. (2015). Panel-based genetic diagnostic testing for inherited eye diseases is highly accurate and reproducible, and more sensitive for variant detection, than exome sequencing. *Genetics in Medicine*, 17(4), 253-261.
21. Covert DJ. Retinal vein occlusion: Epidemiology, clinical manifestations, and diagnosis. In: UpToDate, Trobe J(Ed), UpToDate, Waltham, MA (Accessed March 13, 2017).
22. Culp, M. B., Benard, V., Dowling, N. F., King, J., Lu, H., Rao, C., ... & Wu, M. (2021). Ocular melanoma incidence rates and trends in the United States, 2001–2016. *Eye*, 35(2), 687-689.
23. Dada T, Sihota R, Gadia R, et al. Comparison of anterior segment optical coherence tomography and ultrasound biomicroscopy for assessment of the anterior segment. *J Cataract Refract Surg*. 2007; 33(5): 837-840.
24. de Seze J, Blanc F, Jeanjean L, et al. Optical coherence tomography in neuromyelitis optica. *Arch Neurol* 2008; 65:920.
25. Douglas PS. Spontaneous coronary artery dissection. In: UpToDate, Kaski JC (Ed), UpToDate, Waltham, MA (Accessed March 13, 2017).
26. Duker, J.S., Kaiser, P.K., Binder, S., et al. (2013). The International Vitreomacular Traction Study Group classification of vitreomacular adhesion, traction, and macular hole. *Ophthalmology*, 120(12), 2611-2619. doi: 10.1016/j.ophtha.2013.07.042. Epub 2013 Sep 17.
27. Duncan, J. L., Bernstein, P. S., Birch, D. G., Fishman, G. A., Heon, E., & Jacobson, S. G. (2016). Recommendations on clinical assessment of patients with inherited retinal degenerations-2016. American Academy of Ophthalmology Clinical Education/Guidelines/Clinical Statements. <https://www.aao.org/clinical-statement/recommendations-on-clinical-assessment-of-patients>
28. Evans JA, Bouma BE, Bressner J, et al. Identifying intestinal metaplasia at the squamocolumnar junction by using optical coherence tomography. *Gastrointest Endosc* 2007; 65:50.
29. Evans JA, Poneris JM, Bouma BE, et al. Optical coherence tomography to identify intramucosal carcinoma and high-grade dysplasia in Barrett's esophagus. *Clin Gastroenterol Hepatol* 2006; 4:38.
30. Fabian, I. D., Abdallah, E., Abdullahi, S. U., Abdulqader, R. A., Boubacar, S. A., Ademola-Popoola, D. S., ... & Gole, G. A. (2020). Global retinoblastoma presentation and analysis by national income level. *JAMA oncology*, 6(5), 685-695.
31. Familiari L, Strangio G, Consolo P, et al. Optical coherence tomography evaluation of ulcerative colitis: the patterns and the comparison with histology. *Am J Gastroenterol* 2006; 101:2833.
32. Fernández-Vigo, J. I., Shi, H., Burgos-Blasco, B., Fernández-Aragón, S., De-Pablo-Gómez-de-Liaño, L., Kudsieh, B., ... & Ángel Fernández-Vigo, J. (2022). Anterior scleral thickness dimensions by swept-source optical coherence tomography. *Clinical and Experimental Optometry*, 105(1), 13-19.

33. Fisher JB, Jacobs DA, Markowitz CE, et al. Relation of visual function to retinal nerve fiber layer thickness in multiple sclerosis. *Ophthalmology* 2006; 113:324.
34. FDA Drug Safety Communications. Potiga (Ezogabine): Drug Safety Communication - Linked To Retinal Abnormalities And Blue Skin Discoloration. November 1, 2013.
35. Friedland S. Optical coherence tomography in the gastrointestinal tract. In: UpToDate, Saltzman JR (Ed), UpToDate, Waltham, MA (Accessed March 13, 2017).
36. Fujino Y et al. Frequency-domain optical coherence tomography assessment of unprotected left main coronary artery disease-a comparison with intravascular ultrasound. *Catheter Cardiovasc Interv* 2013;82:E173-83
37. Hatta W, Uno K, Koike T, et al. A prospective comparative study of optical coherence tomography and EUS for tumor staging of superficial esophageal squamous cell carcinoma. *Gastrointest Endosc* 2012; 76:548.
38. Hatta W, Uno K, Koike T, et al. Optical coherence tomography for the staging of tumor infiltration in superficial esophageal squamous cell carcinoma. *Gastrointest Endosc* 2010; 71:899.
39. Hayes, Inc. Evidence Analysis Research Brief. *Optical Coherence Tomography Angiography with AngioPlex for Macular Degeneration*. Lansdale, PA: Hayes, Inc.; Sep 17, 2020.
40. Hayes, Inc., Evidence Analysis Research Brief. *Optical Coherence Tomography for Diagnosis and Management of Open Angle Glaucoma*. Lansdale, PA: Hayes, Inc.; Jun 22, 2023.
41. Hayes, Inc. Evidence Analysis Research Brief. *Optical Coherence Tomography for Plaque Characterization and Stent Implantation*. Lansdale, PA: Hayes, Inc.; Jan 11, 2023.
42. Huang, J., Belmadani, K., Chatot, M., Ecarnot, F., Chopard, R., Wang, M., ... & Meneveau, N. (2018). Clinical significance of optical coherence tomography-guided angioplasty on treatment selection. *Experimental and Therapeutic Medicine*, 16(2), 483-492.
43. Huang-Link, Y., Eleftheriou, A., Yang, G., Johansson, J. M., Apostolou, A., Link, H., & Jin, Y. P. (2019). Optical coherence tomography represents a sensitive and reliable tool for routine monitoring of idiopathic intracranial hypertension with and without papilledema. *European journal of neurology*, 26(5), 808-e57.
44. Hui, V. W., Szeto, S. K., Tang, F., Yang, D., Chen, H., Lai, T. Y., ... & Cheung, C. Y. (2022). Optical coherence tomography classification systems for diabetic macular edema and their associations with visual outcome and treatment responses—an updated review. *Asia-Pacific Journal of Ophthalmology*, 11(3), 247-257.
45. Hunter RS, Skondra D, Papaliadis G, Sobrin L. Role of OCT in the diagnosis and management of macular edema from uveitis. *Seminars in Ophthalmology* 2012;27(56):23641.
46. Idiopathic Macular Hole. Preferred Practice Pattern. (PPP) - Updated 2017. American Academy of Ophthalmology. Available at: <https://www.aao.org/Assets/647f6f55-24ba-46f3-86f3-91798ca05724/636277037392430000/idiopathic-macular-hole-updated-2017-pdf>
47. Inoue T, Shite J, Yoon J, et al. Optical coherence evaluation of everolimus-eluting stents 8 months after implantation. *Heart*. 2011; 97(17):1379-1384.

48. Jäckle S, Gladkova N, Feldchtein F, et al. In vivo endoscopic optical coherence tomography of esophagitis, Barrett's esophagus, and adenocarcinoma of the esophagus. *Endoscopy* 2000; 32:750.
49. Jager RD, Mieler WF, Miller JW. Age-related macular degeneration. *New England Journal of Medicine* 2008;358(24):2606-17.
50. Jang IK. Who should Get Optical Coherence Tomography? *JACC* Feb 2013.
51. Jang IK, Bouma BE, Kang DH et al. Visualization of coronary atherosclerotic plaques in patients using optical coherence tomography: comparison with intravascular ultrasound. *J Am Coll Cardiol.* 2002; 39(4):604-609.
52. Jang IK, Tearney GJ, MacNeill B, et al. In vivo characterization of coronary atherosclerotic plaque by use of optical coherence tomography. *Circulation.* 2005; 111(12):1551-1555.
53. Jia H, Abtahian F, Aguirre AD, et al. In vivo diagnosis of plaque erosion and calcified nodule in patients with acute coronary syndrome by intravascular optical coherence tomography. *J Am Coll Cardiol.* 2013; 62(19):1748-1758.
54. Jittpoonkuson T, Garcia PM, Rosen RB. Correlation between fluorescein angiography and spectral-domain optical coherence tomography in the diagnosis of cystoid macular edema. *British Journal of Ophthalmology* 2010;94(9):1197-200
55. John R, Adie SG, Chaney EJ, et al. Three-dimensional optical coherence tomography for optical biopsy of lymph nodes and assessment of metastatic disease. *Ann Surg Oncol.* 2013;20(11):3685-3693.
56. Johnson PM, Patel J, Yeung M, Kaul P. Intra-coronary imaging modalities. *Curr Treat Options Cardiovasc Med.* 2014; 16(5):304.
57. Kennedy BF, McLaughlin RA, Kennedy KM, et al. Investigation of optical coherence microelastography as a method to visualize cancers in human breast tissue. *Cancer Res.* 2015;75(16):3236-3245.
58. Kubo T, Nakamura N, Matsuo Y, et al. Virtual histology intravascular ultrasound compared with optical coherence tomography for identification of thin-cap fibroatheroma. *Int Heart J.* 2011; 52(3):175-179.
59. Kuo WC, Kim J, Shemonski ND, et al. Real-time three-dimensional optical coherence tomography image-guided core-needle biopsy system. *Biomed Opt Express.* 2012;3(6):1149-1161.
60. Lawton, J. S., Tamis-Holland, J. E., Bangalore, S., Bates, E. R., Beckie, T. M., Bischoff, J. M., ... & Zwischenberger, B. A. (2022). 2021 ACC/AHA/SCAI guideline for coronary artery revascularization: executive summary: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*, 145(3), e4-e17.
61. Levine GN et al., 2011 ACCF/AHA/ SCAI Guideline for Percutaneous Coronary Intervention. A report of the American College of Cardiology Foundation/ American Heart Association Task Force on Practice Guidelines and the Society for Cardiovascular Angiography and Interventions, *J Am Coll Cardiol*, 2011;58:e44–122.

62. Li EY, Mohamed S, Leung CK, et al. Agreement among 3 methods to measure corneal thickness: ultrasound pachymetry, Orbscan II, and Visante anterior segment optical coherence tomography. *Ophthalmology*. 2007; 114(10):1842-1847.
63. Liu XY, Peng XY, Wang S, et al. Features of optical coherence tomography for the diagnosis of Vogt- Koyanagi-Harada disease. *Retina*. 2016;36(11):2116- 2123.
64. LopezLopez F, GomezUlla F. Update on the imaging techniques in the diagnosis of diabetic retinopathy. *Current Diabetes Reviews* 2012;8(3):2008.
65. Lotfi A, Jeremias A, Fearon WF, et al. Society of Cardiovascular Angiography and Interventions. Expert consensus statement on the use of fractional flow reserve, intravascular ultrasound, and optical coherence tomography: a consensus statement of the Society of Cardiovascular Angiography and Interventions. *Catheter Cardiovasc Interv*. 2014; 83(4):509-418.
66. Marin, Y. F. G., Alonso-Caneiro, D., Vincent, S. J., & Collins, M. J. (2022). Anterior segment optical coherence tomography (AS-OCT) image analysis methods and applications: A systematic review. *Computers in Biology and Medicine*, 146, 105471.
67. Marmor MF, Kellner U, Lai TY, Melles RB, Mieler WF, American Academy of Ophthalmology. Recommendations on screening for chloroquine and hydroxychloroquine retinopathy (2016 Revision). *Ophthalmology* 2016;123(6).
68. McLaughlin, R. A., Noble, P. B., & Sampson, D. D. (2014). Optical coherence tomography in respiratory science and medicine: from airways to alveoli. *Physiology (Bethesda, Md.)*, 29(5), 369–380. <https://doi.org/10.1152/physiol.00002.2014>
69. Meneveau, N., Souteyrand, G., Motreff, P., Caussin, C., Amabile, N., Ohlmann, P., ... & Schiele, F. (2016). Optical coherence tomography to optimize results of percutaneous coronary intervention in patients with non–ST-elevation acute coronary syndrome: results of the multicenter, randomized DOCTORS study (Does Optical Coherence Tomography Optimize Results of Stenting). *Circulation*, 134(13), 906-917.
70. Michelessi M, Lucenteforte E, Oddone, et al. Optic nerve head and fibre layer imaging for diagnosing glaucoma. *Cochrane Database Syst Rev*. 2015;11:CD008803.
71. Mintz GS, Nissen SE, Anderson WD, et al. American College of Cardiology Clinical Expert Consensus Document on Standards for Acquisition, Measurement and Reporting of Intravascular Ultrasound Studies (IVUS). A report of the American College of Cardiology Task Force on Clinical Expert Consensus Documents. *J Am Coll Cardiol* 2001; 37:1478.
72. Murakami T, Yoshimura N. Structural changes in individual retinal layers in diabetic macular edema. *Journal of Diabetes Research* 2013;2013:920713.
73. Naismith RT, Tutlam NT, Xu J, et al. Optical coherence tomography is less sensitive than visual evoked potentials in optic neuritis. *Neurology* 2009; 73:46.
74. Naismith RT, Tutlam NT, Xu J, et al. Optical coherence tomography differs in neuromyelitis optica compared with multiple sclerosis. *Neurology* 2009; 72:1077.
75. National Comprehensive Cancer Network. (May 23, 2024). NCCN Clinical Practice Guidelines in Oncology. Melanoma: Uveal. Version 1.2024. Retrieved from https://www.nccn.org/professionals/physician_gls/pdf/uveal.pdf

76. Papaliodis, GN. (Oct 2022). Uveitis: Etiology, clinical manifestations, and diagnosis. UpToDate, Waltham, MA. Last reviewed: May 2024. Retrieved from https://www.uptodate.com/contents/uveitis-etiology-clinical-manifestations-and-diagnosis?search=panuveitis&source=search_result&selectedTitle=1%7E21&usage_type=default&display_rank=1#H3185467976
77. Patel R, Khan A, Quinlan R, Yaroslavsky AN. Polarization-sensitive multimodal imaging for detecting breast cancer. *Cancer Res.* 2014;74(17):4685-4693
78. Pasarikovski, C. R., Ku, J. C., Priola, S. M., da Costa, L., & Yang, V. X. D. (2020). Endovascular optical coherence tomography imaging in cerebrovascular disease. *Journal of clinical neuroscience : official journal of the Neurosurgical Society of Australasia*, 80, 30–37. <https://doi.org/10.1016/j.jocn.2020.07.064>
79. Pavlin CJ, Vásquez LM, Lee R, et al. Anterior segment optical coherence tomography and ultrasound biomicroscopy in the imaging of anterior segment tumors. *Am J Ophthalmol.* 2009; 147(2):214-219.
80. Pfau PR, Sivak MV Jr, Chak A, et al. Criteria for the diagnosis of dysplasia by endoscopic optical coherence tomography. *Gastrointest Endosc* 2003; 58:196.
81. Poneros J. Optical coherence tomography and the detection of dysplasia in Barrett's esophagus. *Gastrointest Endosc* 2005; 62:832.
82. Poneros JM, Brand S, Bouma BE, et al. Diagnosis of specialized intestinal metaplasia by optical coherence tomography. *Gastroenterology* 2001; 120:7.
83. Poneros JM, Tearney GJ, Shiskov M, et al. Optical coherence tomography of the biliary tree during ERCP. *Gastrointest Endosc* 2002; 55:84.
84. Prati F, Guagliumi G, Mintz GS, et al. Expert review document part 2: methodology, terminology and clinical applications of optical coherence tomography for the assessment of interventional procedures. *Eur Heart J* 2012; 33:2513.
85. Prati F, Regar E, Mintz GS, et al. Expert review document on methodology, terminology, and clinical applications of optical coherence tomography: physical principles, methodology of image acquisition, and clinical application for assessment of coronary arteries and atherosclerosis. *Eur Heart J* 2010;31:401-415.
86. Preetam Peraka, R., & Murthy, S. I. (2023, April). Role of anterior segment optical coherence tomography in scleral diseases: A review. In *Seminars in Ophthalmology* (Vol. 38, No. 3, pp. 238-247). Taylor & Francis.
87. Primary Angle Closure. Preferred Practice Pattern. (PPP). American Academy of Ophthalmology. 2015. Available at: <https://www.aao.org/Assets/0cec3943-ea25-4775-ac1dd8414570dcaa/636125680057700000/primary-angle-closure-ppp-pdf>
88. Primary Open-Angle Glaucoma. Preferred Practice Pattern. (PPP) - 2015. American Academy of Ophthalmology. Available at: <https://www.aao.org/Assets/964d5b3f-a551-46da-bf47-ecacd9ce909c/636125693862470000/primary-open-angle-glaucoma-ppp-pdf>

89. Primary Open-Angle Glaucoma Suspect. Preferred Practice Pattern (PPP) - 2015. American Academy of Ophthalmology. Available at:
<https://www.aao.org/Assets/82113fbf-efea-4afd-a6eb-3646686613f7/636125690676100000/primary-open-angle-glaucoma-suspect-ppp-pdf>
90. Pujari, A., Agarwal, D., & Sharma, N. (2021, November). Clinical role of swept source optical coherence tomography in anterior segment diseases: a review. In *Seminars in Ophthalmology* (Vol. 36, No. 8, pp. 684-691). Taylor & Francis.
91. Qi X, Sivak MV, Isenberg G, et al. Computer-aided diagnosis of dysplasia in Barrett's esophagus using endoscopic optical coherence tomography. *J Biomed Opt* 2006; 11:044010.
92. Regar E, van Soest G, Bruining N, et al. Optical coherence tomography in patients with acute coronary syndrome. *EuroIntervention* 2010; 6 Suppl G:G154.
93. Regar E. Intravascular ultrasound, optical coherence tomography, and angiography of coronary circulation. In: *UpToDate*, Cutlip D (Ed), *UpToDate*, Waltham, MA (Accessed March 13, 2017)
94. Ribeiro, J., & Pires-Morais, G. (2023). IVUS versus OCT: Calibrating the use of intracoronary imaging. *Revista Portuguesa de Cardiologia*.
95. Savastru D, Chang EW, Miclos S, et al. Detection of breast surgical margins with optical coherence tomography imaging: A concept evaluation study. *J Biomed Opt*. 2014;19(5):056001
96. Seitz U, Freund J, Jaeckle S, et al. First in vivo optical coherence tomography in the human bile duct. *Endoscopy* 2001; 33:1018.
97. Shen B, Zuccaro G, Gramlich TL, et al. Ex vivo histology-correlated optical coherence tomography in the detection of transmural inflammation in Crohn's disease. *Clin Gastroenterol Hepatol* 2004; 2:754.
98. South FA, Chaney EJ, Marjanovic M, et al. Differentiation of ex vivo human breast tissue using polarization-sensitive optical coherence tomography. *Biomed Opt Express*. 2014;5(10):3417-3426.
99. Sun JG, Adie SG, Chaney EJ, Boppart SA. Segmentation and correlation of optical coherence tomography and X-ray images for breast cancer diagnostics. *J Innov Opt Health Sci*. 2013;6(2):1350015.
100. Tearney GJ, Regar E, Akasaka, T et al. Consensus standards for acquisition, measurement, and reporting of intravascular optical coherence tomography studies: a report from the International Working Group for Intravascular Optical Coherence Tomography Standardization and Validation. *J Am Coll Cardiol*. 2012; 59(12):1058-1072.
101. Testoni PA, Mangiavillano B, Albarello L, et al. Optical coherence tomography to detect epithelial lesions of the main pancreatic duct: an Ex Vivo study. *Am J Gastroenterol* 2005; 100:2777.
102. Triolo, G., Barboni, P., Savini, G., De Gaetano, F., Monaco, G., David, A., & Scialdone, A. (2021). The use of anterior-segment optical-coherence tomography for the assessment of the iridocorneal angle and its alterations: update and current evidence. *Journal of Clinical Medicine*, 10(2), 231.
103. Trip SA, Schlottmann PG, Jones SJ, et al. Retinal nerve fiber layer axonal loss and visual dysfunction in optic neuritis. *Ann Neurol* 2005; 58:383.

104. Tsai, T. H., Leggett, C. L., Trindade, A. J., Sethi, A., Swager, A. F., Joshi, V., Bergman, J. J., Mashimo, H., Nishioka, N. S., & Namati, E. (2017). Optical coherence tomography in gastroenterology: a review and future outlook. *Journal of biomedical optics*, 22(12), 1–17. <https://doi.org/10.1117/1.JBO.22.12.121716>
105. Tubert, D. (June 2021). What Is Optical Coherence Tomography? AAO.org. <https://www.aao.org/eye-health/treatments/what-is-optical-coherence-tomography>
106. U.S. National Library of Medicine. ClinicalTrials.gov. ILUMIEN IV: OPTIMAL PCI. NCT03507777. Start date March 26, 2018, Completion Date Feb 28, 2023. Retrieved from <https://clinicaltrials.gov/ct2/show/NCT03507777>
107. Van de Werf F, Bax J, Betriu A, et al. Management of acute myocardial infarction in patients presenting with persistent ST-segment elevation: the Task Force on the Management of ST-Segment Elevation Acute Myocardial Infarction of the European Society of Cardiology. *Eur Heart J* 2008; 29:2909.
108. Virgili G, et al. Optical coherence tomography (OCT) for detection of macular oedema in patients with diabetic retinopathy. *Cochrane Database of Systematic Reviews* 2015, Issue 1.
109. Vijay, V., Mollan, S. P., Mitchell, J. L., Bilton, E., Alimajstorovic, Z., Markey, K. A., ... & Sinclair, A. J. (2020). Using optical coherence tomography as a surrogate of measurements of intracranial pressure in idiopathic intracranial hypertension. *JAMA ophthalmology*, 138(12), 1264-1271.
110. Wang, M. T., Bhatti, M. T., & Danesh-Meyer, H. V. (2022). Idiopathic intracranial hypertension: pathophysiology, diagnosis and management. *Journal of Clinical Neuroscience*, 95, 172-179.
111. Wielders, L. H., Schouten, J. S., Winkens, B., van den Biggelaar, F. J., Veldhuizen, C. A., Murta, J. C., ... & Sisquella, M. (2018). Randomized controlled European multicenter trial on the prevention of cystoid macular edema after cataract surgery in diabetics: ESCRS PREMED Study Report 2. *Journal of Cataract & Refractive Surgery*, 44(7), 836-847.
112. Wolf S, WolfSchnurrbusch U. Spectraldomain optical coherence tomography use in macular diseases: a review. *Ophthalmologica* 2010;224(6):33340.
113. Xie, J. S., Donaldson, L., & Margolin, E. (2022). Papilledema: a review of etiology, pathophysiology, diagnosis, and management. *Survey of Ophthalmology*, 67(4), 1135-1159.
114. Yeh EA, Marrie RA, Reginald YA, et al. Functional-structural correlations in the afferent visual pathway in pediatric demyelination. *Neurology* 2014; 83:2147.
115. Zuccaro G, Gladkova N, Vargo J, et al. Optical coherence tomography of the esophagus and proximal stomach in health and disease. *Am J Gastroenterol* 2001; 96:2633.
116. Kitahata M, Suzuki T, Oka N, Toriyama K, Ohashi Y. Anterior Segment Optical Coherence Tomography of Patients With Late-Onset Tunnel Fungal Infections With Endophthalmitis After Cataract Surgery. *Cornea*. 2016;35(8):1138-40.
117. Pihlblad MS, Erenler F, Sharma A, Manchandia A, Reynolds JD. Anterior Segment Optical Coherence Tomography of the Horizontal and Vertical Extraocular Muscles With Measurement of the Insertion to Limbus Distance. *J Pediatr Ophthalmol Strabismus*. 2016;53(3):141-5.
118. Ngo CS, Smith D, Kraft SP. The accuracy of anterior segment optical coherence tomography (AS-OCT) in localizing extraocular rectus muscles insertions. *J AAPOS*. 2015;19(3):233-6.

119. Park KA, Lee JY, Oh SY. Reproducibility of horizontal extraocular muscle insertion distance in anterior segment optical coherence tomography and the effect of head position. *J AAPOS*. 2014;18(1):15-20.
120. Marion KM, Maram J, Pan X, et al. Reproducibility and Agreement Between 2 Spectral Domain Optical Coherence Tomography Devices for Anterior Chamber Angle Measurements. *J Glaucoma*. 2015;24(9):642-6.
121. Maram J, Pan X, Sadda S, et al. Reproducibility of angle metrics using the time-domain anterior segment optical coherence tomography: Intra-observer and inter-observer variability. *Curr Eye Res*. 2015;40(5):496-500.
122. Montero JA, Ruiz-Moreno J. Optical coherence tomography characterisation of idiopathic central serous chorioretinopathy. *Br J Ophthalmol*. 2005;89(5): 562-564.
doi:10.1136/bjo.2004.049403
123. Piotrowiak I, Kaluzny BJ, Danek B, Chwiędacz A, Sikorski BL, Malukiewicz G. Spectral optical coherence tomography vs. fluorescein pattern for rigid gas-permeable lens fit. *Med Sci Monit*. 2014;20:1137-41.
124. Pavlin, CJ., Vasquez, LM., Lee, R., Simpson, ER., Ahmed, II. Anterior segment optical coherence tomography and ultrasound biomicroscopy in the imaging of anterior segment tumors. *Am J Ophthalmol*. 2009; 147(2):214-219.
125. Russell S, Bennett J, Wellman JA, et al. (2017). Efficacy and safety of voretigene neparvovec (AAV2-hRPE65v2) in patients with RPE65-mediated inherited retinal dystrophy: a randomised, controlled, open-label, phase 3 trial. *The Lancet*. 390 (10097), P849-860. Doi: [https://doi.org/10.1016/S0140-6736\(17\)31868-8](https://doi.org/10.1016/S0140-6736(17)31868-8)
126. Bussell II, Wollstein G, Schuman JS. OCT for glaucoma diagnosis, screening and detection of glaucoma progression. *Br J Ophthalmol*. 2014;98 Suppl 2:ii15-9.
127. Li G, Fansi AK, Boivin JF, Joseph L, Harasymowycz P. Screening for glaucoma in high risk populations using optical coherence tomography. *Ophthalmology* 2010;117(3):45361. DOI: 10.1016/j.ophtha.2009.07.033.
128. Waksman R, et al. Intravascular Ultrasound Versus Optical Coherence Tomography Guidance. *Journal of the American College of Cardiology* Vol. 62, No. 17, Suppl S, 2013
129. Zhang Y, et al. Comparison of intravascular ultrasound versus angiography-guided drug-eluting stent implantation: a meta-analysis of one randomised trial and ten observational studies involving 19,619 patients. *EuroIntervention* 2012;8:855–65.
130. Kim JS, et al. Randomized comparison of clinical outcomes between intravascular ultrasound and angiography-guided drug-eluting stent implantation for long coronary artery stenoses. *J Am Coll Cardiol Intv* 2013;6:369–76.
131. Witzensbichler B, et al. Use of IVUS reduces stent thrombosis: Results from the prospective, multicenter ADAPT-DES study. Available at: <http://www.tctmd.com/show.aspx?id¼116703>.
132. Prati F, Di Vito L, Biondi-Zoccai G, et al. Angiography alone versus angiography plus optical coherence tomography to guide decision making during percutaneous coronary intervention: the Centro per la Lotta contro l'Infarto-Optimisation of Percutaneous Coronary Intervention (CLI-OPCI) study. *EuroIntervention* 2012;8:823–9.

133. Sakata LM, Lavanya R, Friedman DS, et al. Comparison of gonioscopy and anterior segment ocular coherence tomography in detecting angle closure in different quadrants of the anterior chamber angle. *Ophthalmology*. 2008;115(5):769-774.
134. American Academy of Ophthalmology. Primary angle closure preferred practice patterns. San Francisco, CA: AAO. October 2010. Available at:
http://one.aao.org/CE/PracticeGuidelines/PPP_Content.aspx?cid=92bea8f6-5459-49a6-9233-4528343dc4c3.
135. Smith SD, Singh K, Lin SC, et al. Evaluation of the anterior chamber angle in glaucoma: A report by the American Academy of Ophthalmology. *Ophthalmology*. 2013;120(10):1985-1997.
136. Kashani et al. Optical coherence tomography angiography: A comprehensive review of current methods and clinical applications. *Prog Retin Eye Res*. 2017 Sep;60:66-100.
137. Koustenis A, Harris A, Gross J, Januleviciene I, Shah A, Siesky B. Optical coherence tomography angiography: an overview of the technology and an assessment of applications for clinical research. *Br J Ophthalmol*. 2017 Jan;101(1):16-20.
138. Retinal Vein Occlusions. Preferred Practice Pattern (PPP). American Academy of Ophthalmology. November 2015. Available at:
<https://www.aao.org/preferred-practice-pattern/retinal-vein-occlusions-ppp-2015>

Clinical Guideline Revision / History Information

Original Date: 7/31/2017

Reviewed/Revised: 1/18/2018, 7/31/2018, 7/23/2019, 7/21/2020, 08/04/2021, 12/01/2021, 07/26/2022, 07/19/2023, 07/29/2024