

Spravato (esketamine)

Disclaimer

Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.

Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.

Spravato (esketamine)	1
Summary	2
Definitions	3
Clinical Indications	4
Medical Necessity Criteria for Clinical Review	4
General Medical Necessity Criteria	4
Medical Necessity Criteria for Initial Clinical Review	4
Initial Indication-Specific Criteria	4
Treatment Resistant Depression (TRD)	4
Major Depressive Disorder (MDD) with Acute Suicidal Ideation or Behavior	5
Medical Necessity Criteria for Subsequent Clinical Review	5
Subsequent Indication-Specific Criteria	5
Treatment Resistant Depression (TRD)	5
Experimental or Investigational or Unproven / Not Medically Necessary	5
Applicable Billing Codes	6
References	7
Clinical Guideline Revision / History Information	9

Summary

Major depressive disorder is a mental health disorder that causes symptoms of sadness, hopelessness, and loss of interest in things. It can disrupt relationships and everyday activities, such as work, school, and activities that are usually pleasant. There are many medications from several classes that are available to treat major depressive disorder including selective serotonin-reuptake inhibitors (SSRIs), serotonin- and norepinephrine-reuptake inhibitors (SNRIs), tricyclic antidepressants, monoamine oxidase (MAO) inhibitors, and other antidepressants (e.g., bupropion, mirtazapine, trazodone).

Treatment resistant depression (TRD) refers to individuals who may improve partially but do not remit symptomatically nor regain full functional status. Partial response is the presence of residual symptoms.

In studies, TRD generally refers to major depressive episodes that do not respond satisfactorily after two (2) trials of antidepressant monotherapy. Response is generally classified by the amount of improvement from baseline on a depression rating scale:

- No response: improvement <25%
- Partial response: improvement 25% to 49%
- Response: improvement \geq 50% but less than the threshold for remission
- Remission: depression rating scale score less than or equal to a specific cutoff that defines the normal range.

It is estimated that at least 30% of individuals with depression have TRD. However, a notable percentage of individuals with TRD are pseudo-resistant (e.g., due to inadequacy of treatment trials or non-adherence to treatment).

Spravato (esketamine) is a non-competitive N-methyl D-aspartate (NMDA) receptor antagonist indicated for the treatment of:

1. Treatment-resistant depression (TRD) in adults, as monotherapy or in conjunction with an oral antidepressant; or
2. Depressive symptoms in adults with major depressive disorder (MDD) with acute suicidal ideation or behavior in conjunction with an oral antidepressant.

Limitations of use:

- The effectiveness of Spravato (esketamine) nasal spray in preventing suicide or in reducing suicidal ideation or behavior has not been demonstrated. Use of Spravato (esketamine) does not preclude the need for hospitalization if clinically warranted, even in those experiencing improvement after an initial dose.
- Spravato (esketamine) is not approved as an anesthetic agent; the safety and efficacy of Spravato (esketamine) as an anesthetic has not been established.

Due to the risk for sedation, dissociation, respiratory depression, abuse and misuse, Spravato (esketamine) is only available through a Risk Evaluation and Mitigation Strategy (REMS) Program. Spravato (esketamine) can only be administered at healthcare settings certified in the Spravato REMS Program. Patients treated in outpatient healthcare settings (e.g., medical offices and clinics) must be enrolled in the program.

Definitions

“Major depressive disorder”, also known as (MDD), is a psychiatric condition characterized by persistent low mood, low energy, or loss of interest in enjoyable activities causing substantial impairment in daily life. MDD is thought to be caused by a combination of genetic, environmental and psychological 2 factors. Risk factors include family history, major life changes, certain medications, chronic health problems, and substance use disorders.

“Depression assessment instruments” are used to screen and track treatment outcomes. These instruments may include interview and/or self-reported measures. Examples of validated and reliable instruments include Beck Depression Inventory-II [BDI-II], Hamilton Rating Scale for Depression [HAM-D], Inventory of Depressive Symptomatology–Clinician Rating [IDS-C], Montgomery-Asberg Depression Rating Scale [MADRS], Patient Health Questionnaire-9 [PHQ-9], and Quick Inventory of Depressive Symptomatology [QIDS-C16].

“Documentation” refers to written information, including but not limited to:

- Up-to-date chart notes, relevant test results, and/or relevant imaging reports to support diagnoses; or
- Prescription claims records, and/or prescription receipts to support prior trials of formulary alternatives.

“No evidence of” indicates that the reviewer has not identified any records of the specified item or condition within the submitted materials or claims history. In the absence of such evidence, the member is considered eligible. If any evidence of the item or condition is present upon review of the request, the member does not qualify.

“Risk Evaluation and Mitigation Strategy”, also known as (REMS), is a drug safety program that the Food and Drug Administration (FDA) requires for certain medications to ensure the benefits of the medication outweigh its risks.

“[s]” indicates state mandates may apply.

Clinical Indications

Medical Necessity Criteria for Clinical Review

General Medical Necessity Criteria

The Plan considers Spravato (esketamine) medically necessary when ALL of the following criteria are met:

1. The medication is prescribed by or in consultation with a psychiatrist; *AND*
2. The member is 18 years of age or older; *AND*
3. The member meets ALL of the following criteria:
 - a. No evidence of aneurysmal vascular disease (including thoracic and abdominal aorta, intracranial, and peripheral arterial vessels) or arteriovenous malformation; *and*
 - b. No evidence of history of intracerebral hemorrhage; *and*
 - c. No evidence of hypersensitivity to esketamine, ketamine, or any of the excipients; *AND*
4. Spravato (esketamine) is being prescribed at a dose and frequency that is within FDA approved labeling OR is supported by compendia or evidence-based published dosing guidelines for the requested indication; *AND*
5. The member meets the applicable [Medical Necessity Criteria for Initial Clinical Review](#) or [Subsequent Clinical Review](#) listed below.

Medical Necessity Criteria for Initial Clinical Review

Initial Indication-Specific Criteria

Treatment Resistant Depression (TRD)

The Plan considers Spravato (esketamine) medically necessary when ALL of the following criteria are met:

6. The member meets the above [General Medical Necessity Criteria](#); *AND*
7. The member has a diagnosis of TRD; *AND*
8. The member has a diagnosis of major depressive disorder (MDD) that is treatment resistant defined as nonresponse to an adequate trial (dosage, duration, and adherence) of at least TWO (2) antidepressants in an episode; *AND*
9. There is documentation of baseline scoring by a validated rating scale prior to starting therapy (e.g., BDI-II, HAM-D, IDS-C, MADRS, PHQ-9, QIDS-C16); *AND*
10. The member is unable to use all, or has tried and failed TWO (2) antidepressants, each from a different drug class for at least six (6) weeks each^[5]:
 - a. Aminoketones (e.g., bupropion); *and/or*
 - b. Monoamine oxidase inhibitors (e.g., selegiline, tranylcypromine); *and/or*
 - c. Noradrenaline and serotonergic antidepressants (e.g., mirtazapine); *and/or*
 - d. Selective serotonin reuptake inhibitors (e.g., citalopram, fluoxetine, paroxetine, sertraline); *and/or*

- e. Serotonin norepinephrine reuptake inhibitors (e.g., duloxetine, venlafaxine, desvenlafaxine); *and/or*
- f. Tricyclic antidepressants (e.g., amitriptyline, nortriptyline); *and/or*
- g. Serotonin modulators (e.g., nefazodone, trazodone, vilazodone); *and/or*
- h. Augmentation with lithium, thyroid hormone (e.g., liothyronine), atypical antipsychotics, or anticonvulsants.

If the above prior authorization criteria are met, the requested product will be authorized for up to 6 months.^[s]

Major Depressive Disorder (MDD) with Acute Suicidal Ideation or Behavior

The Plan considers Spravato (esketamine) medically necessary when ALL of the following criteria are met:

5. The member meets the above [General Medical Necessity Criteria](#); *AND*
6. The member has a diagnosis of major depressive disorder (MDD); *AND*
7. The member has current suicidal ideation or behavior with intent; *AND*
8. Spravato (esketamine) will be used in conjunction with an oral antidepressant (e.g. venlafaxine, escitalopram, duloxetine, sertraline, quetiapine).

If the above prior authorization criteria are met, the requested product will be authorized for up to 1 month.^[s]

Continued Care

[Medical Necessity Criteria for Subsequent Clinical Review](#)

Subsequent Indication-Specific Criteria

Treatment Resistant Depression (TRD)

Reauthorization for up to 12 months for Spravato (esketamine) will be granted if the member has recent (within the last 3 months) clinical chart documentation demonstrating ALL of the following criteria:

1. The member meets the above applicable [General Medical Necessity Criteria](#); *AND*
2. The member has experienced a documented improvement compared to baseline by a validated rating scale (e.g., BDI-II, HAM-D, IDS-C, MADRS, PHQ-9, QIDS-C16); *AND*
3. There is no evidence of unacceptable toxicity or adverse reactions to Spravato (esketamine).

[Experimental or Investigational or Unproven / Not Medically Necessary](#)^[s]

Spravato (esketamine) for any other indication or use is considered not medically necessary by the Plan, as it is deemed to be experimental, investigational, unproven, or not medically necessary. Non-covered indications include, but are not limited to, the following:

- Anesthetic agent. Spravato (eskatmine) is explicitly not approved as an anesthetic agent.

- Acute pain. Spravato (eskatmine) is explicitly not approved as an anesthetic agent, and should not be used for acute pain.
- Chronic pain. Spravato (eskatmine) is explicitly not approved as an anesthetic agent, and should not be used for chronic pain.
- Migraine headaches. There are no high quality studies to support the use of Spravato (eskatamine) for the use of migraine headaches.

Applicable Billing Codes

Table 1	
CPT/HCPCS Codes considered medically necessary for treatment resistant depression or major depressive disorder (MDD) with acute suicidal ideation or behavior if criteria are met:	
<i>Code</i>	<i>Description</i>
G2082	Office or other outpatient visit for the evaluation and management of an established patient that requires the supervision of a physician or other qualified health care professional and provision of up to 56 mg of esketamine nasal self-administration, includes 2 hours post-administration observation
G2083	Office or other outpatient visit for the evaluation and management of an established patient that requires the supervision of a physician or other qualified health care professional and provision of greater than 56 mg esketamine nasal self-administration, includes 2 hours post-administration observation
J0013	Esketamine, nasal spray, 1 mg

Table 2	
ICD-10 diagnostic codes considered medically necessary for treatment resistant depression or major depressive disorder (MDD) with acute suicidal ideation or behavior with Table 1 (CPT/HCPCS) codes if criteria are met:	
<i>Code</i>	<i>Description</i>
F06.31	Mood disorder due to known physiological condition with depressive features
F06.32	Mood disorder due to known physiological condition with major depressive-like episode
F32.1	Major depressive disorder, single episode, moderate
F32.2	Major depressive disorder, single episode, severe without psychotic features
F32.4	Major depressive disorder, single episode, in partial remission
F32.5	Major depressive disorder, single episode, in full remission

F32.89	Other specified depressive episodes
F32.9	Major depressive disorder, single episode, unspecified
F32.A	Depression, unspecified
F33.1	Major depressive disorder, recurrent, moderate
F33.2	Major depressive disorder, recurrent severe without psychotic features
F33.40	Major depressive disorder, recurrent, in remission, unspecified
F33.41	Major depressive disorder, recurrent, in partial remission
F33.42	Major depressive disorder, recurrent, in full remission
F33.8	Other recurrent depressive disorders
F33.9	Major depressive disorder, recurrent, unspecified

References

1. American Family Physician. Common Questions About the Pharmacologic Management of Depression in Adults. Available at: <https://www.aafp.org/pubs/afp/issues/2015/0715/p94.html#afp20150715p94-b51>. Accessed February 17, 2025.
2. APA. Depression Assessment Instruments. Available at: <https://www.apa.org/depression-guideline/assessment>. Accessed February 18, 2025.
3. ClinicalTrials.gov. A Study of Esketamine Nasal Spray, Administered as Monotherapy, in Adult Participants With Treatment-resistant Depression. <https://clinicaltrials.gov/study/NCT04599855>. Accessed February 17, 2025.
4. Daly EJ, Trivedi MH, Janik A, et al. Efficacy of Esketamine Nasal Spray Plus Oral Antidepressant Treatment for Relapse Prevention in Patients With Treatment-Resistant Depression: A Randomized Clinical Trial. *JAMA Psychiatry*. 2019 Sep 1;76(9):893-903.
5. Department of Veterans Affairs/Department of Defense. VA/DoD Clinical Practice Guideline for Assessment and Management of Patients at Risk for Suicide. Version 3.0 - 2024. Available at: https://www.healthquality.va.gov/HEALTHQUALITY/guidelines/MH/srb/VADOD-CPG-Suicide-Risk-Full-CPG-2024_Final_508.pdf. Accessed 7 April 2026.
6. Department of Veterans Affairs/Department of Defense. VA/DoD Clinical Practice Guideline for the Management of Major Depressive Disorder. Version 3.0 - 2024. Available at: https://www.healthquality.va.gov/HEALTHQUALITY/guidelines/MH/mdd/VADODMDDCPGFinal_508.pdf. Accessed 7 April 2026.
7. Fu DJ, Ionescu DF, Li X, et al. Esketamine Nasal Spray for Rapid Reduction of Major Depressive Disorder Symptoms in Patients Who Have Active Suicidal Ideation With Intent: Double-Blind, Randomized Study (ASPIRE I). *J Clin Psychiatry*. 2020 May 12;81(3):19m13191.
8. Guideline Development Panel for the Treatment of Depressive Disorders. APA Clinical Practice Guideline for the Treatment of Depression Across Three Age Cohorts. *Am Psychol*. 2019.
9. Ionescu DF, Fu DJ, Qiu X, et al. Esketamine Nasal Spray for Rapid Reduction of Depressive Symptoms in Patients With Major Depressive Disorder Who Have Active Suicide Ideation With Intent: Results of a Phase 3, Double-Blind, Randomized Study (ASPIRE II). *Int J Neuropsychopharmacol*. 2021 Jan 20;24(1):22-31.

10. Janik A, Qiu X, Lane R, et al. Esketamine Monotherapy in Adults With Treatment-Resistant Depression: A Randomized Clinical Trial. *JAMA Psychiatry*. 2025 Sep 1;82(9):877-887. doi: 10.1001/jamapsychiatry.2025.1317.
11. McIntyre RS, Alsuwaidan M, Baune BT, et al. Treatment-resistant depression: definition, prevalence, detection, management, and investigational interventions. *World Psychiatry*. 2023 Oct;22(3):394-412.
12. McIntyre RS, Rosenblat JD, Nemeroff CB, et al. Synthesizing the Evidence for Ketamine and Esketamine in Treatment-Resistant Depression: An International Expert Opinion on the Available Evidence and Implementation. *Am J Psychiatry*. 2021 May 1;178(5):383-399.
13. National Institute for Health and Care Excellence. Depression in adults: treatment and management. NICE Guidelines (NH222). Published 29 June 2022. Available at: <https://www.nice.org.uk/guidance/ng222/chapter/Recommendations>. Accessed 7 April 2026.
14. NP Psych Navigator. Understanding the Montgomery-Åsberg Depression Rating Scale (MADRS) Video. Available at: https://www.nppsychnavigator.com/NP_Perspectives/Clinical-Insights/Understanding-the-MADRS. Accessed February 17, 2025.
15. Popova V, Daly EJ, Trivedi M, et al. Efficacy and Safety of Flexibly Dosed Esketamine Nasal Spray Combined With a Newly Initiated Oral Antidepressant in Treatment-Resistant Depression: A Randomized Double-Blind Active-Controlled Study. *Am J Psychiatry*. 2019 Jun 1;176(6):428-438.
16. Qaseem A, Owens DK, Etzeandía-Ikobaltzeta I, et al. Nonpharmacologic and Pharmacologic Treatments of Adults in the Acute Phase of Major Depressive Disorder: A Living Clinical Guideline From the American College of Physicians (Version 1, Update Alert 4). *Ann Intern Med*. 2026 Mar 10. doi: 10.7326/ANNALS-26-00548. Epub ahead of print.
17. Spravato (esketamine hydrochloride solution) [prescribing information]. Titusville, NJ: Janssen Pharmaceuticals, Inc.; April 2025.
18. UptoDate. Unipolar treatment-resistant depression in adults: Epidemiology, risk factors, assessment, and prognosis. Available at: <https://www.uptodate.com/contents/unipolar-treatment-resistant-depression-in-adults-epidemiology-risk-factors-assessment-and-prognosis>. Accessed February 17, 2025.
19. U.S. Food & Drug Administration. Risk Evaluation and Mitigation Strategies REMS. Available at: <https://www.fda.gov/drugs/drug-safety-and-availability/risk-evaluation-and-mitigation-strategies-rems>. Accessed February 17, 2025.
20. Walter HJ, Abright AR, Bukstein OG, et al. Clinical Practice Guideline for the Assessment and Treatment of Children and Adolescents With Major and Persistent Depressive Disorders. *J Am Acad Child Adolesc Psychiatry*. 2023 May;62(5):479-502. doi: 10.1016/j.jaac.2022.10.001. Epub 2022 Oct 21.
21. World Health Organization. Duration of antidepressant treatment. Available at: <https://www.who.int/teams/mental-health-and-substance-use/treatment-care/mental-health-gap-action-programme/evidence-centre/depression/duration-of-antidepressant-treatment>. Accessed February 17, 2025.

Clinical Guideline Revision / History Information

Original Date: 07/01/2025

Reviewed/Revised: 06/01/2026