

## Growth Hormones (Lonapegsomatropin, Somapacitan, Somatropin, Somatrogon)

### Disclaimer

*Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.*

*Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.*

Growth Hormones (Lonapegsomatropin, Somapacitan, Somatropin, Somatrogon)	1
Summary	2
Definitions	2
Medical Necessity Criteria for Clinical Review	3
General Medical Necessity Criteria	3
Medical Necessity Criteria for Initial Clinical Review	3
Initial Indication-Specific Criteria	3
For the treatment of Adult Growth Hormone Deficiency	3
For the treatment of Pediatric Growth Hormone (GH) Deficiency	5
For the treatment of Turner Syndrome	5
For the treatment of persons who are/were Small for Gestational Age	6
For the treatment of Noonan Syndrome	6
For the treatment of Prader-Willi Syndrome	6
For the treatment of Short Stature homeobox-Containing Gene (SHOX) Deficiency	7
For the treatment of Growth Failure Associated with CKD, Cerebral Palsy, Congenital Adrenal Hyperplasia, Cystic Fibrosis, or Russell-Silver Syndrome	7
Medical Necessity Criteria for Subsequent Clinical Review	8
Subsequent Indication-Specific Criteria	8
For the treatment of Adult Growth Hormone Deficiency	8

For the treatment of Pediatric Growth Hormone Deficiency	8
For the treatment of Turner Syndrome	8
For the treatment of persons who are/were Small for Gestational Age	8
For the treatment of Noonan Syndrome	9
For the treatment of Prader-Willi Syndrome	9
For the treatment of Short Stature homeobox-Containing Gene (SHOX) Deficiency	9
For the treatment of Growth Failure Associated with CKD, Cerebral Palsy, Congenital Adrenal Hyperplasia, Cystic Fibrosis, or Russell-Silver Syndrome	9
<b>Experimental or Investigational / Not Medically Necessary</b>	10
References	11
Appendix A	13
Clinical Guideline Revision / History Information	14

## Summary

Growth hormone (GH) is produced by the pituitary somatotroph cells. GH production begins early in fetal life and continues throughout life, although at a progressively lower rate. The plan covers treatment for the following FDA-approved indications: pediatric patients with growth failure due to any of the following - growth hormone (GH) deficiency, Turner syndrome, Noonan syndrome, small for gestational age (SGA), Prader-Willi syndrome, chronic kidney disease (CKD), short stature homeobox-containing gene (SHOX) deficiency and adults with childhood-onset or adult-onset GH deficiency. The plan also covers certain compendial uses for GH including human immunodeficiency, growth failure associated with any of the following - cerebral palsy, congenital adrenal hyperplasia, cystic fibrosis, and Russell-Silver syndrome. All other indications are considered experimental / investigational or not medically necessary.

The Plan's preferred growth hormone products are Humatrope (somatropin) and Norditropin (somatropin).

Coverage for other (Non-Formulary) growth hormone products are preferred for certain conditions:

- Genotropin (somatropin) or Omnitrope (somatropin) is only provided when being prescribed for a member with Prader-Willi syndrome who is unable to use, or has tried and failed Norditropin (somatropin).
- Requests for non-formulary medications are also subject to Medical Necessity Criteria for Non-Formulary Products (PG069) Clinical Guideline.

## Definitions

"Compendia" are summaries of drug information and medical evidence to support decision-making about the appropriate use of drugs and medical procedures. Examples include, but are not limited to:

1. American Hospital Formulary Service Drug Information
2. Clinical pharmacology
3. National Comprehensive Cancer Network Drugs and Biologics Compendium
4. Thomson Micromedex DrugDex
5. United States Pharmacopeia-National Formulary (USP-NF)

“Growth Hormone Deficiency” is a condition characterized by growth failure and can be divided into congenital and acquired forms.

“Noonan Syndrome” is an autosomal dominant condition that is associated with short stature and congenital heart disease (CHD).

“Prader-Willi Syndrome” is a syndrome caused by the absence of expression of the paternally active genes on the long arm of chromosome 15.

“Russell-Silver Syndrome” is a syndrome characterized by severe intrauterine growth restriction and postnatal growth retardation with a prominent forehead, triangular face, downturned corners of the mouth, and body asymmetry (hemihypertrophy).

“Short Stature Homeobox-containing (SHOX) Deficiency” is a syndrome in which there are variants in the SHOX-containing gene on the X chromosome.

“Small for Gestational Age” is defined as a weight and/or length at birth that is at least -2 standard deviations (SD) below the mean for gestational age with lack of catch up growth by the age of 2 years old.

“Turner Syndrome” is a sex chromosome disorder caused by loss of part or all of an X chromosome.

### **Medical Necessity Criteria for Clinical Review**

#### **General Medical Necessity Criteria**

The Plan considers Growth Hormone medically necessary when ALL the following criteria are met:

1. The medication being requested meets BOTH of the following:
  - a. Is being prescribed for an FDA-approved or compendia supported indication; *and*
  - b. Is age-appropriate for the member based on FDA approval or is supported by evidence-based compendia; *AND*
2. Prescribed by, or in consultation with, a physician who specializes in the treatment of the specific condition such as an endocrinologist, pediatric endocrinologist, geneticist, or pediatric nephrologist; *AND*
3. The member is unable to use, or has tried and failed the Plan’s preferred growth hormone product(s); *AND*
4. The member meets the medical necessity criteria for the applicable indication listed below:

### **Medical Necessity Criteria for Initial Clinical Review**

#### **Initial Indication-Specific Criteria**

##### **For the treatment of Adult Growth Hormone Deficiency**

The Plan considers Growth Hormone medically necessary when BOTH of the following criteria are met:

1. The member has ONE (1) of the following:
  - a. TWO (2) pre-treatment pharmacologic provocative GH tests demonstrating GH levels less than (<) 5 ng/mL:
    - i. Unless the agent is Macrilen in which case a GH level of less than (<) 2.8 ng/mL confirms the presence of adult growth hormone deficiency; *or*
    - ii. Unless the agent is a glucagon stimulation test, in which case a peak GH level  $\leq$  3.0 ng/mL in those with a body mass index (BMI)  $\leq$  30 kg/m<sup>2</sup> and a high pretest probability of GHD (e.g., acquired structural abnormalities) OR a BMI < 25 kg/m<sup>2</sup> confirms the presence of adult growth hormone deficiency; *or*
    - iii. Unless the agent is a Glucagon stimulation test, in which case with a peak GH level  $\leq$  1.0 ng/mL in those with a BMI of  $\geq$  25 kg/m<sup>2</sup> and a low pretest probability of GHD (e.g., acquired structural abnormalities) OR a BMI > 30 kg/m<sup>2</sup> confirms the presence of adult growth hormone deficiency; *or*
  - b. ONE (1) pre-treatment pharmacologic provocative GH test demonstrating a GH level less than (<) 5 ng/mL **AND** a pre-treatment IGF-1 level that is low for age and gender:
    - i. Unless the agent is Macrilen in which case a GH level of less than (<) 2.8 ng/mL confirms the presence of adult growth hormone deficiency; *or*
    - ii. Unless the agent is a glucagon stimulation test, in which case a peak GH level  $\leq$  3.0 ng/mL in those with a body mass index (BMI)  $\leq$  30 kg/m<sup>2</sup> and a high pretest probability of GHD (e.g., acquired structural abnormalities) OR a BMI < 25 kg/m<sup>2</sup> confirms the presence of adult growth hormone deficiency; *or*
    - iii. Unless the agent is a Glucagon stimulation test, in which case with a peak GH level  $\leq$  1.0 ng/mL in those with a BMI of  $\geq$  25 kg/m<sup>2</sup> and a low pretest probability of GHD (e.g., acquired structural abnormalities) OR a BMI > 30 kg/m<sup>2</sup> confirms the presence of adult growth hormone deficiency; *or*
  - c. A structural abnormality of the hypothalamus or pituitary **AND BOTH** of the following:
    - i. Greater than > three (3) documented pituitary hormone deficiencies (e.g., adrenocorticotropic hormone [ACTH], antidiuretic hormone [ADH], follicle stimulating hormone [FSH], luteinizing hormone [LH], thyroid stimulating hormone [TSH], prolactin); *and*
    - ii. A pre-treatment IGF-1 level that is low for age and gender; *or*
  - d. A congenital abnormality of the hypothalamus or pituitary (e.g., pituitary transcription factor defects, GH-releasing hormone receptor gene defects) that persist throughout life and was diagnosed in infancy or childhood; **AND**
2. Chart documentation and supporting lab work are provided for review to substantiate the above-listed requirements.

If the above prior authorization criteria are met, the requested medication will be approved for up to 12 months.

### For the treatment of Pediatric Growth Hormone (GH) Deficiency

The Plan considers Growth Hormone medically necessary when BOTH of the following criteria are met:

1. The member meets ONE (1) of the following:
  - a. The member is a neonate or was diagnosed with GH deficiency as a neonate; *or*
  - b. The member is a newborn who has GH deficiency due to congenital hypopituitarism and meets ALL of the following:
    - i. The member is a newborn with hypoglycemia who does not attain a serum GH concentration above 5 mcg/L in the first week of life AND meets ONE (1) of the following:
      1. Deficiency of at least one (1) additional pituitary hormone; *or*
      2. Has classical imaging triad (i.e., congenital malformation) of ectopic posterior pituitary and pituitary hypoplasia with abnormal stalk; *or*
    - ii. The member meets ALL of the following clinical parameters:
      - i. The member has ONE (1) of the following:
        1. Two (2) pre-treatment pharmacologic provocative GH tests with both results demonstrating a peak GH level less than ( $<10$  ng/mL; *or*
        2. A documented pituitary or CNS disorder and a pre-treatment Insulin-like growth factor 1 (IGF-1) level greater than ( $>$  2 standard deviations (SD) below the mean; *and*
      - ii. The member has ONE (1) of the following:
        1. The member is under 2.5 years of age at initiation of treatment, a pre-treatment height greater than ( $>$  2 standard deviations (SD) below the mean; *or*
        2. The member is above 2.5 years of age at initiation of treatment AND ONE (1) of the following:
          - a. A pre-treatment height greater than  $>$  2 SD below the mean and a 1-year height velocity greater than  $>$  1 SD below the mean; *or*
          - b. A pre-treatment 2-year height velocity greater than  $>$  1.5 SD below the mean; *or*
          - c. A pre-treatment 1-year height velocity greater than  $>$  2 SD below the mean; *AND*
  2. Chart documentation and supporting lab work are provided for review to substantiate the above-listed requirements.

If the above prior authorization criteria are met, the requested medication will be approved for up to 12 months.

### For the treatment of Turner Syndrome

The Plan considers Growth Hormone medically necessary when ALL of the following criteria are met:

1. The member has a diagnosis of Turner Syndrome confirmed by karyotyping (genetic condition where one of the X chromosomes is partially or completely absent); *AND*

2. The member's pre-treatment height is less than (<) the 5th percentile for age; *AND*
3. The member's epiphyses are open; *AND*
4. Chart documentation and supporting lab work are provided for review to substantiate the above-listed requirements.

If the above prior authorization criteria are met, the requested medication will be approved for up to 12 months.

#### For the treatment of persons who are/were Small for Gestational Age

The Plan considers Growth Hormone medically necessary when ALL of the following criteria are met:

1. The member meets ONE (1) of the following:
  - a. Birth weight less than < 2500 g at gestational age greater than > 37 weeks; *or*
  - b. Birth weight or length less than less than < the 3rd percentile for gestational age; *or*
  - c. Birth weight or length greater than or equal to  $\geq 2$  SD below the mean for gestational age; *AND*
2. The member's pre-treatment age is greater than or equal to  $\geq 2$  years; *AND*
3. The member failed to manifest catch up growth (i.e. pre-treatment height greater than or equal to  $\geq 2$  SD below the mean); *AND*
4. The member's epiphyses are open; *AND*
5. Chart documentation and supporting laboratory test results are provided for review to substantiate the above-listed requirements.

If the above prior authorization criteria are met, the requested medication will be approved for up to 12 months.

#### For the treatment of Noonan Syndrome

The Plan considers Growth Hormone medically necessary when ALL of the following criteria are met:

1. The member meets ONE (1) of the following:
  - a. A pre-treatment height greater than > 2 SD below the mean and 1 year height velocity is greater than  $> 1$  SD below the mean; *or*
  - b. A pre-treatment 1-year height velocity greater than > 2 SD below the mean; *AND*
2. The member's epiphyses are open; *AND*
3. Chart documentation and supporting lab work are provided for review to substantiate the above-listed requirements.

If the above prior authorization criteria are met, the requested medication will be approved for up to 12 months.

#### For the treatment of Prader-Willi Syndrome

The Plan considers Growth Hormone medically necessary when BOTH of the following criteria are met:

1. The member has a diagnosis of Prader-Willi Syndrome confirmed by genetic testing showing ONE (1) of the following:
  - a. Deletion in the chromosome 15q11.2-q13 region; *or*
  - b. Maternal uniparental disomy in chromosome 15; *or*
  - c. Imprinting defects or translocations involving chromosome 15; *AND*
2. Chart documentation and supporting lab work are provided for review to substantiate the above-listed requirements.

If the above prior authorization criteria are met, the requested medication will be approved for up to 12 months.

For the treatment of Short Stature homeobox-Containing Gene (SHOX) Deficiency

The Plan considers Growth Hormone medically necessary when ALL of the following criteria are met:

1. The member has a diagnosis of SHOX deficiency confirmed by molecular or genetic analyses; *AND*
2. The member meets ONE (1) of the following:
  - a. A pre-treatment height greater than  $> 2$  SD below the mean and 1 year height velocity is greater than  $> 1$  SD below the mean; *or*
  - b. A pre-treatment 1-year height velocity greater than  $> 2$  SD below the mean; *AND*
3. The member's epiphyses are open; *AND*
4. Chart documentation and supporting lab work are provided for review to substantiate the above-listed requirements.

If the above prior authorization criteria are met, the requested medication will be approved for 12 months.

For the treatment of Growth Failure Associated with CKD, Cerebral Palsy, Congenital Adrenal Hyperplasia, Cystic Fibrosis, or Russell-Silver Syndrome

The Plan considers Growth Hormone medically necessary when ALL of the following criteria are met:

1. The member meets ONE (1) of the following:
  - a. The member is less than  $< 2.5$  years of age at initiation of treatment, a pre-treatment height greater than  $> 2$  SD below the mean; *or*
  - b. The member is greater than or equal to ( $\geq$ ) 2.5 years of age at initiation of treatment AND meets ONE (1) of the following:
    - i. A pre-treatment height greater than  $> 2$  SD below the mean and 1 year height velocity is greater than  $> 1$  SD below the mean; *or*
    - ii. A pre-treatment 2-year height velocity greater than  $> 1.5$  SD below the mean; *or*
    - iii. A pre-treatment 1-year height velocity greater than  $> 2$  SD below the mean; *AND*
2. The member's epiphyses are open; *AND*
3. Chart documentation and supporting lab work are provided for review to substantiate the above-listed requirements.

If the above prior authorization criteria are met, the requested medication will be approved for 12 months.

*Continued Care*

#### **Medical Necessity Criteria for Subsequent Clinical Review**

##### **Subsequent Indication-Specific Criteria**

###### **For the treatment of Adult Growth Hormone Deficiency**

Reauthorization for up to 12 months will be granted if clinical chart documentation is provided showing the member meets **ONE** of the following:

1. A clinical improvement, as evidenced by an improvement in the member's serum IGF-1; *OR*
2. Treatment plan indicating that the GH dose will be increased in response to a low IGF-1 serum concentration

###### **For the treatment of Pediatric Growth Hormone Deficiency**

Reauthorization for 12 months will be granted if clinical chart documentation is provided showing the member meets **ALL** of the following:

1. A clinical improvement in symptoms since starting the requested medication; *AND*
2. The member's epiphyses are open; *AND*
3. The member's growth rate is greater than  $> 2$  cm per year unless there is a documented clinical reason for lack of efficacy (i.e. on treatment less than  $<$  a year, nearing final adult height).

###### **For the treatment of Turner Syndrome**

Reauthorization for 12 months will be granted if clinical chart documentation is provided showing the member meets **ALL** of the following:

1. A clinical improvement in symptoms since starting the requested medication; *AND*
2. The member's epiphyses are open; *AND*
3. The member's growth rate is greater than  $> 2$  cm per year unless there is a documented clinical reason for lack of efficacy (i.e. on treatment less than  $<$  a year, nearing final adult height).

###### **For the treatment of persons who are/were Small for Gestational Age**

Reauthorization for 12 months will be granted if clinical chart documentation is provided showing the member meets **ALL** of the following:

1. A clinical improvement in symptoms since starting the requested medication; *AND*
2. The member's epiphyses are open; *AND*

3. The member's growth rate is greater than  $> 2$  cm per year unless there is a documented clinical reason for lack of efficacy (i.e. on treatment less than  $<$  a year, nearing final adult height).

#### For the treatment of Noonan Syndrome

Reauthorization for 12 months will be granted if clinical chart documentation is provided showing the member meets ALL of the following:

1. A clinical improvement in symptoms since starting the requested medication; *AND*
2. The member's epiphyses are open; *AND*
3. The member's growth rate is greater than  $> 2$  cm per year unless there is a documented clinical reason for lack of efficacy (i.e. on treatment less than  $<$  a year, nearing final adult height).

#### For the treatment of Prader-Willi Syndrome

Reauthorization for 12 months will be granted if clinical chart documentation is provided showing the member meeting ONE (1) of the following:

1. Clinical improvement in symptoms since starting the requested medication; *OR*
2. Stabilization in the member's body composition or psychomotor function.

#### For the treatment of Short Stature homeobox-Containing Gene (SHOX) Deficiency

Reauthorization for 12 months will be granted if clinical chart documentation is provided showing the member meets ALL of the following:

1. A clinical improvement in symptoms since starting the requested medication; *AND*
2. The member's epiphyses are open; *AND*
3. The member's growth rate is greater than  $> 2$  cm per year unless there is a documented clinical reason for lack of efficacy (i.e. on treatment less than  $<$  a year, nearing final adult height).

#### For the treatment of Growth Failure Associated with CKD, Cerebral Palsy, Congenital Adrenal Hyperplasia, Cystic Fibrosis, or Russell-Silver Syndrome

Reauthorization for 12 months will be granted if clinical chart documentation is provided showing the member meets ALL of the following:

1. A clinical improvement in symptoms since starting the requested medication; *AND*
2. The member's epiphyses are open; *AND*
3. The member's growth rate is greater than  $> 2$  cm per year unless there is a documented clinical reason for lack of efficacy (i.e. on treatment less than  $<$  a year, nearing final adult height).

### Experimental or Investigational / Not Medically Necessary

Growth hormone products for any other indication or use is considered not medically necessary by the Plan, as it is deemed to be experimental, investigational, unproven, or not medically necessary. Non-covered indications include, but are not limited to, the following:

- Achondroplasia
- Amyotrophic Lateral Sclerosis (ALS)
- Anorexia Nervosa (AN)
- Ataxia-Telangiectasia (A-T)
- Cognitive Developmental Disorder
- Crohn's Disease (CD)
- Dwarfism
- Eating Disorders
- Female Infertility
- Female Infertility Due to Diminished Ovarian Reserve
- Heart Failure
- Hypophosphatemic Rickets
- Idiopathic Short Stature (ISS)
- Juvenile Idiopathic Arthritis (JIA)
- Lipodystrophies
- Mucopolysaccharidosis IH
- Mucopolysaccharidosis Type I (MPS I)
- Mucopolysaccharidosis Type II (MPS II)
- Mucopolysaccharidosis Type VI
- Non-alcoholic Fatty Liver Disease, NAFLD
- Obesity
- Osteogenesis Imperfecta (OI)
- Osteopenia (Disorder)
- Osteoporosis
- Phelan McDermid Syndrome
- Polycystic Ovarian Syndrome (PCOS)
- Preimplantation Genetic Testing (PGT)
- Primary Disease Fascioscapulohumeral Dystrophy (FSHD)
- Septo-Optic Dysplasia
- Short Bowel Syndrome (SBS)
- Short Stature
- Still's Disease, Juvenile Onset
- Type 1 Diabetes Mellitus

## References

1. American Association of Clinical Endocrinologists medical guidelines for clinical practice for growth hormone use in adults and children--2003 update. *Endocr Pract*. 2003; 9(1): p64-76.
2. Biller BM, Samuels MH, Zagar A, Cook DM, Arafah BM, Bonert V, Stavrou S, Kleinberg DL, Chipman JJ, Hartman ML. Sensitivity and specificity of six tests for the diagnosis of adult GH deficiency. *J Clin Endocrinol Metab*. 2002 May;87(5):2067-79. doi: 10.1210/jcem.87.5.8509. PMID: 11994342.
3. Blum WF, Ross JL, Zimmermann AG, et al. GH treatment to final height produces similar height gains in patients with SHOX deficiency and Turner syndrome: results of a multicenter trial. *J Clin Endocrinol Metab*. 2013 Aug;98(8):E1383-92. doi: 10.1210/jc.2013-1222. Epub 2013 May 29. Erratum in: *J Clin Endocrinol Metab*. 2013 Dec;98(12):4992.
4. Bondy CA; Turner Syndrome Study Group. Care of girls and women with Turner syndrome: a guideline of the Turner Syndrome Study Group. *J Clin Endocrinol Metab*. 2007 Jan;92(1):10-25. doi: 10.1210/jc.2006-1374. Epub 2006 Oct 17. PMID: 17047017.
5. Butler MG, Miller JL, Forster JL. Prader-Willi Syndrome - Clinical Genetics, Diagnosis and Treatment Approaches: An Update. *Curr Pediatr Rev*. 2019;15(4):207-244. doi: 10.2174/1573396315666190716120925.
6. Clayton PE, Cianfarani S, Czernichow P, et al. Management of the child born small for gestational age through to adulthood: a consensus statement of the International Societies of Pediatric Endocrinology and the Growth Hormone Research Society. *J Clin Endocrinol Metab* 2007; 92:804.
7. Cohen P, Rogol AD, Deal CL, et al, "Consensus Statement on the Diagnosis and Treatment of Children With Idiopathic Short Stature: A Summary of the Growth Hormone Research Society, the Lawson Wilkins Pediatric Endocrine Society, and the European Society for Paediatric Endocrinology Workshop," *J Clin Endocrinol Metab*, 2008, 93(11):4210-7.
8. Collett-Solberg PF, Ambler G, Backeljauw PF, et al. Diagnosis, genetics, and therapy of short stature in children: a Growth Hormone Research Society International perspective. *Horm Res Paediatr*. 2019;92(1):1-14.
9. Cook DM, Yuen KC, Biller BM, Kemp SF, Vance ML; American Association of Clinical Endocrinologists (AACE). American Association of Clinical Endocrinologists medical guidelines for clinical practice for growth hormone use in growth hormone-deficient adults and transition patients - 2009 update. *Endocr Pract*. 2009;15(suppl 2):S1-S29.
10. Davenport ML, Crowe BJ, Travers SH, et al. Growth hormone treatment of early growth failure in toddlers with Turner syndrome: a randomized, controlled, multicenter trial. *J Clin Endocrinol Metab*. 2007 Sep;92(9):3406-16. doi: 10.1210/jc.2006-2874. Epub 2007 Jun 26.
11. Deal C, Hasselmann C, Pfäffle RW, et al. Associations between pituitary imaging abnormalities and clinical and biochemical phenotypes in children with congenital growth hormone deficiency: data from an international observational study. *Horm Res Paediatr*. 2013;79(5):283-92. doi: 10.1159/000350829. Epub 2013 May 16.
12. Deal CL, Tony M, Höybye C, et al. GrowthHormone Research Society workshop summary: consensus guidelines for recombinant human growth hormone therapy in Prader-Willi syndrome. *J Clin Endocrinol Metab*. 2013 Jun;98(6):E1072-87. doi: 10.1210/jc.2012-3888. Epub 2013 Mar 29.
13. de Boer H, Blok GJ, Popp-Snijders C, et al. Monitoring of growth hormone replacement therapy in adults, based on measurement of serum markers. *J Clin Endocrinol Metab* 1996; 81:1371.
14. Dichtel LE, Yuen KC, Bredella MA, et al. Overweight/Obese adults with pituitary disorders require lower peak growth hormone cutoff values on glucagon stimulation testing to avoid overdiagnosis of growth hormone deficiency. *J Clin Endocrinol Metab*. 2014 Dec;99(12):4712-9. doi: 10.1210/jc.2014-2830.
15. Fleseriu M, Hashim IA, Karavitaki N, et al. Hormonal replacement in hypopituitarism in adults: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2016;101(11):3888-3921. doi:10.1210/jc.2016-2118.
16. Franklin SL, Geffner ME. Growth hormone: the expansion of available products and indications. *Endocrinol Metab Clin North Am*. 2009 Sep;38(3):587-611. doi: 10.1016/j.ecl.2009.06.006.
17. Gallagher MP, Levine LS, Oberfield SE. A review of the effects of therapy on growth and bone mineralization in children with congenital adrenal hyperplasia. *Growth Horm IGF Res*. 2005 Jul;15 Suppl A:S26-30. doi: 10.1016/j.ghir.2005.06.006.

18. Genotropin (somatropin) [prescribing information]. New York, NY: Pharmacia & Upjohn Co; April 2019.
19. Genotropin MiniQuick (somatropin) [prescribing information]. New York, NY: Pharmacia & Upjohn Co; April 2019.
20. Gharib H, Cook DM, Saenger PH, et al. American Association of Clinical Endocrinologists medical guidelines for clinical practice for growth hormone use in adults and children--2003 update. *Endocr Pract.* 2003 Jan-Feb;9(1):64-76. doi: 10.4158/EP.9.1.64. Erratum in: *Endocr Pract.* 2008 Sep;14(6):802-3. Gharib, Hossein [added]; Cook, David M [added]; Saenger, Paul H [added]; Bengtsson, Bengt-Ake [added]; Feld, Stanley [added]; Nippoldt, Todd B [added]; Rodbard, Helena W [added]; Seibel, John A [added]; Vance, Mary Lee [added]; Zimmerman, Donald [added].
21. Gravholt CH, Andersen NH, Christin-Maitre S, et al. Clinical practice guidelines for the care of girls and women with Turner syndrome. *Eur J Endocrinol.* 2024 Jun 5;190(6):G53-G151. doi: 10.1093/ejendo/lvae050.
22. Grimberg A, DiVall SA, Polychronakos C, et al. Guidelines for growth hormone and insulin-like growth factor-I treatment in children and adolescents: growth hormone deficiency, idiopathic short stature, and primary insulin-like growth factor-I deficiency. *Horm Res Paediatr.* 2016;86(6):361-397. doi:10.1159/000452150
23. Humatrope (somatropin) [prescribing information]. Indianapolis, IN: Lilly USA LLC; October 2019.
24. Lin-Su K, Vogiatzi MG, Marshall I, et al. Treatment with growth hormone and luteinizing hormone releasing hormone analog improves final adult height in children with congenital adrenal hyperplasia. *J Clin Endocrinol Metab.* 2005 Jun;90(6):3318-25. doi: 10.1210/jc.2004-2128. Epub 2005 Mar 29.
25. Macrilen (macimorelin) [prescribing information]. Frankfurt, Germany: Aeterna Zentaris GmbH. December 2017.
26. Molitch ME, Clemons DR, Malozowski S, Merriam GR, Vance ML; Endocrine Society (ES). Evaluation and treatment of adult growth hormone deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab.* 2011;96(6):1587-1609.
27. Ngenla (somatrogon) [prescribing information]. New York, NY: Pfizer Labs; June 2023.
28. Norditropin (somatropin) [prescribing information]. Plainsboro, NJ: Novo Nordisk Inc; February 2018.
29. Nutropin AQ (somatropin) [prescribing information]. South San Francisco, CA: Genentech; December 2016.
30. Nutropin AQ NuSpin (somatropin) [product monograph]. Mississauga, Ontario, Canada: Hoffmann-La Roche Limited; June 2018.
31. Omnitrope (somatropin) [prescribing information]. Princeton, NJ: Sandoz Inc; June 2019.
32. Quintos JB, Vogiatzi MG, Harbison MD, New MI. Growth hormone therapy alone or in combination with gonadotropin-releasing hormone analog therapy to improve the height deficit in children with congenital adrenal hyperplasia. *J Clin Endocrinol Metab.* 2001 Apr;86(4):1511-7. doi: 10.1210/jcem.86.4.7412.
33. Russell-Silver Syndrome - National Organization for Rare Diseases, 9th edition. 2020. Available at: <https://rarediseases.org/rare-diseases/russell-silver-syndrome/>
34. Saizen (somatropin) [prescribing information]. Rockland, MA: EMD Serono Inc; February 2020.
35. Serostim (somatropin) [prescribing information]. Rockland, MA: EMD Serono Inc; June 2019.
36. Shim ML, Moshang T Jr, Oppenheim WL, Cohen P. Is treatment with growth hormone effective in children with cerebral palsy? *Dev Med Child Neurol.* 2004 Aug;46(8):569-71. doi: 10.1017/s0012162204000945.
37. Skytrofa (lonapegsomatropin) [prescribing information]. Palo Alto, CA: Ascendis Pharma Inc; August 2021.
38. Sogroya (somapacitan-beco) [prescribing information]. Plainsboro, NJ: Novo Nordisk Inc; April 2023.
39. Thaker V, Carter B, Putman M. Recombinant growth hormone therapy for cystic fibrosis in children and young adults. *Cochrane Database Syst Rev.* 2021 Aug 23;8(8):CD008901. doi: 10.1002/14651858.CD008901.pub5.
40. Wakeling EL, Brioude F, Lokulo-Sodipe O, et al. Diagnosis and management of Silver-Russell syndrome: first international consensus statement. *Nat Rev Endocrinol.* 2017 Feb;13(2):105-124. doi: 10.1038/nrendo.2016.138. Epub 2016 Sep 2.
41. Wilson TA, Rose SR, Cohen P, et al. Update of guidelines for the use of growth hormone in children: the Lawson Wilkins Pediatric Endocrinology Society Drug and Therapeutics Committee. *J Pediatr.* 2003 Oct;143(4):415-21. doi: 10.1067/s0022-3476(03)00246-4.

42. Yuen KCJ, Biller BMK, Radovick S, et al. American Association of Clinical Endocrinologists and American College of Endocrinology Guidelines for Management of Growth Hormone Deficiency in Adults and Patients Transitioning from Pediatric to Adult Care. AACE Guidelines, Nov 2019. 25(11): p1191-1232. DOI: <https://doi.org/10.4158/GL-2019-0405>

43. Yuen KCJ, Biller BMK, Radovick S, et al. American Association of Clinical Endocrinologists and American College of Endocrinology guidelines for management of growth hormone deficiency in adults and patients transitioning from pediatric to adult care. Endocr Pract. 2019;25(11):1191-1232. doi:10.4158/GL-2019-0405.

44. Zomacton (somatropin) [prescribing information]. Parsippany, NJ: Ferring Pharmaceuticals; July 2018.

45. Zomacton [package insert]. Parsippany, NJ: Ferring Pharmaceuticals Inc.; July 2018.

46. Zorbive (somatropin) [prescribing information]. Rockland, MA: EMD Serono Inc; September 2019.

## Appendix A

Table 1: Growth Hormones

Drug	FDA-Approved or Compendia-Supported Indications					
	Growth Hormone Deficiency		Turner Syndrome	SGA	Idiopathic Short Stature	Other
	Pediatric	Adults				
Genotropin (Somatropin)	✓	✓	✓	✓	✓	PWS
Humatrope (Somatropin)	✓	✓	✓	✓*	✓	SHOX
Ngenla (somatrogon-ghla)	✓					
Norditropin (Somatropin)	✓	✓	✓	✓*	✓	NS, PWS
Nutropin (Somatropin)	✓	✓	✓		✓	CKD
Omnitrope (Somatropin)	✓	✓	✓	✓	✓	PWS
Saizen (Somatropin)	✓	✓				
Serostim (Somatropin)						HIV
Skytrofa (Lonapegsomatropin-tcgd)	✓	✓				
Sogroya (Somapacitan-beco)	✓	✓				
Zomacton (Somatropin)	✓	✓	✓	✓*	✓	SHOX
Zorbive <sup>#</sup> (Somatropin)						SBS

NOTE: All Growth Hormone medications require prior authorization.

### Acronyms:

CKD = Growth failure due to chronic kidney disease

HIV = HIV-associated Wasting or Cachexia

NS = Noonan Syndrome

PWS = Prader-Willi Syndrome in children

SBS = Short Bowel Syndrome

SGA = Growth failure in children born Small for Gestational Age who fail to manifest catch-up growth by age 2 years of age (\* – indicated in patients with no catch-up growth by 2 to 4 years of age)

SHOX = Short stature homeobox-containing gene deficiency

# Discontinued by Manufacturer

#### Clinical Guideline Revision / History Information

Original Date: 08/06/2020

Reviewed/Revised: 06/24/2021, 12/01/2021, 06/23/2022, 9/15/2022, 9/21/2023, 12/19/2024,  
02/02/2026