

## Neffy (epinephrine nasal spray)

### Disclaimer

*Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.*

*Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.*

Neffy (epinephrine nasal spray)	1
Summary	1
Definitions	2
Clinical Indications	2
Medical Necessity Criteria for Clinical Review	2
Indication-Specific Criteria	2
Type I Allergic Reactions	2
Experimental or Investigational / Not Medically Necessary	3
References	3
Appendix A	5
Clinical Guideline Revision / History Information	5

### Summary

Type 1 allergic reactions are known as immediate hypersensitivity reactions and include anaphylaxis. Anaphylaxis is a serious, life-threatening allergic reaction that can occur within seconds or minutes of exposure to an allergen. It can be triggered by IgE-mediated reactions to allergens such as foods, drugs, insect stings, and latex. Epinephrine is the first-line treatment for anaphylaxis. Current clinical guidelines

recommend all patients at risk of anaphylaxis be prescribed epinephrine and trained on proper administration.

Epinephrine helps alleviate bronchospasm, wheezing and dyspnea that may occur during anaphylaxis. Epinephrine also alleviates pruritus (itching), urticaria (hives), and angioedema (rapid swelling, typically in the face, lips, tongue and throat). It may also relieve gastrointestinal and genitourinary symptoms associated with anaphylaxis.

Neffy (epinephrine nasal spray) is an intranasal formulation of epinephrine approved by the FDA for the emergency treatment of allergic reactions, including anaphylaxis, in those who weigh 15 kg or greater. It provides a non-injectable option for administration of epinephrine. Neffy (epinephrine nasal spray) is administered as a single spray into one nostril. In absence of clinical improvement or if symptoms worsen after initial treatment, administer a second dose of Neffy (epinephrine nasal spray) in the same nostril with a new nasal spray starting 5 minutes after the first dose. As such, it is recommended that individuals are prescribed and have immediate access to two Neffy (epinephrine nasal spray) nasal sprays at all times.

### Definitions

"Anaphylaxis" refers to a serious, potentially life-threatening allergic reaction that can occur within seconds or minutes of exposure to an allergen and requires immediate treatment with epinephrine.

"Epinephrine" refers to a medication that acts on alpha and beta-adrenergic receptors to quickly reverse the symptoms of anaphylaxis such as airway swelling, hypotension, and wheezing.

"Epinephrine auto-injector" is a medical device used for the automatic injection of a measured dose of epinephrine, most commonly for the treatment of anaphylaxis. Examples include EpiPen, Auvi-Q, generic versions.

"Needle phobia" is an extreme fear of medical procedures involving injections or hypodermic needles, which can lead to avoidance of medically necessary treatment.

"No evidence of" indicates that the reviewer has not identified any records of the specified item or condition within the submitted materials or claims history. In the absence of such evidence, the member is considered eligible. If any evidence of the item or condition is present upon review of the request, the member does not qualify.

"[s]" indicates state mandates may apply.

## Clinical Indications

### Medical Necessity Criteria for Clinical Review

#### Indication-Specific Criteria

##### Type I Allergic Reactions

The Plan considers Neffy (epinephrine nasal spray) medically necessary when ALL of the following criteria are met:

1. The member weighs ( $\geq$ ) 15 kg or greater; *AND*
2. The member has diagnosis of an allergy that may require emergency epinephrine treatment (e.g., food allergy, insect venom allergy, exercise-induced anaphylaxis, idiopathic anaphylaxis); *AND*
3. Neffy (epinephrine nasal spray) is being prescribed for the acute treatment of allergic reactions, including anaphylaxis; *AND*
4. The member meets ONE (1) of the following:
  - a. Has experienced an inadequate response, intolerance, or contraindication to an epinephrine auto-injector product (e.g., EpiPen, generic epinephrine auto-injector)<sup>[s]</sup>; *or*
  - b. Is unable to appropriately use an epinephrine auto-injector due to a clinically valid reason (e.g., needle phobia, lack of dexterity, visual impairment, arthritis); *AND*
5. No evidence of severe obstructive pathology that would reasonably impair drug delivery; *AND*
6. Neffy (epinephrine nasal spray) is being prescribed at a dose and frequency that is within FDA approved labeling OR is supported by compendia or evidence-based published dosing guidelines for the requested indication (see [Appendix A](#), Table 1 below).

*The requested medication is being used within the Plan's Quantity Limit of:*

- Neffy solution nasal spray 1 mg/0.1 ml: 4 devices per 28 days
- Neffy solution nasal spray 2 mg/0.1 ml: 4 devices per 28 days

NOTE: It is recommended that individuals are prescribed and have immediate access to two Neffy (epinephrine nasal sprays) at all times.

If the above prior authorization criteria are met, the requested product will be authorized for up to 12-months.<sup>[s]</sup>

#### Experimental or Investigational or Unproven / Not Medically Necessary<sup>[s]</sup>

Neffy (epinephrine nasal spray) for any other indication or use is considered not medically necessary by the Plan, as it is deemed to be experimental, investigational, unproven, or not medically necessary.

Non-covered indications include, but are not limited to, the following:

- Use in members less than 15 kg. The safety and efficacy of Neffy (epinephrine nasal spray) has not been studied in this population.
- Use for the prevention of anaphylaxis or allergic reactions. Epinephrine is not intended to be used to prevent an allergic or anaphylactic reaction.

- Use for any non-FDA approved indications.

## References

1. Brügggen MC, Walsh S, Ameri MM, et al. Management of Adult Patients With Drug Reaction With Eosinophilia and Systemic Symptoms: A Delphi-Based International Consensus. *JAMA Dermatol.* 2024 Jan 1;160(1):37-44. doi: 10.1001/jamadermatol.2023.4450.
2. Campbell RL, Kelso JM. Anaphylaxis: emergency treatment. Post TW, ed. UpToDate. Waltham, MA: UpToDate Inc.
3. Campbell et al; Members of the Joint Task Force; Practice Parameter Workgroup. Emergency department diagnosis and treatment of anaphylaxis: a practice parameter. *Ann Allergy Asthma Immunol.* 2014 Dec;113(6):599-608. doi: 10.1016/j.anai.2014.10.007. PMID: 25466802.
4. Canadian Society of Allergy and Clinical Immunology. Eight Tests and Treatments to Question: Don't use antihistamines as first-line treatment in anaphylaxis. May 2025. Available at: <https://choosingwiselycanada.org/recommendation/allergy-clinical-immunology/?highlight=anaphylaxis>. Accessed 24 February 2026.
5. Cardona et al, Worm M. World allergy organization anaphylaxis guidance 2020. *World Allergy Organ J.* 2020 Oct 30;13(10):100472. doi: 10.1016/j.waojou.2020.100472. PMID: 33204386; PMCID: PMC7607509.
6. Casale TB, Ellis AK, Nowak-Wegrzyn A, Kaliner M, Lowenthal R, Tanimoto S. Pharmacokinetics/pharmacodynamics of epinephrine after single and repeat administration of neffy, EpiPen, and manual intramuscular injection. *J Allergy Clin Immunol.* 2023 Dec;152(6):1587-1596. doi: 10.1016/j.jaci.2023.08.007. Epub 2023 Aug 19.
7. Crescioli G, Giovannini M, Pessina B, et al. Epinephrine nasal spray for the treatment of anaphylaxis: perspectives in pediatrics. *Curr Opin Allergy Clin Immunol.* 2025 Dec 1;25(6):511-517. doi: 10.1097/ACI.0000000000001109. Epub 2025 Sep 16.
8. Dworaczyk et al. A 13.2 mg epinephrine intranasal spray demonstrates comparable pharmacokinetics, pharmacodynamics, and safety to a 0.3 mg epinephrine autoinjector. *J Allergy Clin Immunol Glob.* 2023 Dec 12;3(2):100200. doi: 10.1016/j.jacig.2023.100200. PMID: 38328805; PMCID: PMC10847913.
9. Dworaczyk et al. Randomized trial of pharmacokinetic and pharmacodynamic effects of 13.2 mg intranasal epinephrine treatment in congestion. *Ann Allergy Asthma Immunol.* 2024 Aug;133(2):186-193.e2. doi: 10.1016/j.anai.2024.04.033. Epub 2024 May 7. PMID: 38719149.
10. Ebisawa M, Takahashi K, Takahashi K, et al. Epinephrine Nasal Spray Improves Allergic Symptoms in Patients Undergoing Oral Food Challenge, Phase 3 Trial. *J Allergy Clin Immunol Pract.* 2025 Oct;13(10):2787-2794. doi: 10.1016/j.jaip.2025.06.038. Epub 2025 Jul 8.
11. Epinephrine injection (auto-injector) [prescribing information]. Lineage Therapeutics Inc.; Horsham, PA: May 2016.
12. EpiPen/EpiPen Jr (epinephrine) [prescribing information]. St. Louis, MO: Meridian Medical Technologies LLC; February 2023.
13. Fleischer DM, Li HH, Talreja N, et al. Pharmacokinetics and Pharmacodynamics of neffy, Epinephrine Nasal Spray, in Pediatric Allergy Patients. *J Allergy Clin Immunol Pract.* 2025 Jun;13(6):1335-1341.e1. doi: 10.1016/j.jaip.2025.03.019. Epub 2025 Mar 20.
14. Gaudio et al. Wilderness Medical Society Clinical Practice Guidelines on Anaphylaxis. *Wilderness Environ Med.* 2022 Mar;33(1):75-91. doi: 10.1016/j.wem.2021.11.009. Epub 2022 Feb 2. Erratum in: *Wilderness Environ Med.* 2022 Jun;33(2):263. doi: 10.1016/j.wem.2022.03.009. PMID: 35120856.
15. Golden et al. Anaphylaxis: A 2023 practice parameter update. *Ann Allergy Asthma Immunol.* 2024 Feb;132(2):124-176. doi: 10.1016/j.anai.2023.09.015. Epub 2023 Dec 18. PMID: 38108678.
16. Iglesia EGA, Kwan M, Virkud YV, Iweala OI. Management of Food Allergies and Food-Related Anaphylaxis. *JAMA.* 2024 Feb 13;331(6):510-521. doi: 10.1001/jama.2023.26857.
17. Li X, Ma Q, Yin J, et al. A Clinical Practice Guideline for the Emergency Management of Anaphylaxis (2020). *Front Pharmacol.* 2022 Mar 28;13:845689. doi: 10.3389/fphar.2022.845689.

18. Lieberman et al. Anaphylaxis--a practice parameter update 2015. *Ann Allergy Asthma Immunol.* 2015 Nov;115(5):341-84. doi: 10.1016/j.anai.2015.07.019. PMID: 26505932.
19. Muraro A, Worm M, Alviani C, et al. EAACI guidelines: Anaphylaxis (2021 update). *Allergy.* 2022 Feb;77(2):357-377. doi: 10.1111/all.15032. Epub 2021 Sep 1.
20. Neffy (epinephrine) nasal spray [prescribing information]. San Diego, CA: ARS Pharmaceuticals Operations Inc; March 2026.
21. Perman SM, Elmer J, Maciel CB, Uzendu A, May T, Mumma BE, Bartos JA, Rodriguez AJ, Kurz MC, Panchal AR, Rittenberger JC; American Heart Association. 2023 American Heart Association Focused Update on Adult Advanced Cardiovascular Life Support: An Update to the American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation.* 2024 Jan 30;149(5):e254-e273. doi: 10.1161/CIR.0000000000001194. Epub 2023 Dec 18. PMID: 38108133.
22. Pflipsen MC, Vega Colon KM. Anaphylaxis: Recognition and Management. *Am Fam Physician.* 2020 Sep 15;102(6):355-362.
23. Pouessel G, Neukirch C. Alternatives to Injectable Adrenaline for Treating Anaphylaxis. *Clin Exp Allergy.* 2025 Jan;55(1):36-51. doi: 10.1111/cea.14598. Epub 2024 Nov 24.
24. Shaker et al. Anaphylaxis-a 2020 practice parameter update, systematic review, and Grading of Recommendations, Assessment, Development and Evaluation (GRADE) analysis. *J Allergy Clin Immunol.* 2020 Apr;145(4):1082-1123. doi: 10.1016/j.jaci.2020.01.017. Epub 2020 Jan 28. PMID: 32001253.
25. Sicherer SH, Simons FER; SECTION ON ALLERGY AND IMMUNOLOGY. Epinephrine for First-aid Management of Anaphylaxis. *Pediatrics.* 2017 Mar;139(3):e20164006. doi: 10.1542/peds.2016-4006. Epub 2017 Feb 13. PMID: 28193791.
26. Takahashi K, Yanagida N, Sato S, Ebisawa M. Intranasal adrenaline: A new treatment for anaphylaxis. *Pediatr Allergy Immunol.* 2026 Mar;37(3):e70316. doi: 10.1111/pai.70316.
27. US Food and Drug Administration (FDA). FDA Drug Safety Communication. MedWatch. FDA alerts patients and health care professionals of EpiPen auto-injector errors related to device malfunctions and user administration. <https://www.fda.gov/drugs/drug-safety-and-availability/fda-alerts-patients-and-health-care-professionals-epipen-auto-injector-errors-related-device>. Published March 24, 2020. Accessed September 5, 2024.

## Appendix A

Table 1: Recommended Dosage of Neffy (epinephrine nasal spray)

Member's Weight	Dosage
30 kg or Greater	One spray of Neffy 2 mg
15 kg to less than 30 kg	One spray of Neffy 1 mg

Each Neffy (epinephrine nasal spray) nasal spray is for single use and delivers the entire dose upon activation.

In absence of clinical improvement or if symptoms worsen after initial treatment, administer a second dose of Neffy (epinephrine nasal spray) in the same nostril with a new nasal spray starting 5 minutes after the first dose. As such, it is recommended that individuals are prescribed and have immediate access to two Neffy (epinephrine nasal spray) nasal sprays at all times.

Clinical Guideline Revision / History Information

Original Date: 09/18/2024  
Reviewed/Revised: 10/01/2025, 05/01/2026