

Extavia (interferon beta-1b)

Disclaimer

Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.

Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.

Summary

Multiple sclerosis (MS) is a chronic, inflammatory, demyelinating disease of the central nervous system. It typically presents in young adults (generally diagnosed before 50 years of age) with symptoms such as vision problems, muscle weakness, numbness, and difficulty with balance and coordination. The most common form is relapsing-remitting MS (occurring in about 85% of patients), characterized by acute attacks followed by periods of remission. Treatment goals include reducing relapses, slowing disability progression, and managing symptoms. Disease-modifying therapies (DMTs) are the primary treatment approach and include injectable medications (e.g., interferons, glatiramer acetate), oral medications (e.g., dimethyl fumarate, fingolimod, teriflunomide, etc.), and infusion therapies (e.g., natalizumab, ocrelizumab).

Extavia (interferon beta-1b) is a first-generation DMT indicated for relapsing forms of MS, including clinically isolated syndrome, relapsing-remitting disease, and active secondary progressive disease in adults. It is administered via subcutaneous injection every other day and works by reducing inflammation and modulating the immune response. Extavia (interferon beta-1b) works by modulating the immune system and reducing inflammation in the central nervous system, though its exact mechanism of action in MS is not fully understood.

Definitions

"Clinically isolated syndrome" refers to a first episode of neurologic symptoms lasting at least 24 hours caused by inflammation or demyelination in the central nervous system.

"Disease-modifying therapy" is a medication that modifies the course of MS by reducing relapses and slowing disability progression.

"Multiple sclerosis" is a chronic autoimmune disease of the central nervous system characterized by inflammation, demyelination, and neurodegeneration.

"Primary progressive MS" refers to worsening neurologic function from the onset of symptoms, without early relapses or remissions.

"Relapse" is defined as the appearance of new symptoms or the worsening of existing symptoms lasting at least 24 hours in the absence of fever or infection.

"Relapsing-remitting MS" refers to a disease course characterized by clearly defined attacks of new or increasing neurologic symptoms followed by periods of partial or complete recovery.

"Secondary progressive MS" is a disease course following relapsing-remitting MS that is characterized by a progressive worsening of neurologic function over time with or without relapses.

Medical Necessity Criteria for Initial Authorization

The Plan considers Extavia (interferon beta-1b) medically necessary when recent (within the last 3 months) clinical chart documentation provided indicates the member meets ALL of the following:

1. Prescribed by or in consultation with a neurologist or physician who specializes in the treatment of multiple sclerosis; *AND*
2. Is 18 years of age or older; *AND*
3. Has ONE of the following forms of multiple sclerosis:
 - a. relapsing-remitting (RRMS); *or*
 - b. active secondary progressive disease (SPMS); *or*
 - c. clinically isolated syndrome (CIS); *AND*
4. Extavia (interferon beta-1b) will be used as monotherapy for multiple sclerosis (i.e., member is not using and will not use other disease-modifying MS therapies while on Extavia); *AND*
5. Extavia (interferon beta-1b) is being prescribed within the manufacturer's published dosing guidelines or falls within dosing guidelines found in a compendia of current literature.
 - o *The recommended dose is 0.25 mg injected subcutaneously every other day.*
 - i. *Up to 15 vials/30 days (1 kit/30 days)*

If the above prior authorization criteria are met, the requested product will be authorized for up to 12-months.

Medical Necessity Criteria for Reauthorization

Reauthorization for up to 12-months will be granted if the member has recent (within the last 6-months) clinical documentation showing BOTH of the following:

1. The requested medication is prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis; *AND*
2. The member has experienced at least ONE of the following:
 - a. Improvement in at least ONE objective measure, such as:
 - i. Reduced disease activity on MRI; *and/or*
 - ii. Improved or stable disability scores; *and/or*
 - iii. Reduced relapse rate; *and/or*
 - iv. Improved fatigue or walking assessments; *AND/OR*
 - b. The member has shown stabilization or improvement in at least ONE MS symptom, such as:
 - i. Motor function; *and/or*
 - ii. Fatigue; *and/or*
 - iii. Vision; *and/or*
 - iv. Bowel/bladder function; *and/or*
 - v. Spasticity; *and/or*
 - vi. Walking/gait; *and/or*
 - vii. Pain/numbness/tingling.

Experimental or Investigational / Not Medically Necessary

Extavia (interferon beta-1b) for any other indication or use is considered not medically necessary by the Plan, as it is deemed to be experimental, investigational, or unproven. Non-covered indications include, but are not limited to, the following:

- Use in combination with other disease-modifying therapies for multiple sclerosis.
- Treatment of non-relapsing forms of multiple sclerosis (e.g., primary progressive MS).
- Use for the treatment of other neurological conditions not related to multiple sclerosis.
- Use as a treatment for cancer or as an adjunct to cancer therapies.
- Use for the treatment of viral infections, including chronic viral hepatitis.
- Use in pediatric patients under 18 years of age.
- Use in the management of other autoimmune disorders not related to multiple sclerosis.

References

1. Bainbridge JL, Miravalle A, Wong PS. Multiple Sclerosis. In DiPiro JT, Yee GC, Posey LM, et al, eds. *Pharmacotherapy: A Pathophysiologic Approach*. 11th ed. New York, NY: McGraw-Hill; 2019.
2. Extavia (interferon beta-1b) [prescribing information]. East Hanover, NJ: Novartis Pharmaceuticals Corporation; July 2023.
3. Hauser SL, Cree BAC. Treatment of Multiple Sclerosis: A Review. *Am J Med*. 2020 Dec;133(12):1380-1390.e2. doi: 10.1016/j.amjmed.2020.05.049. Epub 2020 Jul 17.
4. Interferon beta-1b in the treatment of multiple sclerosis: final outcome of the randomized controlled trial. The IFNB Multiple Sclerosis Study Group and The University of British Columbia MS/MRI Analysis Group. *Neurology*. 1995 Jul;45(7):1277-85.
5. Interferon beta-1b is effective in relapsing-remitting multiple sclerosis. I. Clinical results of a multicenter, randomized, double-blind, placebo-controlled trial. The IFNB Multiple Sclerosis Study Group. *Neurology*. 1993 Apr;43(4):655-61. doi: 10.1212/wnl.43.4.655.
6. Kappos L, Edan G, Freedman MS, et al,. Long-term clinical outcomes in patients with CIS treated with interferon beta-1b: results from the 15-year follow up of the BENEFIT trial. *J Neurol*. 2024 Jul;271(7):4599-4609. doi: 10.1007/s00415-024-12417-x. Epub 2024 May 10.
7. McGinley MP, Goldschmidt CH, Rae-Grant AD. Diagnosis and Treatment of Multiple Sclerosis: A Review. *JAMA*. 2021;325(8):765–779. doi:10.1001/jama.2020.26858
8. Montalban X, Gold R, Thompson AJ, et al.ECTRIMS/EAN guideline on the pharmacological treatment of people with multiple sclerosis [published correction appears in *Eur J Neurol*. 2018;25(3):605]. *Eur J Neurol*. 2018;25(2):215-237. doi:10.1111/ene.13536.
9. Multiple Sclerosis Society of Canada. Disease-modifying therapies. <https://mssociety.ca/managing-ms/treatments/medications/disease-modifying-therapies-dmts>.
10. National MS Society. Disease-modifying therapies for MS (updated March 2022). Available from National MS Society website: <https://nms2cdn.azureedge.net/cmssite/nationalmssociety/media/msnationalfiles/brochures/brochure-the-ms-disease-modifying-medications.pdf>.
11. Paty DW, Li DK. Interferon beta-1b is effective in relapsing-remitting multiple sclerosis. II. MRI analysis results of a multicenter, randomized, double-blind, placebo-controlled trial. UBC MS/MRI Study Group and the IFNB Multiple Sclerosis Study Group. *Neurology*. 1993 Apr;43(4):662-7. doi: 10.1212/wnl.43.4.662.
12. Placebo-controlled multicentre randomised trial of interferon beta-1b in treatment of secondary progressive multiple sclerosis. European Study Group on interferon beta-1b in secondary progressive MS. *Lancet*. 1998 Nov 7;352(9139):1491-7.
13. Rae-Grant A, Day GS, Marrie RA, et al. Practice guideline recommendations summary: Disease-modifying therapies for adults with multiple sclerosis: Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology. *Neurology*. 2018;90(17):777-788.
14. Reich DS, Lucchinetti CF, Calabresi PA. 2018. Multiple sclerosis. *New England Journal of Medicine* 378(2):169-180
15. The use of disease-modifying therapies in multiple sclerosis: principles and current evidence summary. Multiple Sclerosis Coalition. Available from the National MS Society Website: <https://www.nationalmssociety.org/>.
16. Tramacere I, Del Giovane C, Salanti G, et al. Immunomodulators and immunosuppressants for relapsing-remitting multiple sclerosis: a network meta-analysis. *Cochrane Database Syst Rev* 2015;9:CD011381.
17. Yang, J., Rempe, T., Whitmire, N., Dunn-Pirio, A., & Graves, J. (2022). Therapeutic Advances in Multiple Sclerosis. *Frontiers in Neurology*, 13. <https://doi.org/10.3389/fneur.2022.824926>.

Clinical Guideline Revision / History Information

Original Date: 06/27/2024

Reviewed/Revised: 10/01/2025