

### Anesthesia and Sedation in Endoscopic Procedures

#### Disclaimer

*Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.*

*Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.*

#### Summary

The Plan members undergoing certain endoscopic procedures may require different levels of sedation, depending on the procedure and their existing medical conditions. General anesthesia is one type of sedation where a physician anesthesiologist or nurse anesthetist can induce a state of complete unconsciousness. This is different from conscious sedation, where a patient is awake but medications are given for pain and to help with relaxation. The vast majority of patients undergoing endoscopies, where a small camera is inserted to look for disease such as cancer or precancerous masses, are sedated with conscious sedation to make the procedure more tolerable. This guideline provides criteria for members who may require general anesthesia, monitored anesthesia care (MAC), or deep sedation for both upper and lower gastrointestinal endoscopic procedures (i.e., upper endoscopy, colonoscopy). The treating provider should counsel the member on the appropriate anesthesia options and levels of risk.

#### Definitions

"Local Anesthesia" is when a medication is used to induce an absence of pain/sensation to a specific region or part of the body.

"Minimal Sedation" is also referred to as anxiolysis and is defined as the lowest level of drug-induced impaired cognition. Persons who are minimally sedated respond normally to commands and their respirations/cardiovascular system are unaffected. Examples of minimal sedatives include oral benzodiazepines.

“Moderate Sedation” was historically known as “Conscious Sedation” and refers to a deeper depression of consciousness compared to minimal sedation. Persons who are moderately sedated will typically still respond to verbal commands, sometimes requiring tactile stimuli, and their respirations/cardiovascular system are unaffected. Examples of medications used for moderate sedation are IV benzodiazepines and opioids. During moderate sedation the responsible physician typically assumes the dual role of performing the procedure and supervising the sedation (an anesthesia provider is not involved with moderate sedation).

“Monitored Anesthesia Care (MAC)” is a combination of local anesthesia together with sedation and/or analgesia, in a setting monitored by a trained anesthesiologist (physician), an certified registered nurse anesthetist (CRNA), or certified anesthesiologist assistants. While Monitored Anesthesia Care may include the administration of sedatives and/or analgesics often used for Moderate Sedation, the qualified anesthesia provider of MAC is focused exclusively on the member for airway, hemodynamic and physiologic changes. Furthermore, the provider of MAC must be prepared and qualified to convert to general anesthesia. As described by American Society of Anesthesiologists, Monitored Anesthesia Care is a physician service that is clearly distinct from Moderate Sedation due to the expectations and qualifications of the provider who must be able to utilize all anesthesia resources to support life and to provide patient comfort and safety during a diagnostic or therapeutic procedure.

“Deep Sedation/Analgesia” is part of a continuum from monitored anesthesia care to general anesthesia and results in significant central nervous system depression, where patients are no longer conscious and are more difficult to arouse. They will typically respond to painful stimuli but not to verbal or simple tactile stimuli. The respiratory system is depressed and they may need some ventilatory support. Examples of medications used for deep sedation are IV benzodiazepines, opioids, propofol, ketamine, etomidate, and dexmedetomidine. This level of anesthesia can be provided only by qualified anesthesia providers.

“General Anesthesia” is the deepest form of sedation, where there is complete loss of consciousness and no arousability to stimuli. Cardiovascular and respiratory function are often impaired and may require monitoring and assistance such as ventilatory support. General anesthesia always requires monitoring by an anesthesia provider.

“American Society for Anesthesia Physical Status Classes” are used to risk stratify patients prior to procedures. The categories were developed and defined by the American Society for Anesthesiologists and are as follows:

- ASA Physical Status Class I: A normal, healthy person
- ASA Physical Status Class II: A person with mild systemic disease without functional limitations, including but not limited to:
  - Current smoker
  - Social alcohol drinker

- Pregnancy
- Obesity (BMI <40)
- Well-controlled diabetes mellitus, hypertension, well-controlled lung disease, well-controlled COPD, etc.
- **ASA Physical Status Class III:** A person with severe systemic disease or not adequately controlled resulting in substantial functional limitations, including but not limited to:
  - Poorly controlled diabetes mellitus or hypertension
  - Poorly controlled Chronic obstructive pulmonary disease (COPD)
  - Morbid obesity (BMI ≥40)
  - Active hepatitis
  - Active alcohol dependence or abuse
  - Implanted pacemaker
  - Moderate reduction of ejection fraction
  - End stage renal disease (ESRD) undergoing regularly scheduled dialysis
  - Premature infant with post-conceptual age < 60 weeks
  - History (>3 months) of myocardial infarction (MI), cerebrovascular accident (CVA or stroke), transient ischemic attack (TIA or mini-stroke), or coronary artery disease (CAD)/stents.
- **ASA Physical Status Class IV:** A person with severe systemic disease that is a constant threat to life; including but not limited to:
  - Recent (< 3 months) MI, CVA, TIA, or CAD/stents
  - Ongoing cardiac ischemia
  - Severe valve dysfunction
  - Severe reduction of ejection fraction
  - Sepsis
  - Disseminated intravascular coagulation (DIC)
  - ESRD not undergoing regularly scheduled dialysis
- **ASA Physical Status Class V:** A moribund person who is not expected to survive without the operation; including but not limited to:
  - Ruptured abdominal/thoracic aneurysm
  - Massive trauma
  - Intracranial bleed with mass effect
  - Ischemic bowel in the face of significant cardiac pathology
  - Multiple organ/system dysfunction
- **ASA Physical Status Class VI:** A declared brain-dead person whose organs are being removed for donor purposes.

“Modified Mallampati Classification” is used to assess airways and risk for difficult tracheal intubation.

The original Mallampati Classification has class I-III. The modified Mallampati Classification has class I-IV.

- Class 0: Ability to see any part of the epiglottis upon mouth opening and tongue protrusion
- Class I: Soft palate, fauces, uvula, pillars visible

- Class II: Soft palate, fauces, uvula visible
- Class III: Soft palate, base of uvula visible
- Class IV: Soft palate not visible at all

“Endoscopy” refers to a procedure where a small camera is inserted to visualize internal parts of the body. Examples include the following:

- Colonoscopy (looking at the colon)
- Sigmoidoscopy (looking at the sigmoid portion of the colon)
- Esophagoduodenoscopy (i.e., EGD, where the upper GI system is visualized)
- ERCP (endoscopic retrograde cholangiopancreatography)

### Clinical Indications

The Plan considers general anesthesia, MAC, or deep sedation requiring anesthesiologist or anesthesiologist attendance medically necessary for use in upper or lower gastrointestinal endoscopy in “high-risk” members, defined as those with ONE of the following criteria:

1. Significant medical condition, as defined by at least ONE of the following:
  - a. ASA Physical Status Class III-V; *or*
    - i. *Note:* In specific cases of ETOH dependence/abuse, stable chronic ETOH dependence can be performed under moderate sedation for ASA III. As per American Society of Gastrointestinal Endoscopy minimal and/or moderate sedation can be delivered safely by endoscopists to patients who are ASA Class I, II, or III. Although alcohol dependence or abuse falls within ASA III, the concern for complications and the request for higher level of anesthesia is appropriate when alcohol has affected other organs (e.g., cirrhosis, end-stage liver disease), or systemic effects such as acute alcohol intoxication.
  - b. Pregnancy; *or*
  - c. Epilepsy; *or*
  - d. Age less than 18; *or*
  - e. Age greater than 70, one other criteria must also be met under 1-4.
2. Risk for airway compromise, as defined by at least ONE of the following:
  - a. Current evidence of obstructive sleep apnea or stridor; *or*
  - b. Dysmorphic facial features, such as Pierre-Robin or Down syndrome; *or*
  - c. Modified Mallampati classification Class III-IV; *or*
  - d. Neck abnormalities, such as neck obesity, short neck, limited neck extension, cervical spine instability, cervical spine disease or trauma, neck mass, tracheal deviation, advanced rheumatoid arthritis (risk of cervical instability), cranial nerve IX or X dysfunction; *or*
  - e. Jaw abnormalities, such as micrognathia, retrognathia, trismus, significant macro-occlusion; *or*

- f. Morbid obesity (defined as body mass index [BMI] >40 or >35 with hypertension, coronary artery disease, obstructive sleep apnea, or Type 2 diabetes).
3. Anticipated intolerance to standard sedatives, as defined by at least **ONE** of the following:
    - a. Previous reaction or complication with sedation or anesthesia; *or*
    - b. Opiate, sedative, or hypnotic dependence or tolerance due to chronic use; *or*
    - c. Active alcohol or substance abuse.
  4. Situations where deep sedation or general anesthesia may be required, as defined by at least **ONE** of the following:
    - a. Uncooperative or combative patients (e.g., those with dementia, psychiatric disorders, or young children); *or*
    - b. Complex or prolonged procedures or invasive therapeutic endoscopies, such as the following:
      - i. Endoscopic retrograde cholangiopancreatography [ERCP]
      - ii. Endoscopic ultrasound [EUS]
      - iii. Esophageal stenting
      - iv. Emergency therapeutic procedures such as acute GI bleeding
      - v. A patient with a complex condition who may experience more pain or discomfort
      - vi. A patient with a history of prolonged procedures or history of difficult intubation
      - vii. A patient requiring a repeat endoscopic procedure (e.g., long or tortuous colon) due to the inability to be sedated with conscious/moderate sedation at the initial endoscopic procedure or due to the result of another issue from the initial endoscopic procedure that a repeat was medically warranted.

#### Experimental or Investigational / Not Medically Necessary

Sedation requiring anesthesiologist or anesthesiologist care (e.g., general anesthesia, MAC, or deep sedation) is not considered medically necessary in members with an “average risk” of surgical complications, as defined as ASA Class I-II and not meeting “high-risk” criteria above, who are undergoing standard upper or lower endoscopic procedures. These members should undergo procedures with moderate sedation/analgesia as indicated and provided by the endoscopist, where it is not medically necessary for an anesthesiologist to be present.

Applicable Billing Codes (HCPCS/CPT Codes)

<i>Anesthesia and Sedation in Endoscopy</i>	
CPT/HCPCS Codes considered medically necessary if criteria are met:	
<i>Code</i>	<i>Description</i>
00731	Anesthesia for upper gastrointestinal endoscopic procedures, endoscope introduced proximal to duodenum; not otherwise specified
00732	Anesthesia for upper gastrointestinal endoscopic procedures, endoscope introduced proximal to duodenum; endoscopic retrograde cholangiopancreatography (ERCP)
00811	Anesthesia for lower intestinal endoscopic procedures, endoscope introduced distal to duodenum; not otherwise specified
00812	Anesthesia for lower intestinal endoscopic procedures, endoscope introduced distal to duodenum; screening colonoscopy
00813	Anesthesia for combined upper and lower gastrointestinal endoscopic procedures, endoscope introduced both proximal to and distal to the duodenum
99100	Anesthesia for patient of extreme age, younger than 1 year and older than 70 (List separately in addition to code for primary anesthesia procedure)
99140	Anesthesia complicated by emergency conditions (specify) (List separately in addition to code for primary anesthesia procedure)
99151	Moderate sedation services provided by the same physician or other qualified health care professional performing the diagnostic or therapeutic service that the sedation supports, requiring the presence of an independent trained observer to assist in the monitoring of the patient's level of consciousness and physiological status; initial 15 minutes of intraservice time, patient younger than 5 years of age
99152	Moderate sedation services provided by the same physician or other qualified health care professional performing the diagnostic or therapeutic service that the sedation supports, requiring the presence of an independent trained observer to assist in the monitoring of the patient's level of consciousness and physiological status; initial 15 minutes of intraservice time, patient age 5 years or older

99153	Moderate sedation services provided by the same physician or other qualified health care professional performing the diagnostic or therapeutic service that the sedation supports, requiring the presence of an independent trained observer to assist in the monitoring of the patient's level of consciousness and physiological status; each additional 15 minutes intraservice time (List separately in addition to code for primary service)
99155	Moderate sedation services provided by a physician or other qualified health care professional other than the physician or other qualified health care professional performing the diagnostic or therapeutic service that the sedation supports; initial 15 minutes of intraservice time, patient younger than 5 years of age
99156	Moderate sedation services provided by a physician or other qualified health care professional other than the physician or other qualified health care professional performing the diagnostic or therapeutic service that the sedation supports; initial 15 minutes of intraservice time, patient age 5 years or older
99157	Moderate sedation services provided by a physician or other qualified health care professional other than the physician or other qualified health care professional performing the diagnostic or therapeutic service that the sedation supports; each additional 15 minutes intraservice time (List separately in addition to code for primary service)
G0500	Moderate sedation services provided by the same physician or other qualified health care professional performing a gastrointestinal endoscopic service that sedation supports, requiring the presence of an independent trained observer to assist in the monitoring of the patient's level of consciousness and physiological status; initial 15 minutes of intra-service time; patient age 5 years or older (additional time may be reported with 99153, as appropriate)
ICD-10 codes considered medically necessary if criteria are met:	
<i>Code</i>	<i>Description</i>
E66.01	Morbid (severe) obesity due to excess calories
E66.2	Morbid obesity
F10.10 - F19.99	Mental and behavioral disorders due to psychoactive substance use
G40.0-G40.8	Epilepsy and recurrent seizures

G47.30	Sleep apnea, unspecified
G47.31	Primary central sleep apnea
G47.33	Obstructive sleep apnea (adult) (pediatric)
G47.37	Central sleep apnea in conditions classified elsewhere
G47.39	Other sleep apnea
J35.1	Hypertrophy of tonsils
M26.02	Maxillary hypoplasia
M26.04	Mandibular hypoplasia
M26.09	Other specified anomalies of jaw size [micrognathia]
M26.19	Other specified anomalies of jaw-cranial base relationship [retrognathia]
P28.30 - P28.39	Primary sleep apnea of newborn
Q18.8	Other specified congenital malformations of face and neck
Q38.2	Macroglossia
Q75.0- Q75.9	Other congenital malformations of the skull and face bones
Q90.0 - Q90.9	Down syndrome
R06.1	Stridor
R22.1	Localized swelling, mass and lump, neck
R25.2	Cramp and spasm
S12.000A - S12.9XXS	Fracture of cervical vertebra and other parts of neck
S14.2XXA - S14.2XXS	Injury of nerve root of cervical spine
Z33.1	Pregnant state, incidental
Z33.3	Pregnant state, gestational carrier



## References

1. Adams MA, Prenovost KM, Dominitz JA, et al. National Trends in Use of Monitored Anesthesia Care for Outpatient Gastrointestinal Endoscopy in the Veterans Health Administration. *JAMA Intern Med.* 2017;177(3):436-438.
2. Adams MA, Saleh A, Rubenstein JH. A systematic review of factors associated with utilization of monitored anesthesia care for gastrointestinal endoscopy. *Gastroenterol Hepatol (N Y).* 2016; 12(6):361-370.
3. Aisenberg J, Miller KM. Propofol for endoscopic sedation: A protocol for safe and effective administration by the gastroenterologist. *Gastrointest Endosc.* 2003;58(5):725-732.
4. American Society for Gastrointestinal Endoscopy. Guidelines for sedation and anesthesia in GI endoscopy. *Gastrointest Endosc.* 2018;87(2):327-337. doi: 10.1016/j.gie.2017.07.018
5. American Society for Gastrointestinal Endoscopy. ASGE guideline: modifications in endoscopic practice for the elderly. *Gastrointestinal Endoscopy.* 2013; 78(1):1-7.
6. American Society for Gastrointestinal Endoscopy. Modifications in endoscopic practice for pediatric patients. *Gastrointestinal Endoscopy.* 2014; 79(5):699-710.
7. American Society for Gastrointestinal Endoscopy. Sedation and anesthesia in GI endoscopy. *Gastrointestinal endoscopy.* 2008; 68(5):815-826.
8. American Society of Anesthesiologists task force on sedation and analgesia by non-anesthesiologists. Practice guidelines for sedation and analgesia by non-anesthesiologists. *Anesthesiology.* 2002; 96(4):1004-1017.
9. American Society of Anesthesiologists. Continuum of depth of sedation definition of general anesthesia and levels of sedation/analgesia (approved by ASA House of Delegates on October 13, 1999, and amended on October 15, 2014). For additional information visit the ASA website: <http://www.asahq.org>. Accessed on August 12, 2017.
10. American Society of Anesthesiologists. (Oct 2014). ASA Physical Status Classification System. Last amended December 13, 2020. Accessed Aug 25, 2023 <https://www.asahq.org/standards-and-guidelines/asa-physical-status-classification-system>
11. American Society of Anesthesiologists. (2019, October 23). Continuum of Depth of Sedation: Definition of General Anesthesia and Levels of Sedation/Analgesia. <https://www.asahq.org/standards-and-guidelines/continuum-of-depth-of-sedation-definition-of-general-anesthesia-and-levels-of-sedationanalgesia>
12. American Society of Anesthesiologists. (2018, October 17). Distinguishing Monitored Anesthesia Care ("MAC") from Moderate Sedation/Analgesia (Conscious Sedation). <https://www.asahq.org/standards-and-guidelines/distinguishing-monitored-anesthesia-care-mac-from-moderate-sedationanalgesia-conscious-sedation>
13. American Society of Anesthesiologists. Position on monitored anesthesia care (approved by the House of Delegates on October 25, 2005; last amended on October 16, 2013). For additional information visit the ASA website: <http://www.asahq.org>. Accessed on August 12, 2017
14. American Society of Anesthesiologists. Practice guidelines for the management of the difficult airway: an updated report by the American Society of Anesthesiologists task force on management of the difficult airway. *Anesthesiology.* 2013; 118(2):251-271.

15. American Society of Anesthesiologists. Statement on anesthesia care for endoscopic procedures. Approved by the ASA House of Delegates on October 15, 2014. For additional information visit the ASA website: <http://www.asahq.org>. Accessed on August 12, 2017.
16. American Society of Anesthesiologists. Statement on respiratory monitoring during endoscopic procedures (approved by the ASA House of Delegates on October 21, 2009; last amended on October 15, 2014). For additional information visit the ASA website: <http://www.asahq.org>. Accessed on August 12, 2017.
17. Amornyotin S. Sedation and monitoring for gastrointestinal endoscopy. *World J Gastrointest Endosc* 2013; 5:47.
18. Brown CA. Approach to the anatomically difficult airway in adults outside the operating room. UpToDate.com. Last updated May 6, 2022.  
[https://www.uptodate.com/contents/approach-to-the-anatomically-difficult-airway-in-adults-outside-the-operating-room?search=modified%20mallampati%20classification&source=search\\_result&selectedTitle=1~150&usage\\_type=default&display\\_rank=1](https://www.uptodate.com/contents/approach-to-the-anatomically-difficult-airway-in-adults-outside-the-operating-room?search=modified%20mallampati%20classification&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1)
19. Carlsson U, Grattidge, P. Sedation for upper gastrointestinal endoscopy: A comparative study of propofol and midazolam. *Endoscopy*. 1995;27:240-243.
20. Cataldo PA. Colonoscopy without sedation. *Dis Colon Rectum*. 1996;39:257.
21. Chen WX, Lin HJ, Zhang WF, et al. Sedation and safety of propofol for therapeutic endoscopic retrograde cholangiopancreatography. *Hepatobiliary Pancreat Dis Int*. 2005;4(3):437-440.
22. Cohen J. Overview of procedural sedation for gastrointestinal endoscopy. UpToDate. Waltham, MA: UpToDate; 2017
23. Cohen LB, Delegee MH, Aisenberg J, et al. AGA Institute review of endoscopic sedation. *Gastroenterology*. 2007; 133(2):675-701.
24. Cohen LB, Hightower CD, Wood DA, et al. Moderate level sedation during endoscopy: A prospective study using low-dose propofol, meperidine/fentanyl, and midazolam. *Gastrointest Endosc*. 2004;59(7):795-803.
25. Cooper GS, Kou TD, Rex DK. Complications following colonoscopy with anesthesia assistance: a population-based analysis. *JAMA Intern Med*. 2013 Apr 8;173(7):551-6.
26. De Gregorio BT, Poorman JC, Katon RM. Peroral ultrathin endoscopy in adult patients. *Gastrointest Endosc*. 1997;45:303.
27. Dhir V, Swaroop VS, Vazifdar KF, Wagle SD. Topical pharyngeal anesthesia without intravenous sedation during upper gastrointestinal endoscopy. *Indian J Gastroenterol*. 1997;16:10.
28. Early DS, Lightdate JR, Vargo JJ, et al. Guidelines for sedation and anesthesia in GI. *American Society for Gastrointestinal Endoscopy. Gastrointestinal Endoscopy*, 2018;87(2): 327 - 337.
29. Hajouli, S., & Ludhwani, D. (2020). Heart failure and ejection fraction.
30. Hinkelbein J, Lamperti M, Akeson J, Santos J, Costa J, De Robertis E, Longrois D, Novak-Jankovic V, Petrini F, Struys MMRF, Veyckemans F, Fuchs-Buder T, Fitzgerald R. European Society of Anaesthesiology and European Board of Anaesthesiology guidelines for procedural sedation and analgesia in adults. *Eur J Anaesthesiol*. 2018;35(1):6.
31. Hocevar, L. A., & Fitzgerald, B. M. (2019). American society of anesthesiologists staging.

32. Faigel DO, Baron TH, Goldstein JL, et al. Guidelines for the use of deep sedation and anesthesia for GI endoscopy. *Gastrointest Endosc.* 2002;65(5):613-617.
33. Froehlich F, Schwizer W, Thorens J, et al. Conscious sedation for gastroscopy; Patient tolerance and cardiorespiratory parameters. *Gastroenterology.* 1995;108:697.
34. Geller E. From oral midazolam to propofol: A perspective. *Gastrointest Endosc.* 2005;61(2).
35. Goudra B, Nuzat A, Singh PM, et al. Association between Type of Sedation and the Adverse Events Associated with Gastrointestinal Endoscopy: An Analysis of 5 Years' Data from a Tertiary Center in the USA. *Clin Endosc.* 2016;50(2):161–169. doi:10.5946/ce.2016.019.
36. Harjai M, Alam S, Bhaskar P. Clinical Relevance of Mallampati Grading in Predicting Difficult Intubation in the Era of Various New Clinical Predictors. *Cureus.* 2021 Jul 14;13(7):e16396. doi: 10.7759/cureus.16396. PMID: 34408949; PMCID: PMC8362876.
37. Herman FN. Avoidance of sedation during total colonoscopy. *Dis Colon Rectum.* 1990;33:70.
38. Jeffrey L. Apfelbaum, Carin A. Hagberg, Richard T. Connis, Basem B. Abdelmalak, Madhulika Agarkar, Richard P. Dutton, John E. Fiadjoe, Robert Greif, P. Allan Klock, David Mercier, Sheila N. Myatra, Ellen P. O'Sullivan, William H. Rosenblatt, Massimiliano Sorbello, Avery Tung; 2022 American Society of Anesthesiologists Practice Guidelines for Management of the Difficult Airway. *Anesthesiology* 2022; 136:31–81 doi: <https://doi.org/10.1097/ALN.0000000000004002>
39. Jung M, Hofmann C, Kiesslich R, Brakertz A. Improved sedation in diagnostic and therapeutic ERCP: Propofol is an alternative to midazolam. *Endoscopy* 2000;32:233-238.
40. Koshy G, Nair S, Norkus EP, et al. Propofol versus midazolam and meperidine for conscious sedation in GI endoscopy. *Am J Gastroenterol.* 2000;95:1476-1479.
41. Lieber SR, Heller BJ, Howard CW, Sandler RS, Crockett S, Barritt AS 4th. Complications Associated With Anesthesia Services in Endoscopic Procedures Among Patients With Cirrhosis. *Hepatology.* 2020 Dec;72(6):2109-2118. doi: 10.1002/hep.31224. PMID: 32153048; PMCID: PMC7483314.
42. The Joint Commission. (2021, October 19). Sedation and Anesthesia - Understanding the Assessment Requirements. What are the assessment requirements organizations need to understand? <https://www.jointcommission.org/standards/standard-faqs/ambulatory/provision-of-care-treatment-and-services-pc/000001645/>
43. Mallampati SR, Gatt SP, Gugino LD, Desai SP, Waraksa B, Freiburger D, Liu PL. A clinical sign to predict difficult tracheal intubation: a prospective study. *Can Anaesth Soc J.* 1985 Jul;32(4):429-34. doi: 10.1007/BF03011357. PMID: 4027773.
44. Paspatis GA, Tribonias G, Manolaraki MM, et al. Deep sedation compared with moderate sedation in polyp detection during colonoscopy: a randomized controlled trial. *Colorectal Dis.* 2011; 13(6):e137-144.
45. Patterson KW, Casey PB, Murray JP, et al. Propofol sedation for outpatient upper gastrointestinal endoscopy: Comparison with midazolam. *Br J Anaesth* 1991;67:108-111
46. Qadeer MA, Vargo JJ, Khandwala F, et al. Propofol versus traditional sedative agents for gastrointestinal endoscopy: A meta-analysis. *Clin Gastroenterol Hepatol.* 2005;3(11):1049- 1056

47. Rex DK, Imperiale TF, Portish V. Patients willing to try colonoscopy without sedation: Associated clinical factors and results of a randomized controlled trial. *Gastrointest Endosc.* 1999;49(5):554-559
48. Rex DK, Overlay CA, Walker J. Registered nurse-administered propofol sedation for upper endoscopy and colonoscopy: Why? When? How? *Rev Gastroenterol Disord.* 2003; 3(2):70-80.
49. Ristikankare M, Hartikainen J, Heikkinen M, et al. Is routinely given conscious sedation of benefit during colonoscopy? *Gastrointest Endosc.* 1999;49:566.
50. Rosero EB. Monitored anesthesia care in adults. UpToDate.com. Last updated Dec 1, 2020. [https://www.uptodate.com/contents/monitored-anesthesia-care-in-adults?search=monitored%20anesthesia%20care&source=search\\_result&selectedTitle=1~131&usage\\_type=default&display\\_rank=1](https://www.uptodate.com/contents/monitored-anesthesia-care-in-adults?search=monitored%20anesthesia%20care&source=search_result&selectedTitle=1~131&usage_type=default&display_rank=1)
51. Samssoon GL, Young JR. Difficult tracheal intubation: a retrospective study. *Anaesthesia* 1987; 42:487.
52. Seow-Choen F, Leong AF, Tsang C. Selective sedation for colonoscopy. *Gastrointest Endosc.* 1994;40:661.
53. Sipe BW, Rex DK, Latinovich D. Propofol versus midazolam/meperidine for outpatient colonoscopy: Administration by nurses supervised by endoscopist. *Gastrointest Endosc.* 2002;55(7):815-825
54. Solomon SA, Kajla VK, Banerjee AK. Can the elderly tolerate endoscopy without sedation? *J R Coll Physicians Lond.* 1994;28:407
55. Tan CC, Freeman JG. Throat spray for upper gastrointestinal endoscopy is quite acceptable to patients. *Endoscopy.* 1996;28:177
56. Vargo JJ, Cohen LB, Rex DK, Kwo PY. Position statement: Nonanesthesiologist administration of propofol for GI endoscopy. *Am J Gastroenterol* 2009; 104(12):2886–2892.
57. Vargo JJ, Niklewski PJ, Williams JL, et al. Patient safety during sedation by anesthesia professionals during routine upper endoscopy and colonoscopy: an analysis of 1.38 million procedures. *Gastrointest Endosc.* 2017 Jan;85(1):101-108
58. Vargo JJ, Zuccaro G, Dumot JA, et al. Gastroenterologist-administered propofol versus meperidine and midazolam for advanced upper endoscopy: A prospective, randomized trial. *Gastroenterology.* 2002;123(1):8-16
59. Waring JP, Baron TH, Hirota WK, et al. Guidelines for conscious sedation and monitoring during gastrointestinal endoscopy. *Gastrointest Endosc.* 2003;58(3):317-322.
60. Wernli KJ, Brenner AT, Rutter CM, et al. Risks Associated With Anesthesia Services During Colonoscopy. *Gastroenterology* Volume 150, Issue 4, April 2016, Pages 888-894.

#### Clinical Guideline Revision / History Information

Original Date: 10/11/2017

Reviewed/Revised: 1/18/2018, 11/6/2018, 10/21/2019, 10/21/2020, 10/21/2021, 12/01/2021, 10/20/2022, 10/19/2023, 11/01/2024