

Balloon Ostial Dilation

Disclaimer

Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.

Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.

Balloon Ostial Dilation	1
Summary	1
Definitions	2
Medical Necessity Criteria for Clinical Review	3
Indication-Specific Criteria	3
Chronic Rhinosinusitis	3
Recurrent Acute Rhinosinusitis	3
Experimental, Investigational, or Unproven / Not Medically Necessary	4
Acute Rhinosinusitis	5
Chronic Rhinosinusitis in Patients 17 Years-of-Age and Younger	5
Applicable Billing Codes	5
References	7
Clinical Guideline Revision / History Information	10

Summary

The Plan considers surgical treatment for chronic rhinosinusitis medically necessary in patients who fail attempts at medical therapy such as antibiotics, intranasal steroids, systemic steroids, and/or saline irrigation. Traditional functional endoscopic sinus surgery (FESS) is a minimally invasive surgical technique that opens and drains the sinus air cells and sinus passages under direct visualization.

Balloon ostial dilation (BOD), also known as balloon sinuplasty or balloon catheter sinusotomy, is a newer minimally invasive endoscopic technique that has been proposed as an alternative to or in addition to FESS for the surgical management of chronic rhinosinusitis. This technique involves using a balloon to dilate the sinus ostia to improve sinus drainage. Potential benefits of BOD include outflow tract enlargement with preservation of normal anatomy, excellent mucosal sparing, minimal intraoperative bleeding, and less discomfort than traditional sinus surgery. BOD may be used as a standalone procedure, but it is most frequently used in combination with FESS, which is also known as a “hybrid” technique.

Studies have confirmed the safety and efficacy of balloon ostial dilation in patients who have failed medical therapy. Balloon dilation also has some advantages when compared to FESS in terms of recovery time and degree of postoperative pain.

Definitions

“Rhinosinusitis” is defined as an inflammatory condition of one or more of the paranasal sinuses and nasal passages. Rhinosinusitis can be further classified based on acuity (acute vs. chronic), anatomical location (involvement of the maxillary, ethmoid, frontal, or sphenoid sinuses), and severity (uncomplicated vs. complicated).

- “Acute rhinosinusitis (ARS)” is an inflammatory condition of one or more of the paranasal sinuses and nasal passages (“rhinosinusitis”) that lasts up to 4 weeks.
- “Recurrent acute rhinosinusitis (RARS)” is defined as at least 4 episodes of acute rhinosinusitis in a 1-year period. Patients are asymptomatic between episodes. Even though the summed duration of all episodes may be greater than 12 weeks, this is differentiated from chronic rhinosinusitis in that each single episode is still less than 4 weeks long.
- “Chronic rhinosinusitis (CRS)” is an inflammatory condition of one or more of the paranasal sinuses and nasal passages (“rhinosinusitis”) that persists for 12 weeks or longer.

“Functional endoscopic sinus surgery (FESS)” is a minimally invasive surgical technique to improve the drainage pathways of the paranasal sinuses. It is performed with an endoscope inserted into the nasal cavity to allow for direct visualization of sinus anatomy.

“Hybrid technique” is the combination of traditional endoscopic sinus surgery and balloon ostial dilation techniques during a single procedure.

Medical Necessity Criteria for Clinical Review

Indication-Specific Criteria

Chronic Rhinosinusitis

The Plan considers balloon ostial dilation medically necessary when ALL of the following criteria are met:

1. Patient 18 years of age and older; *and*
2. Documented persistent chronic rhinosinusitis for at least 12 weeks with two or more of the following signs or symptoms:
 - a. Mucopurulent drainage (not clear drainage); *and/or*
 - b. Nasal obstruction (congestion); *and/or*
 - c. Facial pain, pressure, or fullness (differentiated from headaches); *and/or*
 - d. Decreased sense of smell; *and*
3. Documented failure of medical therapy despite treatment with:
 - a. A trial of at least TWO different oral antibiotics (e.g., amoxicillin-clavulanate, cefpodoxime, clindamycin, doxycycline, levofloxacin, metronidazole, etc.) unless there is a specific contraindication, severe adverse reaction, or cause of rhinosinusitis is a viral illness; *and*
 - b. A trial of at least TWO of the following topical intranasal therapies for a minimum of 4+ weeks each:
 - i. Intranasal steroids; *or*
 - ii. Saline irrigations; *or*
 - iii. Antihistamine spray.
4. Allergic rhinitis has been adequately ruled out or treated appropriately with medications and/or documented lifestyle changes; *and*
5. Confirmation of chronic rhinosinusitis by CT scan radiology report as demonstrated by the presence of at least ONE of the following:
 - a. Mucosal thickening; *or*
 - b. Bony remodeling or thickening; *or*
 - c. Sinus opacification; *or*
 - d. Obstruction of the ostiomeatal complex.
6. Dilation is limited to the maxillary, frontal, or sphenoid sinuses; *and*
7. If a balloon ostial dilation is performed in conjunction with a functional endoscopic sinus surgery (FESS) in the same sinus cavity for a member without nasal polyps, it must meet above Chronic Rhinosinusitis criteria and MCG (A-0185) Functional Endoscopic Sinus Surgery (FESS) criteria.

Recurrent Acute Rhinosinusitis

The Plan considers balloon ostial dilation medically necessary when ALL of the following criteria are met:

1. Patient 18 years of age and older; *and*
2. Documented recurrent acute rhinosinusitis of at least 4 episodes in a 12-month rolling period with asymptomatic periods in between episodes; *and*
3. Documented recurrence of symptoms despite optimal prior medical therapy with:

- a. Oral antibiotic therapy (e.g., amoxicillin, amoxicillin-clavulanate, etc.) unless specific contraindication, severe adverse reaction, or cause of rhinosinusitis is viral illness; *and*
- b. A trial of at least TWO of the following topical intranasal therapies for a minimum of 4+ weeks each:
 - i. Intranasal steroids; *or*
 - ii. Saline irrigations; *or*
 - iii. Antihistamine spray.
4. Allergic rhinitis has been adequately ruled out or treated appropriately with medications and/or documented lifestyle changes; *and*
5. Confirmation of acute rhinosinusitis that would be reasonably expected to improve after surgical treatment, based on CT imaging radiology report results during an acute episode; *and*
6. Dilation is limited to the maxillary, frontal, or sphenoid sinuses; *and*
7. If a balloon ostial dilation is performed in conjunction with a functional endoscopic sinus surgery (FESS) in the same sinus cavity for a member without nasal polyps, it must meet above Recurrent Acute Rhinosinusitis criteria and MCG (A-0185) Functional Endoscopic Sinus Surgery (FESS) criteria.

Experimental, Investigational, or Unproven / Not Medically Necessary

The Plan considers balloon ostial dilation experimental, investigational, or unproven for all other conditions not listed in this guideline, including but not limited to the following, as its safety and efficacy have not been clearly established in the published clinical literature:

1. Nasal polyps
2. Tumors
3. Acute rhinosinusitis
4. Aspirin sensitivity
5. Sinusitis secondary to autoimmune or connective tissue disorders
6. Sinusitis secondary to ciliary dysfunction (e.g., cystic fibrosis)
7. Isolated ethmoid sinus disease
8. Sleep apnea without chronic sinusitis symptoms and imaging findings
9. Headache without chronic sinusitis symptoms and imaging findings

The Plan considers balloon ostial dilation experimental, investigational, or unproven in children 17 years of age and younger as there is insufficient evidence to determine its effectiveness in this population.

The Plan considers the use of self-expanding absorptive sinus ostial dilation devices experimental, investigational, or unproven as the safety and efficacy have not been clearly established in the published clinical literature.

Balloon sinus ostial dilation used adjunctively during functional endoscopic sinus surgery (FESS) in the same sinus cavity is considered integral to the primary procedure and not separately reimbursable.

Similarly, when balloon dilation and sinus tissue removal are performed as part of the same surgery, only one code will be considered reimbursable.

Acute Rhinosinusitis

There are several case reports in the literature (Roland et al. 2015; Zhao et al. 2016; Hopkins et al. 2009) and one small retrospective case series (Wittkopf et al. 2009) addressing the use of balloon dilation in cases of acute sinusitis, with or without associated complication. The results of these reports are interesting but do not provide objective data, statistics, or long-term follow up to allow a conclusion that is generalizable to the standard acute rhinosinusitis patient. Further studies that provide clinical data based on randomized trials comparing medical therapy to balloon ostial dilation for acute rhinosinusitis and traditional FESS to balloon dilation for complicated acute disease are needed to make any scientific conclusions.

Chronic Rhinosinusitis in Patients 17 Years-of-Age and Younger

A consensus statement from the American Academy of Otolaryngology-Head and Neck Surgery (AAO-HNS) on the use of balloon sinuplasty for the treatment of pediatric chronic rhinosinusitis concluded that the effectiveness cannot be determined based on current evidence. The current literature includes a nonrandomized prospective study (Ramadan et al. 2011), a retrospective case control study (Wang et al. 2015), a retrospective cohort study (Thottam et al. 2016), and several case reports that have suggested that balloon catheter sinuplasty is safe and might be as effective as FESS in children with chronic or acute rhinosinusitis, but larger, prospective, randomized control trials with long term follow up are needed to determine the true efficacy in this population.

Applicable Billing Codes

Table 1	
CPT/HCPCS codes considered medically necessary if criteria are met:	
<i>Code</i>	<i>Description</i>
31295	Nasal/sinus endoscopy, surgical, with dilation (eg, balloon dilation); maxillary sinus ostium, transnasal or via canine fossa
31296	Nasal/sinus endoscopy, surgical, with dilation (eg, balloon dilation); frontal sinus ostium
31297	Nasal/sinus endoscopy, surgical, with dilation (eg, balloon dilation); sphenoid sinus ostium
31298	Nasal/sinus endoscopy, surgical, with dilation (eg, balloon dilation); frontal and sphenoid sinus ostia
C1726	Catheter, balloon dilatation, nonvascular

Table 2	
CPT/HCPCS codes <u>not considered medically necessary</u> for indications listed in this guideline:	
<i>Code</i>	<i>Description</i>
31299	Unlisted procedure, accessory sinuses

Table 3	
ICD-10 codes considered experimental, investigational or unproven with Table 1 codes:	
<i>Code</i>	<i>Description</i>
C31.0 - C31.9	Malignant neoplasm of accessory sinuses
D14.0	Benign neoplasm of middle ear, nasal cavity and accessory sinuses
E84.0 - E84.9	Cystic fibrosis • <u>Due to the broad nature of these codes, specific exclusions are indicated:</u> • When billed for sinusitis secondary to ciliary dysfunction (e.g., cystic fibrosis), it is considered experimental, investigational, or unproven
G44.001 - G44.89	Other headache syndromes
G47.30 - G47.39	Sleep apnea
J01.00	Acute maxillary sinusitis, unspecified
J01.10	Acute frontal sinusitis, unspecified
J01.20	Acute ethmoidal sinusitis, unspecified
J01.21	Acute recurrent ethmoidal sinusitis
J01.30	Acute sphenoidal sinusitis, unspecified
J01.40	Acute pansinusitis, unspecified
J01.80	Other acute sinusitis
J01.90	Acute sinusitis, unspecified
J32.2	Chronic ethmoidal sinusitis
J33.0 - J33.9	Nasal polyps
M30.0 - M36.8	Systemic connective tissue disorders

Table 3	
ICD-10 codes considered experimental, investigational or unproven with Table 1 codes:	
<i>Code</i>	<i>Description</i>
R51.0 - R51.9	Headache
T39.015A - T39.015S	Adverse effect of aspirin
Z88.6	Allergy status to analgesic agent • <u>Due to the broad nature of this code, specific exclusions are indicated:</u> • When billed for aspirin sensitivity, it is considered experimental, investigational, or unproven

References

1. Ahmed, J., Pal, S., Hopkins, C., & Jayaraj, S. (2011). Functional endoscopic balloon dilation of sinus ostia for chronic rhinosinusitis. *Cochrane Database of Systematic Reviews*, (7).
2. American Academy of Otolaryngology-Head and Neck Surgery (AAO-HNS). Statement on Reimbursement of Balloon Sinus Ostial Dilation. September 2014.
3. American Academy of Otolaryngology-Head and Neck Surgery (AAO-HNS). Position Statement: Dilation of sinuses, any method (e.g., balloon, etc.). March 2017.
4. American Rhinologic Society. Ostial balloon dilation position statement. March 14, 2017. Available at: <https://www.american-rhinologic.org/position-statements>. Accessed on May 23, 2019.
5. Achar P, Duvvi S, Kumar BN. Endoscopic dilatation sinus surgery (FEDS) versus functional endoscopic sinus surgery (FESS) for treatment of chronic rhinosinusitis: a pilot study. *Acta Otorhinolaryngol Ital*. 2012; 32(5):314-319.
6. Albritton FD 4th, Casiano RR, Sillers MJ. Feasibility of in-office endoscopic sinus surgery with balloon sinus dilation. *Am J Rhinol Allergy*. 2012; 26(3):243-248.
7. Bikhazi N, Light J, Truitt T, et al.; REMODEL Study Investigators. Standalone balloon dilation versus sinus surgery for chronic rhinosinusitis: a prospective, multicenter, randomized, controlled trial with 1-year follow-up. *Am J Rhinol Allergy*. 2014; 28(4):323-329.
8. Brietzke SE, Shin JJ, Choi S, et al. Clinical consensus statement: pediatric chronic rhinosinusitis. *Otolaryngol Head Neck Surg*. 2014 Oct;151(4):542-53.
9. Brodner D, Nachlas N, Mock P, et al. Safety and outcomes following hybrid balloon and balloon-only procedures using a multifunction, multisinus balloon dilation tool. *Int Forum Allergy Rhinol*. 2013; 3(8):652-658.
10. Brook, I. (April 2022). *Microbiology and antibiotic management of chronic rhinosinusitis*. UpToDate Retrieved from

<https://www.uptodate.com/contents/microbiology-and-antibiotic-management-of-chronic-rhino-sinusitis>

11. Chandra RK, Kern RC, Cutler JL, et al. REMODEL larger cohort with long-term outcomes and meta-analysis of standalone balloon dilation studies. *Laryngoscope*. 2016; 126(1):44-50.
12. Cingi, C., Muluk, N. B., & Lee, J. T. (2019). Current indications for balloon sinuplasty. *Current Opinion in Otolaryngology & Head and Neck Surgery*, 27(1), 7-13.
13. Bikhazi N, Light J, Truitt T, et.al. Standalone balloon dilation versus sinus surgery for chronic rhinosinusitis: A prospective, multicenter, randomized, controlled trial with 1-year follow-up. *Am J Rhinol Allergy* 2014; 28(4): 323-329.
14. Gould J, Alexander I, Tomkin E, Brodner D. In-office, multisinus balloon dilation: 1-Year outcomes from a prospective, multicenter, open label trial. *Am J Rhinol Allergy*. 2014; 28(2):156-163.
15. Hathorn IF, Pace-Asciak P, Habib AR, et al. Randomized controlled trial: hybrid technique using balloon dilation of the frontal sinus drainage pathway. *Int Forum Allergy Rhinol*. 2015; 5(2):167-173.
16. Hamilos DL, Holbrook EH. (2022). Chronic rhinosinusitis: Management. UpToDate. Retrieved from <https://www.uptodate.com/contents/chronic-rhinosinusitis-management>
17. Hayes, Inc. Health Technology Assessment. Balloon Sinuplasty for Treatment of Chronic Rhinosinusitis in Adult Patients. Lansdale, PA: Hayes, Inc.; January 4, 2021.
18. Higgins Jr, T. S., Öcal, B., Adams, R., & Wu, A. W. (2020). In-Office balloon Sinus ostial dilation with concurrent antiplatelet and anticoagulant therapy for chronic rhinosinusitis without nasal polyps. *Annals of Otolaryngology, Rhinology & Laryngology*, 129(3), 280-286.
19. Hoying, D., Kabalan, M. J., Kaelber, D. C., & Sindwani, R. (2026). Comparison of long-term revision rates and health care utilization between standalone balloon sinus dilation and endoscopic sinus surgery in chronic rhinosinusitis patients without nasal polyps: Is the balloon value proposition overinflated?. *American Journal of Rhinology & Allergy*, 40(1), 31–37. <https://doi.org/10.1177/19458924251377259>
20. Hubbell, R. D., Toivonen, J., Kawai, K., Kim, H. J., Nieman, C. L., Ward, B. K., & Poe, D. S. (2023). Patulous Eustachian Tube Dysfunction Symptoms Following Balloon Dilation. *The Laryngoscope*, 133(11), 3152-3157.
21. Huisman, J. M. L., Verdam, F. J., Stegeman, I., & de Ru, J. A. (2018). Treatment of Eustachian tube dysfunction with balloon dilation: a systematic review. *The Laryngoscope*, 128(1), 237-247.
22. Karanfilov B, Silvers S, Pasha R, et al. Office-based balloon sinus dilation: a prospective, multicenter study of 203 patients. *Int Forum Allergy Rhinol*. 2013; 3(5):404-411.
23. Levine AB, Truitt TT, Schwartz M, Atkins J. In-office stand-alone balloon dilation of maxillary sinus ostia and ethmoid infundibula in adults with chronic or recurrent acute rhinosinusitis: a prospective, multi-institutional study with 1-year follow-up. *Ann Otol Rhinol Laryngol*. 2013; 122(11):665-671.
24. Levy JM, Marino MJ, McCoul ED. Paranasal sinus balloon catheter dilation for treatment of chronic rhinosinusitis: a systematic review and meta-analysis. *Otolaryngol Head Neck Surg*. 2016; 154(1):33-40.

25. Lofgren, D. H., & Shermetaro, C. (2023, July 24). *Balloon sinuplasty*. StatPearls - NCBI Bookshelf. <https://www.ncbi.nlm.nih.gov/books/NBK546671/>
26. Marzetti A, Tedaldi M, Passali FM. The role of balloon sinuplasty in the treatment of sinus headache. *Otolaryngol Pol*. 2014; 68(1):15-19.
27. Peters AT, Spector S, Hsu J, et al. Diagnosis and management of rhinosinusitis: a practice parameter update. *Ann Allergy Asthma Immunol*. 2014 Oct;113(4):347-85.
28. Piccirillo JF, Merritt MG Jr, Richards ML. Psychometric and clinimetric validity of the 20-Item Sino-Nasal Outcome Test (SNOT-20). *Otolaryngol Head Neck Surg*. 2002; 126(1):41-47.
29. Piccirillo, J. F., Payne, S. C., Rosenfeld, R. M., Baroody, F. M., Batra, P. S., DelGaudio, J. M., ... & Corrigan, M. D. (2018). Clinical consensus statement: balloon dilation of the sinuses. *Otolaryngology–Head and Neck Surgery*, 158(2), 203-214.
30. Roland LT, Wineland AM, Leonard DS. Balloon frontal sinuplasty for intracranial abscess in a pediatric acute sinusitis patient. *Int J Pediatr Otorhinolaryngol*. 2015 Mar;79(3):432-4.
31. Rosenfeld RM, Piccirillo JF, Chandrasekhar SS, et al. Clinical practice guideline (update): adult sinusitis. *Otolaryngol Head Neck Surg*. 2015;152(2 Suppl):S1-S39.
32. Sikand A, Silvers SL, Pasha R, et al.; ORIOS 2 Study Investigators. Office-based balloon sinus dilation: 1-year follow-up of a prospective, multicenter study. *Ann Otol Rhinol Laryngol*. 2015; 124(8):630-637.
33. Sinha, P., Tharakan, T., Payne, S., & Piccirillo, J. F. (2023). Balloon sinus dilation versus functional endoscopic sinus surgery for chronic rhinosinusitis: Systematic review and meta-analysis. *Annals of Otolaryngology, Rhinology, and Laryngology*, 132(5), 578–588. <https://doi.org/10.1177/00034894221104939>
34. Soler ZM, Rosenbloom JS, Skarada D, Gutman M, Hoy MJ, Nguyen SA. Prospective, multicenter evaluation of balloon sinus dilation for treatment of pediatric chronic rhinosinusitis. *Int Forum Allergy Rhinol*. 2016 Nov 26.
35. Stolovitzky, J. P., Mehendale, N., Matheny, K. E., Brown, W. J., Rieder, A. A., Liepert, D. R., Tseng, E., & Gould, A. (2018). Medical therapy versus balloon sinus dilation in adults with chronic rhinosinusitis (MERLOT): 12-month follow-up. *American Journal of Rhinology & Allergy*, 32(4), 294–302. <https://doi.org/10.1177/1945892418773623>
36. Thottam PJ, Metz CM, Kieu MC, Dworkin J, Jagini J, Bangiyev JN, Mehta D. Functional Endoscopic Sinus Surgery Versus Balloon Sinuplasty with Ethmoidectomy: A 2-year Analysis in Pediatric Chronic Rhinosinusitis. *Indian J Otolaryngol Head Neck Surg*. 2016 Sep;68(3):300-6.
37. Tomazic PV, Stammberger H, Braun H, et al. Feasibility of balloon sinuplasty in patients with chronic rhinosinusitis: the Graz experience. *Rhinology*. 2013; 51(2):120-127.
38. Tucci D, McCoul E, Rosenfeld R, et al. Clinical Consensus Statement: Balloon Dilation of the Eustachian Tube. American Academy of Otolaryngology–Head and Neck Surgery Foundation 2019. April 15, 2019:1-12.
39. Zhao Y, Chen K, Wang Z. Sinus Balloon Dilation as Treatment for Acute Sphenoid Sinusitis with Impaired Vision for a Child. *Case Rep Med*. 2016;2016:5209243. doi: 10.1155/2016/5209243. Epub 2016 Feb 24.

Clinical Guideline Revision / History Information

Original Date: 8/1/2017

Reviewed/Revised: 1/18/2018, 7/31/2018, 7/23/2019, 07/21/2020, 8/4/2021, 12/01/2021, 07/26/2022, 07/19/2023, 07/29/2024, 11/01/2025, 11/02/2026