

Roctavian (valoctocogene roxaparvovec-rvox)

Disclaimer

Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.

Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.

Summary

Hemophilia A is a rare, X-linked genetic bleeding disorder caused by coagulation factor VIII deficiency. Based on the level of clotting factor (factor VIII) activity detected in the blood, hemophilia A severity is classified into the following categories:

1. Severe disease - Factor VIII activity less than 1% of normal (less than 1 unit/dL)
2. Moderate disease - Factor VIII activity between 1% to 5% of normal (1 to 5 units/dL)
3. Mild disease - Factor VIII activity more than 5% but less than 40% of normal (more than 5 but less than 40 units/dL)

Patients with severe hemophilia A are at high risk for spontaneous bleeding episodes, particularly into joints and muscles, leading to chronic pain, joint damage, and disability. The current standard of care is routine intravenous prophylactic infusions of factor VIII concentrate to prevent bleeding events.

However, breakthrough bleeding still occurs due to the short 8-12 hour half-life of factor VIII products.

Roctavian (valoctocogene roxaparvovec-rvox) is the first gene therapy approved for adults with severe hemophilia A. It delivers a functional F8 gene via an AAV5 viral vector to enable endogenous factor VIII production. It is indicated for the treatment of adults with severe hemophilia A (congenital factor VIII deficiency with factor VIII activity < 1 IU/dL) without antibodies to adeno-associated virus serotype 5 (AAV5) detected by an FDA-approved test

- It is administered as a one-time intravenous infusion.
- It provides an alternative to intravenous factor VIII prophylaxis, but is not a cure for hemophilia A.
- In the phase 3 trial, Roctavian (valoctocogene roxaparvovec-rvox) reduced annualized bleeding rates by 52% over 3 years. However, factor VIII activity wanes over time.

Definitions

“AAV5 Viral Vector” is a tool used in gene therapy to deliver a functional F8 gene to cells. AAV5 is a specific type of adeno-associated virus.

“Annualized Bleeding Rates” refers to the expected number of bleeding episodes within a year.

“Antibodies to AAV5” are proteins produced by the immune system that recognize and neutralize the AAV5 virus, potentially impacting the efficacy of Roctavian.

“Bethesda Assay” is a laboratory test used to measure the level of factor VIII inhibitors in the blood.

“Factor VIII Inhibitors” are proteins produced by the immune system that neutralize factor VIII, making treatment less effective.

“Factor VIII” is a protein involved in blood clotting. Its deficiency leads to Hemophilia A.

“FDA” refers to the United States Food and Drug Administration, responsible for approving drugs and medical treatments.

“Hemophilia A” is a rare, X-linked genetic bleeding disorder resulting from a deficiency in coagulation factor VIII.

“Prophylactic Infusions” refers to preventive treatment given to avoid the onset of a disease or condition, in this case, bleeding episodes.

“**Thrombocytopenia**” is a condition characterized by low platelet count, leading to increased risk of bleeding.

“**Vector Genomes**” are complete sets of viral DNA. In this context, it's used to measure the dosage of Roctavian.

“**Von Willebrand Disease**” is a genetic disorder caused by missing or defective von Willebrand factor (VWF), a clotting protein.

“**X-linked**” refers to a gene carried on the X chromosome. In the context of hemophilia, it primarily affects males since they have only one X chromosome.

Medical Necessity Criteria for Authorization

The Plan considers **Roctavian (valoctocogene roxaparvovec-rvox)** medically necessary when **ALL** of the following criteria are met:

1. Prescribed by or in consultation with a physician experienced in the treatment of hemophilia and/or bleeding disorders (e.g., hematologist); **AND**
 - a. **IF** there is documentation indicating any radiological liver irregularities or liver function test anomalies (ALT, AST, GGT, ALP or total bilirubin $> 1.25 \times \text{ULN}$ or $\text{INR} \geq 1.4$), a consultation with a hepatologist is required.
2. The member is 18 years of age or older; **AND**
3. The member has a diagnosis of severe hemophilia A, defined as congenital factor VIII deficiency with factor VIII activity level less than or equal to (\leq) 1 IU/dL; **AND**
4. Documentation of **ALL** of the following :
 - a. baseline testing for **BOTH** of the following:
 - i. factor VIII inhibitor titer; **and**
 - ii. preexisting antibodies to adeno-associated virus serotype 5 (AAV5)^{2/}; **and**
 - b. at least 6 months prophylactic use of factor VIII before considering Roctavian; **and**
 - c. liver health assessments, which include:
 - i. Liver function tests, i.e., ALT, AST, GGT, ALP, total bilirubin, and INR; **and**
 - ii. Liver condition evaluations through ultrasound, elastography, or laboratory assessments for liver fibrosis; **and**
 - d. A comprehensive review of the member’s concurrent therapy considerations, inclusive of:

- i. alcohol consumption **AND** the member has received guidance to refrain from alcohol consumption for a minimum of one year after Roctavian administration and is encouraged to moderate their alcohol intake thereafter; **and**
- ii. herbal products and dietary supplements; **and**
- iii. medications that might induce hepatotoxicity, reduce factor VIII activity, or alter plasma corticosteroid levels which can influence liver enzymes or factor VIII activity; **and**
- iv. member's capability to undergo prolonged corticosteroids and/or other immunosuppressive treatments; **AND**

⁷*Information on approved tests is available at fda.gov/CompanionDiagnostics.*

5. The member does **NOT** have **ANY** of the following:
 - a. Active infections, either acute (such as acute respiratory infections or acute hepatitis) or uncontrolled chronic (such as chronic active hepatitis B); **and/or**
 - b. Known hypersensitivity to mannitol; **and/or**
 - c. Presence of antibodies to AAV5 capsid, as confirmed by an FDA-approved/cleared or CLIA-compliant test⁷, such as ARUP Laboratories' AAV5 DetectCDx™; **and/or**
 - d. presence of factor VIII inhibitors (i.e., results from a Bethesda assay or Bethesda assay with Nijmegen modification of less than 0.6 Bethesda Units (BU)); **and/or**
 - e. Stage 3 or 4 liver fibrosis, or cirrhosis; **AND**
6. Authorization is for no more than:
 - a. 6×10^{13} vector genomes per kilogram (vg/kg) body weight, administered as a single intravenous infusion; **or**
 - b. one dose of Roctavian (valoctocogene roxaparvovec-rvox) per lifetime.

If the above criteria are met, one dose of Roctavian (valoctocogene roxaparvovec-rvox) will be approved for the one-time treatment.

Experimental or Investigational / Not Medically Necessary

Roctavian (valoctocogene roxaparvovec-rvox) for any other indication or use is considered not medically necessary by the Plan, as it is deemed to be experimental, investigational, or unproven. Non-covered indications include, but are not limited to, the following:

1. hemophilia A (congenital factor VIII deficiency):
 - a. Retreatment after receiving one prior dose

- b. dosage exceeding the FDA-approved dose of 6×10^{13} vector genomes per kilogram (vg/kg) body weight
- c. Use in females
- d. Use in patients with pre-existing antibodies to adeno-associated virus serotype 5 (AAV5)
- e. Use in patients with current or prior factor VIII inhibitors
- f. Use in pediatric patients
- g. Mild or moderate hemophilia A
 - i. Mild: Factor VIII level >5% to <40%
 - ii. Moderate: Factor VIII level 1% to 5%
- 2. Treatment of other coagulation factor deficiencies (e.g. factors IX, XI, XII)
- 3. Treatment of thrombocytopenia or platelet dysfunction
- 4. Treatment of vitamin K deficiency
- 5. Treatment of von Willebrand disease

Applicable Billing Codes (HCPCS/CPT Codes)

Service(s) name	
CPT/HCPCS Codes considered medically necessary if criteria are met:	
<i>Code</i>	<i>Description</i>
96365	Intravenous infusion, for therapy, prophylaxis, or diagnosis (specify substance or drug); initial, up to 1 hour
96366	Intravenous infusion, for therapy, prophylaxis, or diagnosis (specify substance or drug); each additional hour (List separately in addition to code for primary procedure)
99221	Initial hospital inpatient or observation care, per day, for the evaluation and management of a patient, which requires a medically appropriate history and/or examination and straightforward or low level medical decision making. When using total time on the date of the encounter for code selection, 40 minutes must be met or exceeded.
99222	Initial hospital inpatient or observation care, per day, for the evaluation and management of a patient, which requires a medically appropriate history and/or examination and moderate level of medical decision making. When using total time on the date of the encounter for code selection, 55 minutes must be met or exceeded.

99223	Initial hospital inpatient or observation care, per day, for the evaluation and management of a patient, which requires a medically appropriate history and/or examination and high level of medical decision making. When using total time on the date of the encounter for code selection, 75 minutes must be met or exceeded.
99234	Hospital inpatient or observation care, for the evaluation and management of a patient including admission and discharge on the same date, which requires a medically appropriate history and/or examination and straightforward or low level of medical decision making. When using total time on the date of the encounter for code selection, 45 minutes must be met or exceeded.
C9399	Unclassified drugs or biologicals
E0780	Ambulatory infusion pump, mechanical, reusable, for infusion less than 8 hours
J3490	Unclassified drugs
J3590	Unclassified biologics
ICD-10 codes considered medically necessary if criteria are met:	
<i>Code</i>	<i>Description</i>
D66	Hereditary factor VIII deficiency

References

1. Guidelines for the Management of Hemophilia. 3rd Edition. World Federation of Hemophilia 2020. Available at: <https://www1.wfh.org/publications/files/pdf-1863.pdf>.
2. Mahlangu J, Kaczmarek R, von Drygalski A, et al. Two-year outcomes of valoctocogene roxaparovec therapy for hemophilia A. *N Engl J Med*. 2023;388(8):694-705. doi:10.1056/NEJMoa2211075
3. Medical and Scientific Advisory Council (MASAC), National Hemophilia Foundation. MASAC recommendation concerning prophylaxis for hemophilia A and B with and without inhibitors (revised April 27, 2022). MASAC recommendation #267. Available at: https://www.hemophilia.org/sites/default/files/document/files/267_Prophylaxis.pdf
4. Medical and Scientific Advisory Council (MASAC), National Hemophilia Foundation. MASAC recommendations concerning products licensed for the treatment of hemophilia and other bleeding disorders (revised March 2022). MASAC recommendation #272. Available at: https://www.hemophilia.org/sites/default/files/document/files/272_Treatment.pdf
5. Ozelo MC, Mahlangu J, Pasi KJ, et al. Valoctocogene roxaparovec gene therapy for hemophilia A. *N Engl J Med*. 2022;386(11):1013-1025. doi:10.1056/NEJMoa2113708
6. Pasi KJ, Rangarajan S, Mitchell N, et al. Multiyear follow-up of AAV5-hFVIII-SQ gene therapy for hemophilia A. *N Engl J Med*. 2020;382(1):29-40. doi:10.1056/NEJMoa1908490
7. Rangarajan S, Walsh L, Lester W, et al. AAV5-Factor VIII gene transfer in severe hemophilia A. *N Engl J Med*. 2017;377(26):2519-2530. doi:10.1056/NEJMoa1708483

8. Roctavian (valoctocogene roxaparvovec) [prescribing information]. Novato, CA: BioMarin Pharmaceuticals Inc; June 2023.

Clinical Guideline Revision / History Information

Original Date: 9/21/2023

Reviewed/Revised: