

## Ivermectin 1% Topical Cream

### Disclaimer

*Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.*

*Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.*

### Summary

Rosacea is a chronic inflammatory skin condition primarily affecting the face. It is characterized by redness, swelling, telangiectasia (dilated blood vessels that appear on the surface of the skin), inflammatory papules and pustules, and rhinophyma (characterized by an enlarged, red nose). The exact cause is unknown, but factors such as genetics, immune system dysfunction, and microbial imbalances may contribute. Rosacea is managed by non-pharmacological (e.g., avoiding triggers of flushing, gentle skin care, sun-protection), laser and light therapy, and pharmacological therapy such as topical brimonidine, topical antimicrobials, topical azelaic acid, topical oxymetazoline, and oral antibiotics. The first-line treatment for mild-to-moderate disease, including inflammatory papules and pustules, include topical agents such as metronidazole, azelaic acid gel, topical ivermectin, sulfacetamide-sulfur, and erythromycin. Topical ivermectin can be used to improve symptoms of stinging, burning, and eyelid inflammation.

Ivermectin 1% cream (Soolantra) is a semi-synthetic derivative with both anti-inflammatory and antiparasitic properties. It is FDA-approved for the treatment of inflammatory lesions of rosacea. Ivermectin cream is typically used after failure of, or in conjunction with, other topical therapies and may be an alternative to systemic treatments in some patients.

## Definitions

"Inflammatory lesions of rosacea" refers to papules and pustules on the face associated with rosacea.

"Papulopustular rosacea" is a subtype of rosacea characterized by papules (red bumps) and pustules (pus-filled bumps) in addition to persistent central facial erythema.

"Persistent erythema" is lasting redness of the facial skin that is characteristic of rosacea.

"Rosacea" is a skin condition primarily affecting the face and can present as redness, swelling, inflammation that includes papules and pustules, dilated blood vessels that appear on the surface of the skin, and rhinophyma (characterized by an enlarged, red nose).

"Telangiectasia" refers to visible dilated blood vessels near the surface of the skin, commonly seen in rosacea.

## Medical Necessity Criteria for Initial Authorization

The Plan considers Ivermectin 1% Topical Cream medically necessary when ALL of the following criteria are met:

1. The member is 18 years of age or older; *AND*
2. The member has a diagnosis of rosacea with inflammatory lesions (papules and pustules); *AND*
3. The member is unable to use, or has tried and failed at least TWO (2) of the following:
  - a. Metronidazole 0.75% or 1%; *and/or*
  - b. Azelaic acid 15%<sup>§</sup>; *and/or*  
*§NOTE: Prior authorization may be required.*
  - c. Oral antibiotics (e.g., doxycycline, minocycline).

If the above prior authorization criteria are met, the requested product will be authorized for up to 12-months.

## Medical Necessity Criteria for Reauthorization

Reauthorization for up to 12 months will be granted if the member has recent (within the last 3 months) clinical chart documentation indicating the member is responding positively to therapy as evidenced by reduction in inflammatory lesions.

## Experimental or Investigational / Not Medically Necessary

Ivermectin 1% Topical Cream for any other indication or use is considered not medically necessary by the Plan, as it is deemed to be experimental, investigational, or unproven. Non-covered indications include, but are not limited to, the following:

- Treatment of acne vulgaris. There is not enough high quality evidence to support the safety and efficacy of ivermectin 1% topical cream for the management of acne vulgaris.
- Treatment of flushing or persistent erythema associated with rosacea without inflammatory lesions. Ivermectin 1% topical cream is only approved for those with inflammatory lesions (papules and pustules). There is no high quality evidence to support the safety and efficacy of ivermectin 1% topical cream for the management of flushing or persistent erythema associated with rosacea without inflammatory lesions.
- Treatment of perioral dermatitis. Only low quality evidence (case reports, case series) have supported the use of ivermectin 1% topical cream for perioral dermatitis. There is not enough evidence to support the safety and efficacy of ivermectin 1% topical cream for the management of perioral dermatitis.
- Treatment of rhinophyma. There is not enough high quality evidence to support the safety and efficacy of ivermectin 1% topical cream for the management of rhinophyma.
- Treatment of seborrheic dermatitis. There is no high quality evidence to support the safety and efficacy of ivermectin 1% topical cream for the management of seborrheic dermatitis.

## References

1. Asai Y, Tan J, Baibergenova A, Barankin B, Cochrane CL, Humphrey S, Lynde CW, Marcoux D, Poulin Y, Rivers JK, Sapijaszko M, Sibbald RG, Toole J, Ulmer M, Zip C. Canadian Clinical Practice Guidelines for Rosacea. *J Cutan Med Surg*. 2016 Sep;20(5):432-45. doi: 10.1177/1203475416650427. Epub 2016 May 17. Erratum in: *J Cutan Med Surg*. 2021 Jul-Aug;25(4):466. doi: 10.1177/1203475417715265. PMID: 27207355.
2. Barańska-Rybak W, Kowalska-Olędzka E. New indications for topical ivermectin 1% cream: a case series study. *Postepy Dermatol Alergol*. 2019 Feb;36(1):58-62. doi: 10.5114/ada.2019.82825. Epub 2019 Feb 22.
3. Del Rosso JQ, Tangheiti E, Webster G, Stein Gold L, Thiboutot D, Gallo RL. Update on the Management of Rosacea from the American Acne & Rosacea Society (AARS). *J Clin Aesthet Dermatol*. 2020 Jun;13(6 Suppl):S17-S24. Epub 2020 Jun 1. PMID: 33282106; PMCID: PMC7710291.
4. Feaster B, Cline A, Feldman SR, Taylor S. Clinical effectiveness of novel rosacea therapies. *Curr Opin Pharmacol*. 2019 Jun;46:14-18. doi: 10.1016/j.coph.2018.12.001. Epub 2019 Jan 11. PMID: 30639950.
5. Hampton PJ, Berth-Jones J, Duarte Williamson CE, Hay R, Leslie TA, Porter I, Rauz S, Seukeran D, Winn RT, Hashme M, Exton LS, Mohd Mustapa MF, Manounah L; British Association of Dermatologists' Clinical Standards Unit. British Association of Dermatologists guidelines for the management of people with rosacea 2021. *Br J Dermatol*. 2021 Oct;185(4):725-735. doi: 10.1111/bjd.20485. Epub 2021 Jul 5. PMID: 33993465.
6. Reynolds RV, Yeung H, Cheng CE, Cook-Bolden F, Desai SR, Druby KM, Freeman EE, Keri JE, Stein Gold LF, Tan JKL, Tollefson MM, Weiss JS, Wu PA, Zaenglein AL, Han JM, Barbieri JS. Guidelines of care for the management of acne vulgaris. *J Am Acad Dermatol*. 2024 May;90(5):1006.e1-1006.e30. doi: 10.1016/j.jaad.2023.12.017. Epub 2024 Jan 30. PMID: 38300170.
7. Schaller M, Almeida LMC, Bewley A, et al. Recommendations for rosacea diagnosis, classification and management: update from the global ROSacea COnsensus 2019 panel. *Br J Dermatol*. 2020 May;182(5):1269-1276. doi: 10.1111/bjd.18420. Epub 2019 Oct 16.

8. Schaller M, Dirschka T, Lonne-Rahm SB, et al. The Importance of Assessing Burning and Stinging when Managing Rosacea: A Review. *Acta Derm Venereol*. 2021 Oct 31;101(10):adv00584. doi: 10.2340/actadv.v101.356.
9. Schaller M, Lenders D, Handgretinger G, Gawaz A. Topical ivermectin 10 mg/g cream alone or in combination with oral doxycycline for patients with perioral dermatitis (POD): A retrospective case series. *J Dtsch Dermatol Ges*. 2025 Jun;23(6):759-762. doi: 10.1111/ddg.15691. Epub 2025 Mar 22.
10. Siddiqui K, Stein Gold L, Gill J. The efficacy, safety, and tolerability of ivermectin compared with current topical treatments for the inflammatory lesions of rosacea: a network meta-analysis. *Springerplus*. 2016 Jul 22;5(1):1151. doi: 10.1186/s40064-016-2819-8.
11. Soolantra (ivermectin) [prescribing information]. Dallas, TX: Galderma Laboratories LP; October 2022.
12. Stein L, Kircik L, Fowler J, et al. Efficacy and safety of ivermectin 1% cream in treatment of papulopustular rosacea: results of two randomized, double-blind, vehicle-controlled pivotal studies. *J Drugs Dermatol*. 2014 Mar;13(3):316-23.
13. Stein Gold L, Kircik L, Fowler J, et al. Long-term safety of ivermectin 1% cream vs azelaic acid 15% gel in treating inflammatory lesions of rosacea: results of two 40-week controlled, investigator-blinded trials. *J Drugs Dermatol*. 2014 Nov;13(11):1380-6.
14. Stone DU, Chodosh J. Ocular rosacea: an update on pathogenesis and therapy. *Curr Opin Ophthalmol*. 2004 Dec;15(6):499-502.
15. Taieb A, Ortonne JP, Ruzicka T, et al. Superiority of ivermectin 1% cream over metronidazole 0.75% cream in treating inflammatory lesions of rosacea: a randomized, investigator-blinded trial. *Br J Dermatol*. 2015 Apr;172(4):1103-10. doi: 10.1111/bjd.13408. Epub 2015 Feb 11.
16. van Zuuren EJ, Fedorowicz Z, Carter B, van der Linden MM, Charland L. Interventions for rosacea. *Cochrane Database Syst Rev*. 2015 Apr 28;2015(4):CD003262. doi: 10.1002/14651858.CD003262.pub5.
17. Zaenglein AL, Pathy AL, Schlosser BJ, et al. Guidelines of care for the management of acne vulgaris. *J Am Acad Dermatol*. 2016;74(5):945-973.e33. doi:10.1016/j.jaad.2015.12.037

#### Clinical Guideline Revision / History Information

Original Date: 09/18/2024

Reviewed/Revised: 01/01/2026