

Bonjesta (doxylamine/pyridoxine extended-release)

Diclegis (doxylamine/pyridoxine delayed-release)

Disclaimer

Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.

Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.

Summary

Nausea and vomiting during pregnancy (also commonly referred to as "morning sickness") is a common condition that affects the health of a pregnant individual and their developing baby. It can diminish the pregnant individual's quality of life and significantly contribute to health care costs and time lost from work. Modifications in diet and medications to prevent or reduce nausea and vomiting in pregnancy are beneficial to prevent hyperemesis gravidarum. Treatment in the early stages may prevent more serious complications, including hospitalization. Safe and effective treatments are available for more severe cases, and mild cases of nausea and vomiting of pregnancy may be resolved with lifestyle (e.g., avoiding triggers such as heat, humidity, odors, noises or motion, use of acupressure wristbands) and dietary changes (e.g., small, frequent meals, substituting triggering foods for bland, salty, low-fat, protein-dominant, and/or dry foods). Nausea and vomiting of pregnancy should be distinguished from nausea and vomiting related to other causes, as the mechanism of nausea/vomiting is distinctly unique in this circumstance.

Pyridoxine, vitamin B6, is recommended as a first line treatment for pregnant individuals who have mild nausea and infrequent vomiting. Individuals who have nausea with frequent vomiting, or have inadequate relief from dietary changes, avoidance of triggering factors, and pyridoxine may benefit from the addition of medications with antihistamine properties such as doxylamine (Unisom), dimenhydrinate (Dramamine), diphenhydramine (Benadryl), prochlorperazine (Compazine), or promethazine (Phenergan).

Both pyridoxine and doxylamine are available over-the-counter (OTC) as individual products. Both drugs, taken alone or together, have been found to be safe during pregnancy with no known harmful effects to the developing baby.

Bonjesta (doxylamine/pyridoxine extended release) and Diclegis (doxylamine/pyridoxine delayed release) are fixed dose combinations of doxylamine and pyridoxine approved by the Food and Drug Administration (FDA) for the treatment of pregnancy-related nausea and vomiting in those who have not responded to conservative management.

- Diclegis (doxylamine/pyridoxine delayed release) contains 10 mg doxylamine succinate and 10 mg pyridoxine hydrochloride.
- Bonjesta (doxylamine/pyridoxine extended release) contains 20 mg doxylamine succinate and 20 mg pyridoxine hydrochloride.

Definitions

“Conservative management” refers to changes to diet and lifestyle that might help an individual feel better. Examples include taking vitamins, adjusting meal times or changing the types of foods eaten.

“Hyperemesis gravidarum” (HG) refers to the most severe form of nausea and vomiting of pregnancy. HG may be diagnosed with the loss of 5 percent (%) of their pre-pregnancy weight and/or has other problems related to dehydration or loss of body fluids. Those with hyperemesis gravidarum often need acute treatment in a hospital setting to stop the vomiting and restore body fluids.

“Nausea” refers to an uneasy feeling in one’s stomach that might indicate the need to vomit. Other symptoms that can accompany nausea include weakness, sweating, or dizziness.

“Vomiting” refers to the act of ejecting stomach contents from the mouth in an uncontrolled manner. Vomiting is a common action associated with “morning sickness” during the early stages of pregnancy.

Medical Necessity Criteria for Initial Authorization

The Plan considers Bonjesta or Diclegis (doxylamine/pyridoxine) medically necessary when ALL of the following criteria are met:

1. The member is 18 years of age or older; *AND*
2. The member has a diagnosis of pregnancy-related nausea and vomiting, confirmed by laboratory work and/or medical records; *AND*
3. The prescribing provider submits documentation or attestation stating that the member is unable to use, or has inadequate relief from lifestyle modifications such as dietary changes or avoidance of triggers; *AND*
4. The member is unable to use, or has not benefited from, the individual components (OTC doxylamine and pyridoxine) taken concomitantly as separate products; *AND*

5. The requested medication is being prescribed for use within Food and Drug Administration (FDA) approved dosing (*See Table 1 for FDA-approved dosing*) or as supported by evidence-based literature; **AND**
6. Clinical chart documentation is provided for review to substantiate the above listed requirements..

If the above prior authorization criteria are met, Bonjesta or Diclegis (doxylamine/pyridoxine) will be approved for 9 months (or the duration of the pregnancy).

Table 1: Dosage Information

| Product | Initial dose | Maximum dose |
|--|--|---|
| Bonjesta (doxylamine/pyridoxine extended release) | Initially, take one BONJESTA extended-release tablet orally at bedtime (Day 1). If this dose adequately controls symptoms the next day, continue taking one tablet daily at bedtime only. However, if symptoms persist on Day 2, increase the daily dose to one tablet in the morning and one tablet at bedtime. | The maximum recommended dose is two tablets per day, one in the morning and one at bedtime. |
| Diclegis (doxylamine/pyridoxine delayed release) | Initially, take two DICLEGIS delayed-release tablets orally at bedtime (Day 1). If this dose adequately controls symptoms the next day, continue taking two tablets daily at bedtime. However, if symptoms persist into the afternoon of Day 2, take the usual dose of two tablets at bedtime that night then take three tablets starting on Day 3 (one tablet in the morning and two tablets at bedtime). If these three tablets adequately control symptoms on Day 4, continue taking three tablets daily. Otherwise take four tablets starting on Day 4 (one tablet in the morning, one tablet mid-afternoon and two tablets at bedtime). | The maximum recommended dose is four tablets (one in the morning, one in the mid-afternoon and two at bedtime) daily. |

Experimental or Investigational / Not Medically Necessary

Bonjesta or Diclegis (doxylamine/pyridoxine) for any other indication is considered not medically necessary by the Plan, as it is deemed to be experimental, investigational, or unproven.

References

1. ACOG (American College of Obstetrics and Gynecology): Nausea and Vomiting of Pregnancy FAQs. Available at: <https://www.acog.org/womens-health/faqs/morning-sickness-nausea-and-vomiting-of-pregnancy>. Accessed 21 July 2025
2. American College of Obstetricians and Gynecologists (ACOG) Committee on Practice Bulletins-Obstetrics. ACOG Practice Bulletin No. 189: nausea and vomiting of pregnancy. *Obstet Gynecol.* 2018;131(1):e15-e30.
3. Bonjesta (doxylamine and pyridoxine) [prescribing information]. Bryn Mawr, PA: Duchesnay USA; March 2022.
4. Campbell K, Rowe H, Azzam H, Lane CA. The management of nausea and vomiting of pregnancy. *J Obstet Gynaecol Can.* 2016;38(12):1127-1137. doi:10.1016/j.jogc.2016.08.009
5. Diclegis ((doxylamine succinate and pyridoxine hydrochloride) [prescribing information]. Bryn Mawr, PA: Duchesnay USA; March 2022.
6. Herrell HE. Nausea and vomiting of pregnancy. *Am Fam Physician* 2014 Jun 15;89(12):965-970.
7. Koren G, Clark S, Hankins GD, et al. Effectiveness of delayed-release doxylamine and pyridoxine for nausea and vomiting of pregnancy: a randomized placebo controlled trial. *Am J Obstet Gynecol.* 2010;203(6):571.e1-571.e5717. doi:10.1016/j.ajog.2010.07.030
8. Koren G, Clark S, Hankins GD, et al. Maternal safety of the delayed-release doxylamine and pyridoxine combination for nausea and vomiting of pregnancy; a randomized placebo controlled trial. *BMC Pregnancy Childbirth.* 2015;15:59. doi:10.1186/s12884-015-0488-1
9. Kothari S, Afshar Y, Friedman LS, Ahn J. AGA Clinical Practice Update on Pregnancy-Related Gastrointestinal and Liver Disease: Expert Review. *Gastroenterology.* 2024 Oct;167(5):1033-1045. doi: 10.1053/j.gastro.2024.06.014. Epub 2024 Aug 12.
10. Micromedex® (electronic version). IBM Watson Health, Greenwood Village, Colorado, USA. Available at: <https://www.micromedexsolutions.com>. Accessed August 16, 2021.
11. Nelson-Piercy C, Dean C, Shehmar M, et al. The Management of Nausea and Vomiting in Pregnancy and Hyperemesis Gravidarum (Green-top Guideline No. 69). *BJOG.* 2024 Jun;131(7):e1-e30. doi: 10.1111/1471-0528.17739. Epub 2024 Feb 4. Erratum in: *BJOG.* 2025 Jun 19. doi: 10.1111/1471-0528.18258.
12. Oliveira LG, Capp SM, You WB, Riffenburgh RH, Carstairs SD. Ondansetron compared with doxylamine and pyridoxine for treatment of nausea in pregnancy: a randomized controlled trial. *Obstet Gynecol.* 2014 Oct;124(4):735-742. doi: 10.1097/AOG.0000000000000479.
13. Sharifzadeh F, Kashanian M, Koohpayehzadeh J, Rezaian F, Sheikhansari N, Eshraghi N. A comparison between the effects of ginger, pyridoxine (vitamin B6) and placebo for the treatment of the first trimester nausea and vomiting of pregnancy (NVP). *J Matern Fetal Neonatal Med.* 2018 Oct;31(19):2509-2514. doi: 10.1080/14767058.2017.1344965. Epub 2017 Jul 7. PMID: 28629250.
14. Wu XK, Gao JS, Ma HL, et al. Acupuncture and Doxylamine-Pyridoxine for Nausea and Vomiting in Pregnancy: A Randomized, Controlled, 2 × 2 Factorial Trial. *Ann Intern Med.* 2023 Jul;176(7):922-933. doi: 10.7326/M22-2974. Epub 2023 Jun 20.

Clinical Guideline Revision / History Information

Original Date: 10/14/2021

Reviewed/Revised: 12/01/2021, 06/23/2022, 06/29/2023, 09/18/2024, 12/01/2025