

CeQur Simplicity Insulin Delivery System

- CeQur Simplicity Inserter
- CeQur Simplicity Patch

Disclaimer

Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.

Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.

CeQur Simplicity Insulin Delivery System	1
Summary	1
Definitions	2
Clinical Indications	4
Medical Necessity Criteria for Clinical Review	5
General Medical Necessity Criteria	5
Medical Necessity Criteria for Initial Clinical Review	5
Initial Indication-Specific Criteria	5
Diabetes Mellitus	5
Medical Necessity Criteria for Subsequent Clinical Review	6
Subsequent Indication-Specific Criteria	6
Diabetes Mellitus	6
Experimental / Investigational, or unproven[s]	7
References	7
Clinical Guideline Revision / History Information	8

Summary

Diabetes mellitus (commonly referred to as diabetes) is a chronic (long-term) metabolic disorder characterized by high blood glucose (sugar). This may be because the pancreas (an organ in the belly) does not make enough insulin (a hormone), or because the body is not responding to insulin the way it should (insulin insensitivity). Insulin helps glucose get into cells in the body, giving it energy. With diabetes, sugar builds up in the blood because the body stops responding to insulin, or because there is not enough of it. Inadequate glycemic control can lead to serious complications including cardiovascular disease, nephropathy, retinopathy, and neuropathy. The goal of diabetes management is to maintain blood glucose levels as close to the normal range as possible to prevent or delay complications.

Diabetes is classified into two main types:

- Type 1 diabetes mellitus (T1DM): The pancreas produces little or no insulin.
- Type 2 diabetes mellitus (T2DM): The body's cells do not respond effectively to insulin, and sometimes the pancreas does not produce enough insulin.

Insulin therapy is essential for all individuals with type 1 diabetes and many with type 2 diabetes to achieve optimal glycemic control. Intensive insulin therapy aims to mimic normal physiologic insulin secretion through the use of basal insulin for control of glucose throughout the day and bolus or prandial insulin to cover food intake during mealtime and correct hyperglycemia. This is most often accomplished through multiple daily injections (MDI) using basal and rapid-acting insulins or continuous subcutaneous insulin infusion (CSII) via an insulin pump.

However, despite advances in insulin formulations and delivery systems, many individuals fail to reach glycemic targets. Adherence to insulin regimens is often suboptimal due to the burden of multiple daily injections. Insulin pumps, while allowing for more precise insulin delivery, require extensive training and are not acceptable to all patients.

The CeQur Simplicity insulin delivery device is a wearable patch that provides mealtime bolus insulin dosing in 1-unit or 2-unit increments with the push of a button. Depending on the patch version, it can be worn for 4 days or for up to 7 days and holds up to 100 (1.0 mL) to 240 (2.4 mL) units of rapid-acting insulin (e.g., Novolog [insulin aspart] or Humalog [insulin lispro]), that is filled by the individual (the patch does not come preloaded with insulin). CeQur Simplicity provides an alternative delivery option for prandial (mealtime) insulin coverage as part of a basal-bolus regimen. CeQur Simplicity insulin delivery device is approved for those 21 years of age and older, and has been studied in those with type 2 or type 1 diabetes.

The CeQur Simplicity insulin delivery device differs from other disposable insulin pump devices as it provides only mealtime/bolus insulin. The CeQur Simplicity insulin delivery device does not provide basal insulin, which is required for those with type 1 diabetes mellitus, and in some with type 2 diabetes mellitus.

Definitions

"Blood Glucose" is the main sugar found in the blood and the body's main source of energy. It is also called glucose or blood sugar. The blood level of glucose is noted in milligrams per deciliter (mg/dL). When blood sugar is too high for long periods of time, complications can occur as a result of blood vessel damage.

"Blood Glucose Monitors" are small, portable machines used to check blood glucose levels in the ambulatory setting. A member will prick his/her fingertip and place a small sample of blood into the device for a glucose reading. There are a number of different types of blood glucose monitors for specialized situations, such as those for members with visual impairments.

"Basal insulin" refers to the low, steady infusion or injection of insulin to control blood glucose between meals and overnight. It mimics the background insulin normally produced by the pancreas.

"Basal-bolus regimen" is an insulin regimen that includes both long-acting basal insulin for control of glucose between meals and bolus doses of rapid-acting insulin (see "bolus insulin") to cover food intake and correct hyperglycemia.

"Bolus insulin" refers to the administration of insulin to cover food intake and correct hyperglycemia. It is typically given before meals or snacks.

"Continuous Glucose Monitoring (CGM)" serves as an alternative to self-monitoring of blood glucose (SMBG) with a home glucose monitor for patients who have diabetes and require multiple daily measurements.

"Continuous subcutaneous insulin infusion (CSII)" is a method of insulin delivery using an insulin pump to provide a continuous infusion of rapid-acting insulin subcutaneously. The basal rate is programmed and the user administers bolus doses for meals or corrections. The rate is typically determined by the individual's provider and is unique to their particular insulin needs.

"Diabetes mellitus" refers to a group of metabolic disorders characterized by hyperglycemia resulting from defects in insulin secretion, insulin action, or both.

"Disposable Insulin Infusion Pumps" are insulin-delivery devices that consist of disposable components. Both the OmniPod and V-Go insulin delivery systems have disposable components, but they differ in that the OmniPod system is programmable, whereas the V-Go system is non-programmable.

"Documentation" refers to written information, including but not limited to:

- Up-to-date chart notes, relevant test results, and/or relevant imaging reports to support diagnoses; or

- Prescription claims records, and/or prescription receipts to support prior trials of formulary alternatives.

"Glucose/Glycemic variability" refers to the fluctuations or swings in blood glucose levels over hours or days. Higher glycemic variability may be associated with increased risk of diabetes complications.

"Hemoglobin A1c (HbA1c)" refers to a form of hemoglobin that is bound to glucose. The HbA1c test reflects average blood glucose levels over the past 2 to 3 months. It is used to monitor overall glycemic control in diabetes management.

"Hyperglycemia" is excessive blood glucose. Fasting hyperglycemia is blood glucose above a desirable level after a person has fasted for at least 8 hours. Postprandial hyperglycemia is blood glucose above a desirable level 1 to 2 hours after a person has eaten.

"Hypoglycemia" is a condition that occurs when one's blood glucose is lower than normal, usually less than 70 mg/dL. Signs include hunger, nervousness, shakiness, perspiration, dizziness or lightheadedness, sleepiness, and confusion. If left untreated, hypoglycemia may lead to unconsciousness. Hypoglycemia is treated by consuming a carbohydrate-rich food such as a glucose tablet or juice. It may also be treated with an injection of glucagon if the person is unconscious or unable to swallow.

"Insulin" is a hormone made by the beta cells of the pancreas. Insulin allows glucose to enter the cells in the body for use in energy production, and when it is inadequate, the sugar remains in the blood leading to diabetes. There are a variety of oral and parenteral medications that can increase insulin production, increase the body's sensitivity to existing insulin and reduce blood sugar. Insulin can also be injected or infused when lifestyle changes and non-insulin medications are inadequate.

"Multiple daily injections (MDI)" refers to an insulin regimen involving the administration of basal insulin once or twice daily and separate bolus injections of rapid-acting insulin with meals and snacks. It aims to mimic normal physiologic insulin secretion.

"Time in range (TIR)" is the percentage of time that a person's blood glucose levels are within the target range, typically 70-180 mg/dL. It is a metric of glycemic control often assessed through continuous glucose monitoring.

"Type 1 diabetes" is a form of diabetes mellitus caused by autoimmune destruction of the insulin-producing beta cells of the pancreas, leading to absolute insulin deficiency.

"Type 2 diabetes" is a form of diabetes mellitus characterized by insulin resistance and relative insulin deficiency. It is often associated with obesity and may be managed with lifestyle changes, oral medications, and/or insulin.

"[s]" indicates state mandates may apply.

Clinical Indications

Medical Necessity Criteria for Clinical Review

General Medical Necessity Criteria

The Plan considers CeQur Simplicity Insulin Delivery System and its components medically necessary when ALL of the following criteria are met:

1. The member is 21 years of age or older; *AND*
2. The member has a documented diagnosis of type 1 or type 2 diabetes mellitus; *AND*
3. The requested quantity of CeQur Simplicity meets ONE (1) of the following:
 - a. The request(s) is/are being prescribed for use within the Plan's Quantity Limit of:
 - i. Meets ONE of the following:
 1. Starter Kit - 1 Kit per 365 days; *or*
 2. Inserter Kit - 1 Inserter per 365 days; *and/or*
 - ii. Meets ONE of the following for patch refills:
 1. IF the request is for the 2U CeQur Simplicity 4-Day Wear Patch the member meets ONE of the following:
 - a. For members using ≤ 180 units of insulin per 96 hours - 10 patches per 30 days (or 30 patches per 90-days); *or*
 - b. For members using > 180 units of insulin per 96 hours - 30 patches per 30-days (or 90 patches per 90 days); *or*
 2. IF the request is for the 2U CeQur Simplicity 2.4ml Extended Wear Patch: 1 box (containing 5 patches) per 30 days; *or*
 3. IF the request is for the 1U CeQur Simplicity 2.4ml Extended Wear Patch: 1 box (containing 4 patches) per 30 days; *or*
 - b. IF the requested quantity exceeds the Plan's Quantity Limit then clinical rationale is provided demonstrating need for increased use (e.g., for more frequent patch changes due to high insulin requirements); *AND*
 4. The member meets the applicable [Medical Necessity Criteria for Initial Clinical Review](#) or [Subsequent Clinical Review](#) listed below.

Medical Necessity Criteria for Initial Clinical Review

Initial Indication-Specific Criteria

Diabetes Mellitus

The Plan considers CeQur Simplicity Insulin Delivery System and its components medically necessary when ALL of the following criteria are met:

5. The member meets the above [General Medical Necessity Criteria](#); *AND*
6. The member requires blood glucose checks daily OR is using a continuous glucose monitor (CGM); *AND*

- a. The member has suboptimal glycemic control (e.g., hemoglobin A1c >7.0%, hypoglycemia, hyperglycemia, glucose variability/fluctuations); *AND*
- b. The member meets ONE (1) of the following:
 - i. The member has been on an insulin pump for at least the past 90 days; *or*
 - ii. The member has been on a basal-bolus insulin regimen with insulin pens or syringes for at least the past 90 days *AND* meets ONE (1) of the following:
 1. Has tried and failed to achieve glycemic control with traditional insulin delivery methods (pens or syringes)^[s]; *or*
 2. Has a contraindication, intolerance, or inability to use traditional insulin delivery methods due to physical limitations, visual impairment, and/or needle phobia/aversion; *and*
7. The member will use a rapid-acting insulin analog (i.e., Humalog [insulin lispro], Novolog [insulin aspart]) with the device for the purpose of meal-time only bolus dosing.

If the above prior authorization criteria are met, the requested product will be authorized for up to 12-months.^[s]

Continued Care

Medical Necessity Criteria for Subsequent Clinical Review

Subsequent Indication-Specific Criteria

Diabetes Mellitus

The Plan considers CeQur Simplicity Insulin Delivery System and its components medically necessary when ALL of the following criteria are met:

1. The member meets the above applicable [General Medical Necessity Criteria](#) and/or [Initial Clinical Review](#); *AND*
2. The member has demonstrated improvement in glycemic control from baseline as evidenced by any ONE (1) of the following:
 - a. Reduction in hemoglobin A1c; *and/or*
 - b. Improved time in range (TIR) on the member's CGM (e.g., percent time in the member's individual goal blood glucose range); *and/or*
 - c. Reduction in frequency or severity of hypoglycemia episodes; *and/or*
 - d. Reduction in diabetic ketoacidosis episodes; *and/or*
 - e. Improvement in glycemic variability; *and/or*
 - f. Achievement of other individual glycemic targets (e.g., average blood glucose, reduction in preprandial/postprandial or fasting blood glucose).

If the above reauthorization criteria are met, the requested product will be authorized for up to 12-months.^[s]

Experimental / Investigational, or unproven^[5]

CeQur Simplicity insulin delivery device for any other indication or use is considered experimental, investigational, or unproven. Non-covered indications include, but are not limited to, the following[†]:

- Use in those under 21 years of age. The safety and efficacy in those 21 years of age and younger has not been adequately studied.
- Use in gestational diabetes (not including preexisting diabetes diagnosis in someone who is pregnant or planning pregnancy). While CeQur requires the use of a rapid-acting insulin product (e.g., Humalog [insulin lispro] or Novolog [insulin aspart]), which are both approved in the setting of gestational diabetes, the CeQur Simplicity insulin delivery device has not been studied in this population.
- Delivery of medications *other* than rapid-acting insulin (i.e., Humalog [insulin lispro], Novolog [insulin aspart]). The CeQur Simplicity insulin delivery has only been studied with these forms of rapid insulin for the purpose of replacing bolus/mealtime insulin.
- Use as a standalone device without a basal insulin regimen. CeQur Simplicity insulin delivery device has only been studied in those who require both basal and bolus insulin.

†NOTE: The safety and effectiveness of CeQur Simplicity have not been established in these situations. The device is only FDA-cleared for use in adult patients with type 1 or type 2 diabetes who require bolus insulin as part of a basal-bolus regimen. Any off-label use of CeQur Simplicity is considered not medically necessary by the Plan, as it is deemed to be experimental, investigational, or unproven. Coverage is limited to the FDA-approved indications when the medical necessity criteria outlined in this policy are met.

References

1. American Diabetes Association Professional Practice Committee; 7. Diabetes Technology: Standards of Care in Diabetes—2024. *Diabetes Care* 1 January 2024; 47 (Supplement_1): S126–S144. <https://doi.org/10.2337/dc24-S007>
2. American Diabetes Association Professional Practice Committee; 7. Diabetes Technology: Standards of Care in Diabetes - 2025. *Diabetes Care*. 2025;48(suppl. 1):S146-S166. <https://doi.org/10.2337/dc25-S007>.
3. American Diabetes Association Professional Practice Committee. 7. Diabetes Technology: Standards of Care in Diabetes-2024. *Diabetes Care*. 2024 Jan 1;47(Suppl 1):S126-S144. doi: 10.2337/dc24-S007.
4. American Diabetes Association Professional Practice Committee et al: 15. Management of diabetes in pregnancy: standards of medical care in diabetes-2025. *Diabetes Care*. 2025; 48(Suppl. 1):S306-320. <https://doi.org/10.2337/dc25-25>.
5. American Diabetes Association Professional Practice Committee et al: 15. Management of diabetes in pregnancy: standards of medical care in diabetes-2026. *Diabetes Care*. 1 January 2026; 49(Suppl. 1):S321-3338. <https://doi.org/10.2337/dc26-S015>.
6. American Diabetes Association Professional Practice Committee; Summary of Revisions: Standards of Care in Diabetes—2024. *Diabetes Care* 1 January 2024; 47 (Supplement_1): S5–S10. <https://doi.org/10.2337/dc24-SREV>
7. American Diabetes Association Professional Practice Committee; Summary of Revisions: Standards of Care in Diabetes—2025. *Diabetes Care* 2025; 48 (Supplement_1): S6–S13. <https://doi.org/10.2337/dc25-SREV>.

8. American Diabetes Association Professional Practice Committee; Summary of Revisions: Standards of Care in Diabetes—2026. *Diabetes Care*. 1 January 2026; 49 (Suppl. 1): S6–S12. <https://doi.org/10.2337/dc26-SREV>.
9. Battelino T, Danne T, Bergenstal RM, et al. Clinical Targets for Continuous Glucose Monitoring Data Interpretation: Recommendations From the International Consensus on Time in Range. *Diabetes Care*. 2019 Aug;42(8):1593-1603. doi: 10.2337/dci19-0028. Epub 2019 Jun 8.
10. Bergenstal RM, Johnson ML, Aroda VR, Brazg RL, Dreon DM, Frias JP, Kruger DF, Molitch ME, Mullen DM, Peyrot M, Richter S, Rosenstock J, Serusclat P, Vance C, Weinstock RS, Levy BL. Comparing Patch vs Pen Bolus Insulin Delivery in Type 2 Diabetes Using Continuous Glucose Monitoring Metrics and Profiles. *J Diabetes Sci Technol*. 2022 Sep;16(5):1167-1173. doi: 10.1177/19322968211016513. Epub 2021 May 19. PMID: 34008442; PMCID: PMC9445326.
11. Bergenstal RM, Peyrot M, Dreon DM, et al. Implementation of Basal-Bolus Therapy in Type 2 Diabetes: A Randomized Controlled Trial Comparing Bolus Insulin Delivery Using an Insulin Patch with an Insulin Pen. *Diabetes Technol Ther*. 2019 May;21(5):273-285. doi: 10.1089/dia.2018.0298. Epub 2019 Apr 26.
12. BioSpace. CeQur® Announces Exciting Advancements: 2 Recent FDA Clearances and New Clinical Evidence to be Presented at ADA. Available at: <https://www.biospace.com/press-releases/cequr-announces-exciting-advancements-2-recent-fda-clearances-and-new-clinical-evidence-to-be-presented-at-ada>. Accessed June 9, 2026.
13. CeQur SA. (2024). CeQur Detail-Aid-APM-02-0005-Rev-F. Available from: <https://myceqursimplicity.com/wp-content/uploads/Detail-Aid-APM-02-0005-Rev-F.pdf>.
14. CeQur Simplicity website. 2024. CeQur Simplicity Features [online]. Available at: <https://www.myceqursimplicity.com>
15. CeQur Simplicity FAQs. 2026. FAQs [online]. Available at: <https://myceqursimplicity.com/hcp-faq/>. Accessed 9 January 2026.
16. Food and Drug Administration. (2024). SPECIAL 510(k) SUBSTANTIAL EQUIVALENCE DETERMINATION DECISION SUMMARY for K233447. Silver Spring, MD: Food and Drug Administration. Available from: https://www.accessdata.fda.gov/cdrh_docs/reviews/K233447.pdf.
17. Isaacs D, Kruger DF, Shoger E, Chawla H. Patient Perceptions of Satisfaction and Quality of Life Regarding Use of a Novel Insulin Delivery Device. *Clin Diabetes*. 2023 Spring;41(2):198-207. doi: 10.2337/cd22-0034. Epub 2022 Sep 9.
18. LifeScan Europe, a division of Cilag GmbH. "510(k) Summary for the OneTouch Via On-Demand Insulin Delivery System (K163357)." U.S. Food and Drug Administration, 7 June 2017, www.fda.gov/media/98547/download. Accessed 03/06/2024.
19. Mader JK, Lilly LC, Aberer F, et al. Improved glycaemic control and treatment satisfaction with a simple wearable 3-day insulin delivery device among people with Type 2 diabetes. *Diabet Med*. 2018 Oct;35(10):1448-1456. doi: 10.1111/dme.13708. Epub 2018 Jul 5.
20. Wahlqvist P, Warner J, Morlock R. Cost-effectiveness of Simple Insulin Infusion Devices Compared to Multiple Daily Injections in Uncontrolled Type 2 Diabetics in the United States Based on a Simulation Model. *J Health Econ Outcomes Res*. 2018 Aug 22;6(1):84-95. doi: 10.36469/9789.
21. Zraick V, Dreon D, Nalk R, Shearer D, Crawford S, Bradford J, Levy B. 2016. Patient User Experience Evaluation of Bolus Patch Insulin Delivery System. Poster presented at the American Diabetes Association's 76th Scientific Sessions. Abstract 995-P. New Orleans, LA, USA.

Clinical Guideline Revision / History Information

Original Date: 3/21/2024

Reviewed/Revised: 10/29/2024, 10/01/2025, 03/02/2026, 08/03/2026