



adefovir dipivoxil (Hepsera)

Disclaimer

Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.

Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.

adefovir dipivoxil (Hepsera)	1
Summary	2
Definitions	2
Clinical Indications	3
Medical Necessity Criteria for Initial Clinical Review	3
Initial Indication-Specific Criteria	3
Chronic Hepatitis B Virus (HBV) Infection	3
Medical Necessity Criteria for Subsequent Clinical Review	3
Subsequent Indication-Specific Criteria	3
Chronic Hepatitis B Virus (HBV) Infection	3
Experimental / Investigational, or unproven	3
References	4
Clinical Guideline Revision / History Information	4

Summary

Hepatitis B (HBV) is a viral infection that targets the liver, leading to either acute or chronic disease. Acute infections are often self-limiting and do not always require treatment; however, a small fraction can progress to chronic hepatitis B. Chronic infections, defined as the persistence of hepatitis B surface antigens present after 6 months, pose significant health risks as they can result in serious liver complications, such as cirrhosis and hepatocellular carcinoma (HCC). The goal of HBV treatment is to reduce the risk of HCC, however initiation of therapy is based on several factors including presence or absence of cirrhosis, alanine aminotransferase levels (a liver enzyme test, with higher levels indicating liver damage or disease) and HBV DNA levels. Management of chronic HBV includes treatment with either pegylated interferon or nucleoside/nucleotide analogs (i.e., entecavir [Baraclude] or Vemlidy [tenofovir alafenamide]). The American Association for the Study of Liver Diseases (AASLD) has published recommendations for management of HBV at <https://www.aasld.org>.

Adefovir dipivoxil (Hepsera) oral tablet is a nucleotide analogue approved for the treatment of chronic hepatitis B virus (HBV) infections in those 12 years and older. It is usually taken once daily, however, dosing interval adjustments may be required in certain individuals with renal impairments, such as those on hemodialysis. While adefovir (Hepsera) is approved to treat hepatitis B infection, according to major guidelines it is not preferred for chronic hepatitis B treatment due to its high rate of resistance with long-term use. Other drugs with a higher barrier to drug resistance are usually recommended, such as tenofovir (Vemlidy) or entecavir (Baraclude).

Definitions

“ALT and AST” refers to liver enzymes aspartate transaminase (AST) and alanine aminotransferase (ALT) that are indicators of liver damage or injury from different types of diseases or conditions.

“Creatinine clearance” (CrCl) is the volume of blood plasma cleared of creatinine, a waste product from the body, per unit of time. It is an indicator or measurement of kidney function.

“DNA” refers to Deoxyribonucleic Acid, a small molecule inside of cells that contain genetic information.

“Documentation” refers to written information, including but not limited to:

- Up-to-date chart notes, relevant test results, and/or relevant imaging reports to support diagnoses; or
- Prescription claims records, and/or prescription receipts to support prior trials of formulary alternatives.

“Nucleos(t)ide analog” is a class of drugs used to treat viral infections that work by inhibiting the ability of the virus to replicate.

“[s]” indicates state mandates may apply.

Clinical Indications

Medical Necessity Criteria for Initial Clinical Review

Initial Indication-Specific Criteria

Chronic Hepatitis B Virus (HBV) Infection

The Plan considers adefovir (Hepsera) medically necessary when ALL of the following criteria are met:

1. The requested medication is being prescribed by or in consultation with a gastroenterologist, hepatologist, or infectious disease specialist; *AND*
2. The member is 12 years of age or older; *AND*
3. The member has a documented diagnosis of chronic hepatitis B virus (HBV) infection confirmed by appropriate laboratory test(s); *AND*
4. The member is unable to use, or has tried and failed entecavir (Baraclude) or tenofovir (Vemlidy)^[s]; *AND*
5. Adefovir (Hepsera) is being prescribed at a dose and frequency that is within FDA approved labeling OR is supported by compendia or evidence-based published dosing guidelines for the requested indication.

If the above prior authorization criteria are met, Adefovir Dipivoxil (Hepsera) will be approved for up to 6 months.^[s]

Continued Care

Medical Necessity Criteria for Subsequent Clinical Review

Subsequent Indication-Specific Criteria

Chronic Hepatitis B Virus (HBV) Infection

The Plan considers adefovir (Hepsera) medically necessary when ALL of the following criteria are met:

1. The member meets the above applicable [Initial Clinical Review](#); *AND*
2. The member has documented clinical improvement from baseline or maintenance of disease activity as evidenced by ONE of the following:
 - a. Decrease or normalization of serum aminotransferases (ALT or AST) levels; *or*
 - b. Loss or reduction of HBeAg or HBsAg; *or*
 - c. Minimal histologic evidence of liver injury; *or*
 - d. Reduction, suppression, or undetectable levels of serum HBV DNA levels.

[Experimental / Investigational, or unproven](#)^[s]

Adefovir dipivoxil (Hepsera) for any other use is considered experimental, investigational, or unproven.

References

1. Adefovir [prescribing information]. Toronto, Ontario: Apotex Inc.; March 2019.

2. Ali FS, Nguyen MH, Hernaez R, et al. AGA Clinical Practice Guideline on the Prevention and Treatment of Hepatitis B Virus Reactivation in At-Risk Individuals. *Gastroenterology*. 2025 Feb;168(2):267-284. doi: 10.1053/j.gastro.2024.11.008.
3. AS Lok et al. Antiviral therapy for chronic hepatitis B viral infection in adults: a systematic review and meta-analysis. *Hepatology* 2016; 63:284.
4. European Association for the Study of the Liver (EASL). European Association for the Study of the Liver. EASL 2017 clinical practice guidelines on the management of hepatitis B virus infection. *J Hepatol*. 2017;67(2):370-398. doi:10.1016/j.jhep.2017.03.021
5. Ghany MG, Pan CQ, Lok AS, Feld JJ, Lim JK, Wang SH, Kim AY, Tang AS, Nguyen MH, Naggie S, Sulkowski MS, Norberto RB, Chen J, Murad MH, Mohammad KS, Terrault NA. AASLD ISDA Practice Guideline on treatment of chronic hepatitis B. *Hepatology*. 2026 Apr 1;83(4):974-997. doi: 10.1097/HEP.0000000000001549. Epub 2025 Nov 4. PMID: 41186418.
6. Guidelines for the prevention, diagnosis, care and treatment for people with chronic hepatitis B infection [Internet]. Geneva: World Health Organization; 2024 Mar. PMID: 40424433.
7. Hadziyannis SJ, Tassopoulos NC, Heathcote EJ, et al. Long-term therapy with adefovir dipivoxil for HBeAg-negative chronic hepatitis B. *N Engl J Med*. 2005 Jun 30;352(26):2673-81. doi: 10.1056/NEJMoa042957.
8. Hepsera (adefovir) [prescribing information]. Foster City, CA: Gilead Sciences Inc; December 2018.
9. Peters MG, Hann HW, Martin P, et al. Adefovir dipivoxil alone or in combination with lamivudine in patients with lamivudine-resistant chronic hepatitis B. *Gastroenterology*. 2004 Jan;126(1):91-101. doi: 10.1053/j.gastro.2003.10.051.
10. Sigmapharm Laboratories. Adefovir dipivoxil tablet [prescribing information]. Bensalem, PA; 2016 Aug.
11. Terrault NA, Bzowej NH, Chang KM et al. AASLD guidelines for treatment of chronic hepatitis B. *Hepatology*. 2016; 63:261-83.
12. Terrault NA, Lok ASF, McMahon BJ, et al. Update on prevention, diagnosis, and treatment of chronic hepatitis B: AASLD 2018 hepatitis B guidance. *Hepatology*. 2018;67(4):1560-1599. doi:10.1002/hep.29800
13. World Health Organization. Guidelines for the prevention, care and treatment of persons with chronic hepatitis B infection. March 2015. Geneva: World Health Organization; 2015.
14. World Health Organization (WHO). Guidelines for the prevention, diagnosis, care and treatment for people with chronic hepatitis B infection. 29 March 2024. Available at: <https://www.who.int/publications/i/item/9789240090903>.

Clinical Guideline Revision / History Information

Original Date: 11/05/2020

Reviewed/Revised: 10/14/2021, 12/01/2021, 9/15/2022, 9/21/2023, 12/19/2024, 01/01/2026, 01/04/2027