

## Auvelity (Dextromethorphan and Bupropion)

### Disclaimer

*Clinical guidelines are developed and adopted to establish evidence-based clinical criteria for utilization management decisions. Clinical guidelines are applicable according to policy and plan type. The Plan may delegate utilization management decisions of certain services to third parties who may develop and adopt their own clinical criteria.*

*Coverage of services is subject to the terms, conditions, and limitations of a member's policy, as well as applicable state and federal law. Clinical guidelines are also subject to in-force criteria such as the Centers for Medicare & Medicaid Services (CMS) national coverage determination (NCD) or local coverage determination (LCD) for Medicare Advantage plans. Please refer to the member's policy documents (e.g., Certificate/Evidence of Coverage, Schedule of Benefits, Plan Formulary) or contact the Plan to confirm coverage.*

### Summary

Auvelity (dextromethorphan hydrobromide and bupropion hydrochloride), is an oral combination product indicated for the treatment of major depressive disorder (MDD) in adults. Major depressive disorder is a mental health disorder that causes symptoms of sadness, hopelessness, and loss of interest in things. It can disrupt relationships and everyday activities, such as work, school, and activities that are usually pleasant. There are many medications from several classes that are available to treat major depressive disorder including selective serotonin-reuptake inhibitors (SSRIs), serotonin- and norepinephrine-reuptake inhibitors (SNRIs), tricyclic antidepressants, monoamine oxidase (MAO) inhibitors, and other antidepressants (e.g., bupropion, mirtazapine, trazodone).

### Definitions

**“Major depressive disorder”**, also known as **(MDD)**, is a psychiatric condition characterized by persistent low mood, low energy, or loss of interest in enjoyable activities causing substantial impairment in daily life. MDD is thought to be caused by a combination of genetic, environmental and psychological

factors. Risk factors include family history, major life changes, certain medications, chronic health problems, and substance use disorders.

**“Neurotransmitter”** is a molecule that sends signals from neurons to different parts of the body (e.g., muscles).

### Medical Necessity Criteria for Initial Authorization

The Plan considers **Auvelity (Dextromethorphan and Bupropion)** medically necessary when **ALL** of the following criteria are met:

1. The member is 18 years of age or older; **AND**
2. The member has a diagnosis of major depressive disorder (MDD); **AND**
3. The member is unable to use or has adequately tried and failed a minimum **ONE** (1) month trial of at least **THREE** (3) therapies, from at least **THREE** (3) of the following classes:
  - a. Noradrenergic and dopaminergic antidepressants (bupropion); **and/or**
  - b. Noradrenergic and specific serotonin antidepressants (e.g., mirtazapine); **and/or**
  - c. Selective serotonin reuptake inhibitors (e.g., citalopram, escitalopram, fluoxetine, paroxetine, sertraline); **and/or**
  - d. Serotonin-norepinephrine reuptake inhibitors (e.g., duloxetine, venlafaxine IR/ER); **and/or**
  - e. Tricyclic antidepressants (e.g., amitriptyline, nortriptyline); **AND**
4. Clinical chart documentation is provided for review to substantiate the above listed requirements.

**If the above prior authorization criteria are met, the requested medication will be approved for 12 months.**

### Medical Necessity Criteria for Reauthorization

Reauthorization for 12 months will be granted if recent chart documentation (within the last 3 months) shows that the member has experienced therapeutic response to the requested medication as evidenced by **ONE** (1) of the following:

1. clinical improvement (e.g., reduction in signs and symptoms, including residual symptoms) in symptoms since starting the requested medication; **OR**
2. stability in condition (e.g., restoration of prior level of psychosocial and occupational function) since starting the requested medication.

## Experimental or Investigational / Not Medically Necessary

Auvelity (Dextromethorphan and Bupropion) for any other indication (including Alzheimer's disease agitation and smoking cessation) is considered not medically necessary by the Plan, as it is deemed to be experimental, investigational, or unproven.

## References

1. American Psychiatric Association: Practice Guideline for the Treatment of Patients with Major Depressive Disorder, 3rd edition, 2010. Available at: [https://psychiatryonline.org/pb/assets/raw/sitewide/practice\\_guidelines/guidelines/mdd.pdf](https://psychiatryonline.org/pb/assets/raw/sitewide/practice_guidelines/guidelines/mdd.pdf). Published October 2010. Accessed October 2022.
2. Amir Qaseem, Douglas K. Owens, Itziar Etxeandia-Ikobaltzeta, et al; Clinical Guidelines Committee of the American College of Physicians. Nonpharmacologic and Pharmacologic Treatments of Adults in the Acute Phase of Major Depressive Disorder: A Living Clinical Guideline From the American College of Physicians. *Ann Intern Med.*2023;176:239-252. [Epub 24 January 2023]. doi:10.7326/M22-2056.
3. Auvelity (dextromethorphan and bupropion) [prescribing information]. New York, NY: Axsome Therapeutics Inc; May 2024.
4. Auvelity (dextromethorphan and bupropion) [prescribing information]. New York, NY: Axsome Therapeutics Inc; December 2022.
5. Bauer M, Pfennig A, Severus E, Whybrow PC, Angst J, Möller HJ; World Federation of Societies of Biological Psychiatry Task Force on unipolar depressive disorders. World Federation of Societies of Biological Psychiatry (WFSBP) guidelines for biological treatment of unipolar depressive disorders. Part 1: update 2013 on the acute and continuation treatment of unipolar depressive disorders. *World J Biol Psychiatry.* 2013;14(5):334-385. doi: 10.3109/15622975.2013.804195.
6. Bauer M, Severus E, Köhler S, Whybrow PC, Angst J, Möller HJ; WFSBP Task Force on treatment guidelines for unipolar depressive disorders. World Federation of Societies of Biological Psychiatry (WFSBP) guidelines for biological treatment of unipolar depressive disorders. Part 2: maintenance treatment of major depressive disorder-update 2015. *World J Biol Psychiatry.* 2015;16(2):76-95. doi:10.3109/15622975.2014.1001786.
7. Iosifescu DV, Jones A, O'Gorman C, et al. Efficacy and safety of AXS-05 (dextromethorphan-bupropion) in patients with major depressive disorder: a phase 3 randomized clinical trial (GEMINI). *J Clin Psychiatry.* 2022;83(4):21m14345. doi:10.4088/JCP.21m14345.
8. Mauri MC, Fiorentini A, Paletta S, Altamura AC. Pharmacokinetics of antidepressants in patients with hepatic impairment. *Clin Pharmacokinet.* 2014;53(12):1069-1081. doi:10.1007/s40262-014-0187-5.
9. Mullish BH, Kabir MS, Thursz MR, Dhar A. Review article: depression and the use of antidepressants in patients with chronic liver disease or liver transplantation. *Aliment Pharmacol Ther.* 2014;40(8):880-892. doi:10.1111/apt.12925.
10. Stahl SM. Dextromethorphan/bupropion: a novel oral NMDA (N-methyl-d-aspartate) receptor antagonist with multimodal activity-Addendum. *CNS Spectr.* 2020;25(6):803. doi:10.1017/S109285291900155X.

11. Tabuteau H, Jones A, Anderson A, Jacobson M, Iosifescu DV. Effect of AXS-05 (dextromethorphan-bupropion) in major depressive disorder: a randomized double-blind controlled trial. *Am J Psychiatry*. 2022;179(7):490-499. doi:10.1176/appi.ajp.21080800.
12. VA/DoD Clinical Practice Guideline. (2022). *The Management of Major Depressive Disorder*. Washington, DC: U.S. Government Printing Office.

#### Clinical Guideline Revision / History Information

Original Date: 12/08/2022

Reviewed/Revised: 9/21/2023, 12/02/2024