




Influenza Vaccine 2020 - 2021

	Name	Supply	CPT Code	Manufacturer	Age	OK for use in children younger than 3 years old	Dosage	Contains Neomycin	Contains Gentamicin	Light Sensitive	Preservative Free	Latex Free	OK for use in Pregnant women	OK for Nursing Mothers
1	 Fluzone Quadrivalent	Prefilled Syringe 0.5ml	90686Z	Sanofi	 6 months and older	Yes	0.5ml/dose (up to 9 years old repeat in 1 month if first year receiving vaccine) ^{1,2}	No	No	No	Yes	Yes	Yes 	Yes
2	 Fluarix Quadrivalent	Prefilled Syringe 0.5ml	90686X	GSK	 6 months and older	Yes	0.5ml/dose (up to 9 years old repeat in 1 month if first year receiving vaccine) ^{1,2}	No	Yes	Yes	Yes	Yes	Yes 	Yes
3	 Flulaval Quadrivalent	Prefilled Syringe 0.5ml	90686V	GSK	 6 months and older	Yes	0.5ml/dose (up to 9 years old repeat in 1 month if first year receiving vaccine) ^{1,2}	No	No	Yes	Yes	Yes	Yes 	Yes
4	 Flucelvac (cell culture derived) Quadrivalent	Prefilled Syringe 0.5ml	90674C	Seqirus	4 years and older	No	0.5ml/dose (up to 9 years old repeat in 1 month if first year receiving vaccine) ^{1,2}	No	No	Yes	Yes	Yes	Yes 	Yes
5	 Fluad Quadrivalent	Prefilled Syringe 0.5ml	90694D	Seqirus	65 years and older	No	0.5ml/dose	Yes	No	Yes	Yes	Yes	Not approved for patients less than 65years old	Not approved for patients less than 65y.o
6	 FluMist Quadrivalent Intranasal Spray	Single-dose prefilled sprayer 0.2ml	90672M	AstraZeneca	2 years to 49 years	Yes	0.2ml dose ³ (up to 9 years old repeat in 1 month if first year receiving vaccine) ^{1,2}	No	Yes	Yes	Yes	Yes	No	Yes, as long as there are no contraindications to FluMist

³ Children aged 6 months through 8 years who have previously received ≥ 2 total doses of trivalent or quadrivalent influenza vaccine before July 1, 2020 require only 1 dose for 2020–21. The 2 doses of influenza vaccine do not have to have been administered in the same season or consecutive seasons.

Children in this age group who have not previously received ≥ 2 doses of trivalent or quadrivalent influenza vaccine before July 1, 2020 or whose previous influenza vaccination history is unknown, require 2 doses for the 2020–21 season. The interval between the 2 doses should be at least 4 weeks apart.

³ Children who require 2 doses of flu vaccine do not need to receive the same flu vaccine both times; any age appropriate brand can be used for either dose. half the contents (0.1ml) of the sprayer into each nostril for a total of 0.2ml