EVOLUTION: BACTERIA

TO BEETHOVEN

KEY TERMS:	evolution fossil record		assailable ural selection	scientific mutation
NOTE-TAKING COLUMN: Complete this section <u>during</u> the video. Include definitions and key terms.			CUE COLUMN: Complete this section <u>after</u> the video.	
Who proposed the Theory of E	Evolution in 1859?		What are the prima of Evolution?	ry tenets of the Theory
What startling discovery did Jamake in the 1950's?	ames Watson and Francis Cr	ick		rian Explosion and the ow do they 'disrupt' utionary theory?
How many atoms are in our ga	alaxy?			

DISCUSSION & REVIEW QUESTIONS:

- Towards the beginning of the video, Dr. Meyer contends that, "...with some modification it [the Theory of Evolution] has been embraced as unassailable by the science community over the last century. As evolutionary biologist Richard Dawkins says, 'If you meet somebody who claims not to believe in evolution, that person is ignorant, stupid or insane.'" Why do you think that the science community treated Mr. Darwin's theory as 'unassailable' for so long? Why do you think that Dr. Dawkins characterizes his sentiment as a 'belief' in evolution- as if science is a belief system? Do you agree with Dr. Dawkins' assertion? Why or why not?
- Later, Dr. Meyer presents the first of two reasons to doubt evolutionary theory, the Cambrian Explosion, by pointing out that, "A weird and wonderful thing happened 530 million years ago: a whole bunch of major groups of animals, what scientists call the 'phyla,' appeared abruptly within a geologically short window of time- about ten million years. These novel animal forms, exhibiting proto-types of most animal body designs we see today, emerged in the fossil record without evidence of earlier ancestors... A huge number of diverse animals appeared, with no discernible antecedents. So where did they come from?" How would you answer Dr. Meyer's question? Why does this evidence present a problem for evolutionary theory?
- Dr. Meyer later reminds us that, "...natural selection only 'selects' sequences that random mutations generate. Yet experiments have established that DNA sequences capable of making stable proteins are extremely rare- and, thus, really hard to stumble on randomly." What does this information mean, in terms of supporting or not supporting Mr. Darwin's theory? Explain. Why is the 'random' factor so important to Dr. Meyer's point here? Explain.
- Further, Dr. Meyer explains that, "...even 4 billion years of life's history is not enough time to overcome a search problem this big [of finding new DNA sequences that can build a functional and/or stable protein]." Why do you think that this is the case? Explain.
- At the end of the video, Dr. Meyer concludes that, "Scientists who know about these problems [the Cambrian Explosion and the DNA Enigma], are not 'ignorant, stupid, or insane,' they are just appropriately... skeptical." Why do you think that Dr. Meyer makes a distinction between scientists who know about the problems and those who don't? Explain. Do you think that scientists should still be skeptical regarding Darwinian theory? Why or why not?

EXTEND THE LEARNING:

CASE STUDY: de novo genes

INSTRUCTIONS: Read the article "How evolution builds genes from scratch," then answer the questions that follow.

- What did scientists long assume, in terms of new genes? Who is Helle Baalsrud, and what did she do? Where have scientists found de novo genes in the last five years? What are de novo genes causing a rethink of? What was the conventional wisdom regarding new genes? What is the starting material for some new genes? What do researchers still have to work out in terms of identifying a gene as de novo, and what questions still remain about de novo genes? What do scientists still wonder, in terms of making genes from scratch? What was Susumu Ohno's thinking about genes? What do all of the gene- duplicating processes have in common? What is 'Junk DNA?' Who is Mar Alba, and what did her research show? What was her speculation on her findings? What could an understanding of what de novo genes are doing for their hosts do to help explain about the existence of de novo genes themselves? What is an ORF, and what could it theoretically do? What does Dr. Carvunis think she has found? What is a proto-gene? How many de novo genes have been found in humans? Who is Claudio Casola, and what does he think about de novo genes, in terms of evolution?
- How do you think the information in this article fits with the information presented in the video? Do you think that de novo genes support Mr. Darwin's theory? Why or why not? Do you think that any of the information in this article helps to explain the Cambrian Explosion? Explain. Do you think that the information in this article helps to solve the DNA Enigma? If no, why not? If yes, in what way?
- Do you think that the Theory of Evolution will prevail as factual truth someday, at least to some extent? Why or why not?



- 1. In what year did Charles Darwin propose his Theory of Evolution?
 - a. 1839
 - b. 1849
 - c. 1859
 - d. 1869

2. The purpose of the conference for evolutionary biologists that Dr. Meyer attended in London was ______.

- a. to address the fundamental principles of prevailing Intelligent Design theory
- b. to address growing doubts about the modern version of Darwin's theory
- c. to present new evidence supporting the Theory of Evolution
- d. to present evidence that supports the opinions of Richard Dawkins

3. About 530 million years ago a whole bunch of novel animal forms, exhibiting prototypes of most animal body designs we see today, emerged in the fossil record without evidence of earlier ancestors.

- a. True
- b. False

4. What startling discovery did Watson and Crick make in the 1950's?

- a. That proteins will mutate when exposed to neon gas.
- b. That cells need to move in order to survive.
- c. That protein sequences are sometimes rejected by certain cells.
- d. That a DNA molecule stores information as a four-character digital code.

5. Finding a new DNA sequence capable of building a functional protein is like searching blindfolded for a single marked atom among a ______ Milky Way galaxies!

- a. million
- b. billion
- c. trillion
- d. novemquinquagintillion



- 1. In what year did Charles Darwin propose his Theory of Evolution?
 - a. 1839
 - b. 1849
 - c. 1859
 - d. 1869

2. The purpose of the conference for evolutionary biologists that Dr. Meyer attended in London was ______.

- a. to address the fundamental principles of prevailing Intelligent Design theory
- b. to address growing doubts about the modern version of Darwin's theory
- c. to present new evidence supporting the Theory of Evolution
- d. to present evidence that supports the opinions of Richard Dawkins

3. About 530 million years ago a whole bunch of novel animal forms, exhibiting prototypes of most animal body designs we see today, emerged in the fossil record without evidence of earlier ancestors.

- a. True
- b. False

4. What startling discovery did Watson and Crick make in the 1950's?

- a. That proteins will mutate when exposed to neon gas.
- b. That cells need to move in order to survive.
- c. That protein sequences are sometimes rejected by certain cells.
- d. That a DNA molecule stores information as a four-character digital code.

5. Finding a new DNA sequence capable of building a functional protein is like searching blindfolded for a single marked atom among a _____ Milky Way galaxies!

- a. million
- b. billion
- c. trillion
- d. novemquinquagintillion





GENES FROM THE JUNKYARD

Scientists long assumed that new genes appear when evolution tinkers with old ones. It turns out that natural selection is much more creative.

BY ADAM LEVY

n the depths of winter, water temperatures in the ice-covered Arctic Ocean can sink below zero. That's cold enough to freeze many fish, but the conditions don't trouble the cod. A protein in its blood and tissues binds to tiny ice crystals and stops them from growing.

Where codfish got this talent was a puzzle that evolutionary biologist Helle Tessand Baalsrud wanted to solve. She and her team at the University of Oslo searched the genomes of the Atlantic cod (*Gadus morhua*) and several of its closest relatives, thinking they would track down the cousins of the antifreeze gene. None showed up. Baalsrud, who at the time was a new parent, worried that her lack of sleep was causing her to miss something obvious.

But then she stumbled on studies suggesting that genes do not always evolve from existing ones, as biologists long supposed. Instead, some are fashioned from desolate stretches of the genome that do not code for any functional molecules. When she looked back at the fish genomes, she saw hints this might be the case: the antifreeze protein — essential to the cod's survival — had seemingly been built from scratch¹.

The cod is in good company. In the past five years, researchers have found numerous signs of these newly minted '*de novo*' genes in every lineage they have surveyed. These include model organisms such as fruit flies and mice, important crop plants and humans; some of the genes are expressed in brain and testicular tissue, others in various cancers.

De novo genes are even prompting a rethink of some portions of evolutionary theory. Conventional wisdom was that new genes

Some cod species have a newly minted gene involved in preventing freezing.

tended to arise when existing ones are accidentally duplicated, blended with others or broken up,

but some researchers now think that *de novo* genes could be quite common: some studies suggest at least one-tenth of genes could be made in this way; others estimate that more genes could emerge *de novo* than from gene duplication. Their existence blurs the boundaries of what constitutes a gene, revealing that the starting material for some new genes is non-coding DNA (see 'Birth of a gene').

The ability of organisms to acquire new genes in this way is testament to evolution's "plasticity to make something seemingly impossible, possible", says Yong Zhang, a geneticist at the Chinese Academy of Sciences' Institute of Zoology in Beijing, who has studied the role of *de novo* genes in the human brain.

But researchers have yet to work out how to definitively identify a gene as being *de novo*, and questions still remain over exactly how — and how often — they are born. Scientists also wonder why evolution would bother making genes from scratch when so much gene-ready material already exists. Such basic questions are a sign of how young the field is. "You don't have to go back that many years before *de novo* gene evolution was dismissed," Baalsrud says.

NEW ARRIVALS

Back in the 1970s, geneticists saw evolution as a rather conservative process. When Susumu Ohno laid out the hypothesis that most genes evolved through duplication², he wrote that "In a strict sense, nothing in evolution is created *de novo*. Each new gene must have arisen from an already existing gene."

Gene duplication occurs when errors in the DNA-replication process produce multiple instances of a gene. Over generations, the versions accrue mutations and diverge, so that they eventually encode different molecules, each with their own function. Since the 1970s, researchers have found a raft of other examples of how evolution tinkers with genes — existing genes can be broken up or 'laterally transferred' between species. All these processes have something in common: their main ingredient is existing code from a welloiled molecular machine.

But genomes contain much more than just genes: in fact, only a few per cent of the human genome, for example, actually encodes genes. Alongside are substantial stretches of DNA — often labelled 'junk DNA' — that seem to lack any function. Some of these stretches share features with proteincoding genes without actually being genes themselves: for instance, they are littered with three-letter codons that could, in theory, tell the cell to translate the code into a protein.

It wasn't until the twenty-first century that

scientists began to see hints that non-coding sections of DNA could lead to new functional codes for proteins. As genetic sequencing advanced to the point that researchers could compare entire genomes of close relatives, they began to find evidence that genes could disappear rather quickly during evolution. That made them wonder whether genes could just as quickly spring into being.

In 2006 and 2007, evolutionary geneticist David Begun at the University of California, Davis, published what many regard as the first papers to make the case for particular genes arising *de novo* in fruit flies^{3.4}. The studies linked these genes to male reproduction: Begun found they were expressed in the testes and the seminal fluid gland, where it seemed the powerful evolutionary force of sexual selection was driving gene birth.

Shortly before that, evolutionary genomicist Mar Albà at the Hospital del Mar Medical Research Institute in Barcelona, Spain, had shown that the younger a gene is, evolutionarily speaking, the faster it tends to evolve⁵. She speculated that this might be because the molecules encoded by younger genes are

"IT'S LIKE A BETA LAUNCH."

less polished and need more tuning, and that this could be a consequence of the genes having arisen *de novo* — they were not tied to a previous function as tightly as those that had evolved from older genes. Both Albà and Begun recall that it was challenging to publish their early work on the topic. "There was a lot of scepticism," says Albà. "It's amazing how things have changed."

Studies have also started to unpick what de novo genes do. One gene allows the thale cress plant (Arabidopsis thaliana) to produce starch, for instance, and another helps yeast cells to grow. Understanding what they are doing for their hosts should help to explain why they exist — why it is advantageous to create from scratch rather than evolve from existing material. "We're not going to understand why these genes are evolving if we don't understand what they're doing," says Begun.

GENES-IN-WAITING

Studying *de novo* genes turns out to be part genetics, part thought experiment. "Why is our field so difficult?" asks Anne-Ruxandra Carvunis at the University of Pittsburgh in Pennsylvania. "It is because of philosophical issues." At its heart is a question that Carvunis has been asking for a decade: what is a gene?

A gene is commonly defined as a DNA or RNA sequence that codes for a functional molecule. The yeast genome, however, has hundreds of thousands of sequences, known as open reading frames (ORFs), that could theoretically be translated into proteins, but that geneticists assumed were either too short or looked too different from those in closely related organisms to have a probable function.

When Carvunis studied yeast ORFs for her PhD, she began to suspect that not all of these sections were lying dormant. In a study⁶ published in 2012, she looked at whether these ORFs were being transcribed into RNA and translated into proteins — and, just like genes, many of them were — although it was unclear whether the proteins were useful to the yeast, or whether they were translated at high enough levels to serve a function. "So what is a gene? I don't know," Carvunis says. What she thinks she has found, though, is "raw material — a reservoir — for evolution".

Some of these genes-in-waiting, or what Carvunis and her colleagues called protogenes, were more gene-like than others, with longer sequences and more of the instructions necessary for turning the DNA into proteins. The proto-genes could provide a fertile testing ground for evolution to convert non-coding material into true genes. "It's like a beta launch," suggests Aoife McLysaght, who works on molecular evolution at Trinity College Dublin.

Some researchers have gone beyond observation to manipulate organisms into expressing non-coding material. Michael Knopp and his colleagues at Uppsala University, Sweden, showed that inserting and expressing randomly generated ORFs into Escherichia coli could enhance the bacterium's resistance to antibiotics, with one sequence producing a peptide that increased resistance 48-fold⁷. Using a similar approach, Diethard Tautz and his team at the Max Planck Institute for Evolutionary Biology in Plön, Germany, showed that half of the sequences slowed the bacterium's growth, and one-quarter seemed to speed it up^8 — although that result is debated. Such studies suggest that peptides from random sequences can be surprisingly functional.

But random sequences of DNA could also code for peptides that are "reactive and nasty and have a tendency to aggregate and do bad things", says evolutionary biologist Joanna Masel of the University of Arizona in Tucson. Expressing these sequences at low levels could help natural selection to weed out potentially dangerous portions — those that create messy or misfolded proteins — so that what remains in a species is relatively benign.

Creating genes from non-coding regions could have some benefits over other genemaking methods, says Albà. Gene duplication is a "very conservative mechanism" she says, producing well-adapted proteins cut from the same cloth as their ancestors; *de novo* genes, by contrast, are likely to produce markedly different molecules. That could make it



difficult for them to fit into wellestablished networks of genes and proteins — but they could also be better suited to certain new tasks.

A newly minted gene could help an organism to respond to a change in its environment, for instance. This seems to have been the case for the cod, which acquired its antifreeze protein as the Northern Hemisphere cooled some 15 million years ago.

BIRTH RATE

To trace which of an organism's genes were made *de novo*, researchers need comprehensive sequences for the organism and its close relatives. One crop plant that fits the bill is rice. The sweltering heat of Hainan, a tropical island in southern China, is the perfect environment for growing the crop — although the working conditions can be trying. "It's horrible," says evolutionary geneticist Manyuan Long of the University of Chicago, Illinois. It's so hot "you can cook your egg in the sand".

Long's team wanted to know how many genes had emerged de novo in the strain Oryza sativa japonica, and what proteins those genes might be making. So the team lined up its genome against those of its close relatives and used an algorithm to pick out regions that contained a gene in some species but lacked it in others. This allowed the researchers to identify the non-coding DNA that led to the gene in question, and track its journey to being a gene. They could also tot up the number of de novo genes that appeared in the strain: 175 genes over 3.4 million years of evolution⁹ (over the same period, the strain gained 8 times as many genes from duplication).

The study gets at one of the field's biggest preoccupations: how to tell whether a gene is truly de novo. Answers vary wildly, and approaches are still evolving. For example, an early study found 15 de novo genes in the whole primate order¹⁰; a later attempt found 60 in humans alone¹¹. One option for finding candidate de novo genes is to use an algorithm to search for similar genes in related species. If nothing shows up, then it's possible that the gene arose de novo. But failing to find a relative doesn't mean no relative is there: the gene could have been lost along the way, or might have shape-shifted far away from its kin. The rice study got around this by explicitly identifying the pieces of non-coding DNA that became de novo genes.

Over long evolutionary timescales — much longer than the few million years of rice evolution — it is hard to distinguish between a *de novo* gene and one that has simply diverged

BIRTH OF A GENE

Scientists long assumed that evolution made new genes from old ones by copying them in error, or by fusing together or breaking apart existing ones. Now, more and more examples are emerging of genes being created '*de novo*', from barren non-coding portions of the genome.

How genes work

Genes are usually considered any stretches of DNA that code for useful molecules. To make a protein, DNA is transcribed into RNA, which is then translated. Three-letter pieces of sequence called codons dictate which portions of the RNA to translate.



Making a gene de novo

Genes can evolve from non-coding portions of DNA by gaining transcription and codons, in either order. At first, the products of these 'proto-genes' might be dysfunctional or disordered.



too far from its ancestors to be recognizable, so determining the absolute number of genes that have arisen *de novo* rather than from duplication "is an almost unanswerable question", says Tautz.

To demonstrate how varied the results of different methods can be, evolutionary geneticist Claudio Casola at Texas A&M University in College Station used alternative approaches to reanalyse the results of previous studies, and failed to verify 40% of the *de novo* genes they had proposed¹². To Casola, this points to the need to standardize tests. Currently, he says, "it seems to be very inconsistent".

Counting *de novo* genes in the human genome comes with the same trail of caveats. But where *de novo* genes have been identified, researchers are beginning to explore their roles in health and disease. Zhang and his colleagues have found that one gene unique

to humans is expressed at a greater level in the brains of people with Alzheimer's disease¹³, and previous work¹⁴ had linked certain variants of the gene to nicotine dependence. For Zhang, research that links de novo genes to the human brain is tantalizing. "We know that what makes us human is our brains," he says, "so there must be some genetic kit to push the evolution of our brain." That suggests an avenue for future studies. Zhang suggests that researchers could investigate the genetic kit through experiments with human organoids - cultured cells that serve as a model organ.

De novo genes could have implications for understanding cancer, too. One such gene — unique to humans and chimpanzees — has been linked to cancer progression in mouse models of neuroblastoma¹⁵. And cancer-causing versions of human papillomavirus include a gene that is not present in noncancer-causing forms¹⁶.

Many *de novo* genes remain uncharacterized, so the potential importance of the process to health and disease is unclear. "It will take some time before we fully understand to what extent it contributes to human health and to what extent it contributes to the origin of the human species," says Carvunis.

Although *de novo* genes remain enigmatic, their existence makes one thing clear: evolution can readily make something from nothing. "One of the beauties of working with *de novo* genes," says Casola, "is that it shows how dynamic genomes are."

Adam Levy *is a science journalist based in London.*

- Baalsrud, H. T. et al. Mol. Biol. Evol. 35, 593–606 (2018).
- Ohno, S. Evolution by Gene Duplication (Springer, 1970).
- Begun, D. J., Lindfors, H. A., Thompson, M. E. & Holloway, A. K. Genetics 172, 1675–1681 (2006)
- Begun, D. J., Lindfors, H. A., Kern, A. D. & Jones, C. D. Genetics 176, 1131–1137 (2007).
- Albà, M. M. & Castresana, J. Mol. Biol. Evol. 22, 598–606 (2005).
- 6. Carvunis, À.-R. et al. Nature 487, 370–374 (2012).
- Knopp, M. *et al. mBio* **10**, e00837-19 (2019).
 Neme, R., Amador, C., Yildirim, B., McConnell, E. &
- Tautz, D. Nature Eco. Evol. **1**, 0127 (2017).
- Zhang, L. et al. Nature Ecol. Evol. 3, 679–690 (2019).
 Toll-Riera, M. et al. Mol. Biol. Evol. 26, 603–612 (2009).
- 11.Wu, D.-D., Irwin, D. M. & Zhang, Y.-P *PLoS Genet.* 7, e1002379 (2011).
- 12.Casola, C. Genome Biol. Evol. 10, 2906-2918 (2018).
- 13.Li, C.-Y. et al. PLoS Comput. Biol. 6, e1000734 (2010).
- 14.Wang, D., Ma, J. Z. & Li, M. D. *Pharmacogenomics J.* **5**, 166–172 (2005).
- 15. Suenaga, Y. *et al. PLoS Genet.* **10**, e1003996 (2014). 16. Willemsen, A., Félez-Sánchez, M. & Bravo, I. G.
- Genome Biol. Evol. **11**, 1602–1617 (2019).