

INSTRUCTIONS FOR USE

EYSUVIS [eye-SU-vis] (loteprednol etabonate ophthalmic suspension) 0.25% for topical ophthalmic use

This Instructions for Use contains information on how to properly administer EYSUVIS.

Important Information You Need to Know Before Using EYSUVIS

- EYSUVIS is for use in the eye.
- Wash your hands before using EYSUVIS.
- **Do not** use if the tamper-evident seal is not intact.
- **Do not** let the EYSUVIS dropper tip touch your eye, fingers, or any other surfaces to avoid contamination or injury to your eye.
- Use EYSUVIS exactly as your doctor tells you to.
- If you are using EYSUVIS with other eye (ophthalmic) medicines, you should wait at least 5 minutes between using EYSUVIS and the other medicine.
- If you wear contact lenses, remove them before using EYSUVIS.
- Put the pink cap back on EYSUVIS after each use.

Before you use EYSUVIS for the first time:

There are two caps on your bottle of EYSUVIS. Hold the bottle firmly by its neck. Remove the white cap by twisting it **clockwise** (See **Figure A**). Throw away the white cap. EYSUVIS is now ready to use.



Figure A

Follow Steps 1 to 6 each time you use EYSUVIS.

- 1) Wash your hands well.
- 2) Shake the EYSUVIS bottle for 2 to 3 seconds before using (See **Figure B**).



Figure B

- 3) Remove the pink cap from the top of the EYSUVIS dropper by turning it **counterclockwise** (See **Figure C**). Keep the pink cap. **Do not** let the EYSUVIS dropper tip touch your eye, fingers, or any other surface.



- 4) Turn the EYSUVIS bottle upside down (See Figure D).



Figure D

- 5) Tilt your head back. Hold the bottle directly above your affected eye. Squeeze the middle of the EYSUVIS bottle gently to put 1 to 2 drops (follow your doctor's instruction) into the affected eye (See Figure E).



Figure E

- 6) Place the pink cap back onto the EYSUVIS bottle and tighten by turning clockwise (See Figure F).



Figure F

If you use contact lenses, wait for 15 minutes before placing them back in.

How should I store EYSUVIS?

- Store EYSUVIS upright between 59°F to 77°F (15°C to 25°C).
- **Do not** freeze.
- After opening, EYSUVIS can be used until the expiration date (EXP) on the bottle. The expiration date can be found on the lower right side of the label on the bottle.

Keep EYSUVIS and all medicines out of the reach of children.

This Instructions for Use has been approved by the U.S. Food and Drug Administration.

Manufactured for:
Kala Pharmaceuticals, Inc.
Watertown, MA 02472

Approved: 10/2020