

# CREATURE COMFORTS VETERINARY HOSPITAL

## BLAND DIET FEEDING INSTRUCTIONS

**GENERAL INSTRUCTIONS:** A bland diet is a diet that is soft and highly digestible. It is low in fiber, fat and protein and high in carbohydrates. Bland diets are generally composed of a single carbohydrate source and a single lean protein source. The most common bland diet is boiled rice and boiled lean chicken breast, without skin and bones. Since bland diets are low in fiber, stool production slows and defecation is less frequent. Bland diets are fed to rest the gastric system and to help promote normal stool formation. Animals that are physically sick should not be fed bland diets as a method of treatment.

**PRIOR TO FEEDING A BLAND DIET:** Pets should be fasted for 12 to 24 hours. Young animals should not be fasted more than 12 hours. Fasting will allow the intestinal system to relax and minimize acid secretions that may irritate and inflame the intestinal lining. During the fast, small amounts of water or ice chips may be provided. Never fast a sick animal, as it may cause disease progression and death.

### BASIC BOILED RICE AND CHICKEN BLAND DIET RECIPE:

**CARBOHYDRATE SOURCE:** Boiled white rice.

**LEAN PROTEIN SOURCE:** Chicken breast, no skin and bones.

**PREPARING BOILED RICE:** 1 part white rice with 3 parts water boiled for 20-25 minutes or until the rice is easily crushed.

**PREPARING BOILED CHICKEN:** De-fat chicken breast and boil in water for 10-15 minutes until the chicken meat is easily pulled apart and cooked all the way through.

**MIXING INSTRUCTIONS :** Finely chop the lean protein and mix 2 cups carbohydrate source with ½ cup lean protein source.

**STORAGE INSTRUCTIONS:** Bland diets can be pre-made and stored in the refrigerator for a maximum of 48 hours. The bland diet can be cooked in a batch and frozen in feeding sized portions to minimize preparation time. Thaw and warm the frozen diet prior to feeding.

**PERMITTED LEAN PROTEIN SUBSTITUTIONS:** Pork loin, egg whites, low fat cottage cheese, 7% low fat hamburger, plain low fat yogurt, tofu.

**PERMITTED CARBOHYDRATE SUBSTITUTIONS:** Boiled potatoes, boiled spaghetti, Minute rice, sweet potato, oatmeal.

**PERMITTED COMMERCIAL BLAND DIETS:** Royal Canin Veterinary Diet Intestinal HE; Science Prescription Diet I/D; Eukanuba Veterinary Diet Low Residue.

**FEEDING INSTRUCTIONS:** Estimate 25% of your animal's diet and feed that amount of the bland diet every 6-8 hours. Smaller animals will require less and larger animals will require more, adhering to the same carbohydrate to lean protein source ratio. The bland diet should be fed for 4-5 days with no treats or other food sources until stools are firm.

**TRANSITION BACK TO A NORMAL DIET:** Never transition back to a regular diet rapidly. Transition back to the regular diet over a 1-week period. Start by adding 25% of the regular diet to 75% of the bland diet and feed that combination for 2 days. If stools continue to be firm, the continue substituting the regular diet in 25% increments and feeding the combination in 2 day time intervals until the diet is 100% regular diet. Minimize treats for 1 week after moving the diet back to 100% regular diet.

4/6/13