



# Intranasal Midazolam Kit

## Contents

- Glass vial with midazolam suspension in a light protected container
- Syringes with hypodermic needles
- MAD device

## Directions: Following a seizure, follow the steps below

**1** Take syringe, ensure needle is screwed on tightly, remove cap, insert into inverted glass vial containing Midazolam solution.


**2** Draw up appropriate amount of Midazolam as directed by Veterinarian.



**3** Use one handed-method to safely re-cap the syringe.



**4** Carefully unscrew the capped needle off of the syringe.



- 5** Screw the MAD device onto the syringe tip.



- 6** Tilt your pet's head back and rest the cone of the MAD device in one nostril.



- 7** Slowly drip the contents of syringe into the nostril by gently pressing down on the plunger.

- 8** This can be repeated after two more seizures (3 total per 24 hours) but wait at least 10 minutes between doses to avoid excessive sedation. After the third dose, if your pet has another seizure, please bring him or her into your nearest emergency veterinary hospital for further treatment.

### What to expect:

Your pet may experience some side effects after receiving Midazolam. Side effects may include sedation, delirium, incoordination, increased hunger (sometimes for non-food items) and vocalization for up to several hours after administration.



**BUSH VETERINARY  
NEUROLOGY SERVICE**

If you have questions about the care of your pet, please feel free to contact us.