

Anxiety, Depression, Stress, and Oral Health



Course Author(s): Maria L. Geisinger, DDS, MS

CE Credits: 1 hour

Intended Audience: Dentists, Dental Hygienists, Dental Assistants, Dental Students, Dental Hygiene Students, Dental Assistant Students

Date Course Online: 12/11/2019

Last Revision Date: 12/08/2022

Course Expiration Date: 12/07/2025

Cost: Free

Method: Self-instructional

AGD Subject Code(s): 153

Online Course: www.dentalcare.com/en-us/ce-courses/ce573

Disclaimer: Participants must always be aware of the hazards of using limited knowledge in integrating new techniques or procedures into their practice. Only sound evidence-based dentistry should be used in patient therapy.

Conflict of Interest Disclosure Statement

- The author reports no conflicts of interest associated with this course. She has no relevant financial relationships to disclose.

Introduction

Anxiety, Depression, Stress, and Oral Health seeks to discuss the interrelationship of oral health and mental health disorders, including anxiety and depression and to provide dental healthcare providers with tools to intervene and help improve overall health for individuals with both oral and mental health disorders.

Course Contents

- Overview
- Learning Objectives
- Introduction
- Anxiety and Chronic Stress: A State of Inflammation
 - Definitions, Prevalence, and Treatment of Anxiety Disorders
 - The Physiology of Anxiety and Chronic Stress
- Depression: Physical Manifestations of Psychological Illness
 - Definitions, Prevalence, and Treatment of Depression
 - The Link Between Stress and Depression
- Associations of Oral Health and Anxiety and Depressive Disorders
 - Influence of Psychosocial Factors on Periodontal Diseases and Tooth Loss
 - Influence of Anxiety and Depression on Dental Caries
 - Implications for Anxiety and Depression Medications on Periodontal Disease Progression and Therapy
- Identification of At-risk Patients in the Dental Office: Screening and Referral
- Summary
- Course Test
- References / Additional Resources
- About the Author

Overview

This course seeks to discuss the interrelationship of oral health and mental health disorders, including anxiety and depression and to provide dental healthcare providers with tools to intervene and help improve overall health for individuals with both oral and mental health disorders.

Learning Objectives

Upon completion of this course, the dental professional should be able to:

- Review the current data on the prevalence, severity, and therapy options for anxiety and depression.
- Identify screening tools and warning signs for anxiety and depression that can be employed by dental healthcare providers in the dental office.
- Understand the current evidence linking anxiety and depression and oral diseases,

- including caries and periodontal disease.
- Discuss mechanisms of interaction between antidepressant and anti-anxiety medication and treatment outcomes for periodontal disease and caries.
- Be prepared to work in an integrated interdisciplinary team to assess risk factors, screen, and refer patients who may be at risk for anxiety and depression.

Introduction

Currently, it is estimated that 322 million people worldwide live with depression and approximately 7.6% of U.S. adults had at least one depressive episode in 2017.^{1,2} Furthermore, the highest prevalence (13.1%) was seen in individuals 18-25 years old.^{1,2} Despite these staggering figures, mental health, anxiety, and substance use disorders are still significantly under-reported and treatment is often delayed or avoided due to stigma or embarrassment. Anxiety, depression, and stress have also been linked to poorer oral health status and treatment outcomes.³⁻⁸ This relationship has been linked to the alterations that stress may induce on the immune system. Additionally, adequate treatment and antidepressant agents have anti-inflammatory functions that may serve to improve outcomes during periodontal therapy,⁹ but their adverse effects on salivary flow may impact caries rates.¹⁰ Because of the high prevalence of anxiety and depression in the population and the importance of adequate treatment for oral and overall health outcomes, identification, screening, and referral of at-risk patients seen in the dental office is a critical part of ensuring overall patient wellness. This course seeks to identify risk screening tools to assess anxiety and depression in the patient population seeking dental care and to evaluate the link between these common mental health disorders and oral health.

Anxiety and depression are common in the United States. In general, 1 out of every 6 adults will have depression, which may be situational and/or clinical, at some time in their lifetime.¹¹ Many people who experience depression also have other mental health conditions, particularly anxiety disorders, with up to 60% of individuals diagnosed with depression also meeting the criteria for an anxiety disorder.^{12,13}

Individuals with psychiatric disorders experience increased risk of systemic diseases, including diabetes mellitus, cardiovascular disease, autoimmune diseases, cancer, and chronic obstructive pulmonary disorder.^{5,14-18} Chronic psychosocial stress and depression have also been shown to adversely effect therapy and outcomes of these conditions.¹⁸ Stress and depression have also been linked to higher overall rates of morbidity and mortality within the population across a range of systemic conditions. The mechanisms for these links may be manifold.¹⁶⁻¹⁸ Chronic stress, which has been associated with anxiety disorders and depression, increases systemic inflammatory burden and is associated with damage to the brain region that controls mood, including the hippocampus and prefrontal cortex, making these conditions difficult to treat.¹⁹⁻²⁴ The interrelationship of anxiety and depression and periodontal disease is generally explained through the effects of these psychosocial conditions on the host immune response through a chronic stress mechanism.²⁵ A recent meta-analysis identified a significant association between anxiety disorders and/or depression and periodontitis.²⁶ The underlying reasons for this may be multifactorial, including increased xerostomia due to medications and/or an upregulation of sympathetic nervous system stimulation, a reduction in

motivation for self-care behaviors, such as oral hygiene, in individuals with anxiety and depressive disorders, and/or a common inflammatory burden that may affect the host immune response.^{5,26} Assessment of patients' psychological well-being is critical to understanding their current disease state and their potential response to therapy.

Anxiety and Chronic Stress: A State of Inflammation

Definitions, Prevalence, and Treatment of Anxiety Disorders

Anxiety disorders are defined as a negative affective state resulting from perceptions of threat, characterized by a perceived inability to predict, control, or obtain desired results in upcoming situations. Overall, it is estimated that an average of 3.8% of the global population are affected by anxiety disorders.²⁷⁻²⁹ This estimate suggests that 284 million people experienced anxiety disorders in 2017, making it the most prevalent mental health disorder.²⁷⁻²⁹ The prevalence varies between countries, and these disorders tend to affect women in greater numbers. Females account for approximately 63% of the individuals with anxiety disorders (Figure 1).²⁹ This gender disparity may relate to hormonal differences and/or societal pressures, including




Figure 1. Global prevalence of anxiety disorder varies by country.²⁹

the overwhelming percentage of at-home work performed by women, even when both partners work outside the home. The prevalence of anxiety disorders in the United States was 7.1% in 2017.²⁹

Treatment for anxiety disorders may include non-pharmacologic therapies like cognitive behavioral therapy (CBT), exposure therapy, interpersonal therapy (IPT), and others, pharmacological treatments with anxiolytic medications, or a combination of the two.^{30,31} In some instances, particularly anxiety that is situational or transient, short-term treatment may be appropriate, but many individuals with more severe anxiety symptoms may require long-term ongoing treatment.^{30,31}

The Physiology of Anxiety and Chronic Stress

In most individuals, stress is a response to an external cause and subsides once the situation has resolved.^{5,32-34} When an individual is under acute stress, activation of the sympathetic nervous system, the body's fight-or-flight response, results in release of epinephrine, norepinephrine, and adrenocorticotrophic hormone (ACTH).³² In a short-term situation, a work deadline or on-stage performance, these transmitters allow our bodies to react quickly and perform well under pressure.³² However, when stress is ongoing and results in chronic exposure to these hormones and neurotransmitters, it can have a deleterious

effect on well-being.³⁴⁻³⁶ Elevated cortisol levels lead to increased systemic inflammation and decreased immune function over time. Multiple stress mediators, including monoamines, neuropeptides (neuropeptide Y, calcitonin gene-related peptide, adrenomedullin), and steroid hormones transfer the stress signal to the central nervous system (CNS) through the limbic-hypothalamic-pituitary-adrenal axis.^{5,34-36} Patients with anxiety disorders have a persistent or disproportional stress reaction to stimuli that is generally not considered threatening.^{37,38} Anxiety is characterized by a "persistent feeling of apprehension or dread" and severe forms of anxiety disorders include generalized anxiety, panic disorder, phobias, social anxiety, obsessive-compulsive disorder, and post-traumatic stress disorder (PTSD).^{32,33,37,38} In these cases, individuals experience chronic stress in normal situations and this produces the deleterious effects on their physiology seen in individuals exposed to long-term stressors, such as civilians in a war zone.^{37,38}

Depression: Physical Manifestations of Psychological Illness

Definitions, Prevalence, and Treatment of Depression

Major depression is defined as a period of at least two weeks when a person experienced a depressed mood or loss of interest/pleasure in daily activities and had a majority of




Figure 2. Global prevalence of depression by WHO region.²⁸

specified symptoms, such as sleep disruption, changes in eating patterns, decreased energy, concentration, and/or feelings of low self-worth.³⁹ The estimated prevalence of at least one major depressive episode among U.S. adults was 7.6% in 2017 and was highest in adults aged 18-25 (13.1%).^{28,40,41} Depression is more common in females than in males and rates vary worldwide (Figure 2).^{28,40,41} Treatment types include non-pharmacologic therapy with a health professional — including development of coping mechanisms and/or support systems, medication treatment, and/or a combination of both.⁴² Approximately 35% of all adults diagnosed with depression did not receive any treatment at all.²⁸ Among adolescents, this number is significantly higher with 60.1% of adolescents reporting a major depressive episode receiving no treatment in 2017.²⁸ While depression and anxiety can be linked, they are not identical and reported prevalence worldwide does not align with those of anxiety, indicating that there are other factors that may play into the diagnosis of depression.

The Link Between Stress and Depression

Both stress and depression have been associated with decreased function of the limbic system and prefrontal cortex as well as systemic vascular inflammation and elevated serum cytokine levels.^{5,22,43} Emotional disorders, including depression/depressive state and anxiety/anxiety state, are the most frequently observed symptoms resulting from psychological stress.^{25,26} In fact, 50-60% of depressed patients also meet the lifetime criteria for an anxiety disorder and anxiety disorders have been implicated in the underlying etiology of depression in many cases.⁴⁴⁻⁴⁶ Current hypotheses regarding the interaction of stress and anxiety suggest that the relationship is bidirectional—psychological stress can lead to depression in susceptible individuals and depression may exacerbate anxiety disorders and stress. It has also been suggested that the elevated pro-inflammatory cytokines associated with psychological stress and poor coping mechanisms may worsen or contribute to depressive symptoms in some patients.^{47,48} Nonpharmacological interventions, including cognitive behavioral therapy (CBT) and acceptance and commitment therapy

(ACT) focus on altering coping mechanisms and mindset to improve depression perception and symptoms in susceptible individuals. Antidepressant medications also have the potential to reduce the effects of these proinflammatory cytokines on the brain and, thus, result in symptom amelioration.⁴⁹ It has also been suggested that additional deleterious habits, such as smoking and alcohol consumption, may be associated with both conditions leading to a worsening of symptoms overall.⁵⁰

Associations of Oral Health and Anxiety and Depressive Disorders

Individuals with psychiatric disorders have higher rates of many systemic diseases than non-affected individuals.^{51,52} Oral health is also reported to be significantly worse in individuals with mental illness when compared to their counterparts.^{8,53} Caries, tooth loss, and periodontal disease rates are higher in those individuals with anxiety and depression compared to healthy controls.^{8,53}

It is also well-established that adults with mental health disorders like anxiety and depression and less likely to engage with dental healthcare professionals and seek dental care and they report less frequent health dental home care habits when compared to individuals without mental health disorders.⁵⁴ Additionally, anxiety and depression have been associated with higher rates of deleterious habits including alcohol and tobacco consumption, which can impact their oral health status.⁵⁵ Furthermore, many individuals report that dental visits are anxiety inducing. It has been estimated that 20-30% of the US population reports being “somewhat” or “very” nervous about dental procedures and up to 12% of adults are characterized as dental phobic.⁵⁶

Oral Health is a major determinant of general health, self-esteem, and quality of life, but for patients with mental health disorders, oral healthcare has a low priority for individuals suffering with mental illness. Further, poor oral health may negatively affect mood, confidence/self-esteem, and impact mental health.^{57,58} In individuals with severe mental illness, the one-year prevalence of suboptimal oral health (oral




Figure 3. Model of the Effects of Psychosocial Stress on Oral and Overall Health.^{5,85}

dryness, dental caries, periodontal disease, DMFT, oral cancer/precancer) was 32% and lifetime prevalence was 61%.⁵⁷

Influence of Psychosocial Factors on Periodontal Diseases and Tooth Loss

Many studies suggest that psychosocial stress and ineffective coping mechanisms can influence the onset and progression of chronic inflammatory conditions, such as periodontitis.^{4,5,43,58-66} The role of stress in some acute periodontal conditions, such as necrotizing periodontal diseases, is well-established through both temporal associations of disease onset and a stressor and biological measurements, e.g., urinary cortisol levels.⁶⁷⁻⁶⁹ High rate of progression (Grade C) periodontitis (formerly called “aggressive periodontitis”) and chronic periodontitis have both been associated with stress and depression.^{58-66,70,71} Increasing salivary cortisol levels have been linked to worsening periodontal disease parameters and tooth loss, adjusted for other behavioral and stress variables and increasing stress/distress and depression were linked to periodontal disease severity in a dose-dependent fashion.⁶² Increased coping mechanisms seemed to mitigate this interaction.⁴ The mechanisms of these interactions is less clear, but several hypotheses have been proposed.⁵ Chronic

stress and depression reduce immune responsiveness, and this may result in more periopathogenic bacteria leading to increased periodontal tissue destruction.⁵ Psychosocial disorders, including anxiety disorders and depression, have also been associated with increased levels of systemic pro-inflammatory cytokines, which may potentiate the action of the initiating bacterial insult and worsen the host-mediated periodontal destruction.⁵ Finally, chronic stress and depression may result in alterations in health-related behaviors, including oral hygiene, smoking, and dietary intake.⁵ Individuals with high stress levels who did not brush their teeth during stressful periods were at increased risk of tooth loss when compared to individuals who maintained oral hygiene despite increasing stress and anxiety.⁶² This may indicate that a combination of factors stemming from underlying psychosocial disorders interact resulting in deterioration of systemic and, potentially, oral health as well (Figure 3).⁵ The interaction of psychosocial stress, anxiety, and depression on periodontal health may be related to an increase in pro-inflammatory markers and systemic diseases.

Influence of Anxiety and Depression on Dental Caries

In addition to the behavioral factors that may reduce oral hygiene measures and result in poor adherence to professional dental

preventative measures in patients suffering with anxiety and depression,^{7,72,73} sympathetic stimulation in patients with anxiety and chronic stress has been associated with reduced salivary flow.^{74,75} Xerostomia is also a known side effect of many antidepressant and anxiolytic medications.¹⁰ Finally, simple carbohydrate intake is elevated in those individuals who are experiencing anxiety and depression.⁷⁶ All of these factors may combine to increase the risk of caries in individuals with depression and anxiety disorders.

A recent study found that in Finnish adults between 35-54 depression was significantly associated with the number of decayed teeth.⁸ This is consistent with previous studies that demonstrated that the severity of depression was associated with increased numbers of decayed teeth in adults.^{7,72,76} It is thus critical that oral health be considered an integral part of physical and psychological health and emphasis be placed upon collaboration between oral and medical healthcare professionals to result in ideal treatment outcomes. Further, in a study that evaluated oral health and depression, controlling for confounders such as alcohol and tobacco consumption, demographic data, mental health treatment, and systemic health, depression was found to be a risk factor for caries [OR 1.27 (1.13,1.44)], tooth loss [OR 1.31 (1.24, 1.37)], and total edentulism [OR 1.17 (1.02, 1.34)].⁷⁷

Implications for Anxiety and Depression Medications on Periodontal Disease Progression and Therapy

Stress and depression have been associated with negative treatment outcomes in patients with periodontal disease.^{4,5,26} Baseline levels of depression have been found to be predictive of the proportion of residual periodontal probing depths and tooth loss between 1 and 5 years post-treatment.⁷⁸ In patients receiving periodontal maintenance, occupational stress was associated with greater clinical attachment loss.⁷⁹ These studies indicate that psychosocial disorders should be identified and addressed prior to and during therapy to insure optimal treatment results.

Recent studies have indicated that antidepressant medications, particularly SSRIs and SNRIs, were associated with reduced levels of bleeding on probing (BOP) and clinical attachment loss in patients with depression and periodontitis.^{80,81} Furthermore, animal studies indicate that administration of antidepressant medications resulted in an amelioration of bone loss in animals with induced periodontitis exposed to stressful stimuli that made it approximately equivalent to those who were not exposed to the stressor.^{82,83} Given the overall underreporting, under-diagnosis, and under-treatment of anxiety disorders and depression, co-management of individuals with periodontitis and depression/anxiety by mental healthcare professionals and the dental team can insure best outcomes for patients.

Identification of At-risk Patients in the Dental Office: Screening and Referral

A readily available screening tool that allows dentists to assess level of depression is critical to identify the high number of individuals with undiagnosed mental disorders. A recent investigation demonstrated that dentists can effectively evaluate patients to identify both undiagnosed and undertreated depression with an eight question survey instrument, the Patient Health Questionnaire Depression Scale (PHQ8) (Figure 4).^{84,85} This may allow for dental healthcare practitioners to evaluate levels of depression in their patients and to refer patients to appropriate medical healthcare professionals. Furthermore, education and resources should be provided for individuals who are treating patients with diagnosed anxiety and depression so that they can discuss the oral implications of these diseases and the medications used to treat the diseases with patients to allow for continued monitoring and treatment of patients' oral health.⁸⁶

Summary

Poor oral health in individuals with common psychological problems remains a prevalent and unaddressed problem. Patients with anxiety, chronic stress, and depression experience increased risk for caries,



Personal Health Questionnaire Depression Scale (PHQ-8)

Over the **last 2 weeks**, how often have you been bothered by any of the following problems?
(circle **one** number on each line)

How often during the past 2 weeks were you bothered by...	Not at all	Several days	More than half the days	Nearly every day
1. Little interest or pleasure in doing things	0	1	2	3
2. Feeling down, depressed, or hopeless	0	1	2	3
3. Trouble falling or staying asleep, or sleeping too much	0	1	2	3
4. Feeling tired or having little energy.....	0	1	2	3
5. Poor appetite or overeating	0	1	2	3
6. Feeling bad about yourself, or that you are a failure, or have let yourself or your family down.....	0	1	2	3
7. Trouble concentrating on things, such as reading the newspaper or watching television.....	0	1	2	3
8. Moving or speaking so slowly that other people could have noticed. Or the opposite – being so fidgety or restless that you have been moving around a lot more than usual	0	1	2	3

Scoring

If two consecutive numbers are circled, score the higher (more distress) number. If the numbers are not consecutive, do not score the item. Score is the sum of the 8 items. If more than 1 item missing, set the value of the scale to missing. A score of 10 or greater is considered major depression, 20 or more is severe major depression.

Figure 4. A Simplified Screening Tool for Depression/Anxiety.⁸⁶

periodontal disease and tooth loss. These conditions can have a significant effect of overall quality of life. Painful, unesthetic dentition and missing teeth may worsen social isolation and low-self esteem as well as result in problems with mastication and phonetics. For individuals who are suffering from psychological illnesses, this could contribute to increased severity of their psychological symptoms. Given the multiple facets associated with the interaction of oral health and anxiety and depression, a multi-pronged, interdisciplinary approach should be applied.

Clinical Recommendations:

1. Physicians and mental healthcare providers should consider the effects of oral health within the focus on the physical health of patients with psychological illnesses. Identification of potential oral health issues, advice on dietary intake and oral hygiene, and early dental referral are critical to address and manage oral health in individuals with psychosocial illnesses.
2. Identification of individuals with depression and anxiety in the dental office and an

understanding of the effects that these conditions have on common oral health conditions is critical, given the stigma associated with mental health disorders and the large number of individuals who are undiagnosed, untreated, or undertreated.

3. Developing relationships with primary care providers and/or mental health professionals to allow for bidirectional referrals to optimize physical and mental wellness of patients is critical.
4. Risk assessment for caries and periodontal disease should be undertaken to allow dental healthcare and medical professionals to discuss artificial salivary products, administration of high fluoride dentifrices and gels, and increased dental maintenance intervals based upon the patients' individualized risks and conditions.
5. Additional recommendations to reduce overall levels of stress in patients with depression and anxiety may include addressing coping mechanisms, increased social networks, and physical exercise and should be reinforced by both medical and oral healthcare providers.

Course Test Preview

To receive Continuing Education credit for this course, you must complete the online test. Please go to: www.dentalcare.com/en-us/ce-courses/ce573/test

- 1. Approximately how many individuals live with depression worldwide?**
 - A. 102 million
 - B. 257 million
 - C. 332 million
 - D. 549 million
- 2. Anxiety and depression are often comorbid. What percentage of individuals diagnosed with depression also meet the criteria for an anxiety disorder?**
 - A. 20%
 - B. 40%
 - C. 60%
 - D. 80%
- 3. Individuals with anxiety and depression have higher rates of many systemic diseases. Which of the following has NOT been shown to be higher in individuals with psychosocial disorders?**
 - A. Diabetes Mellitus
 - B. Autoimmune Disorders
 - C. Cardiovascular Disease
 - D. All of the above are linked to depression and anxiety.
- 4. Periodontitis and anxiety/depressive disorders have been linked. What is thought to be the underlying etiology of this association?**
 - A. Increased inflammatory burden through a stress mechanism
 - B. Decreased self-care, including oral hygiene
 - C. Increased xerostomia through sympathetic nerve activation and/or antidepressive medications
 - D. All of the above have been thought to play a role.
- 5. Stress signals are transmitted through the limbic-hypothalamic-pituitary-adrenal axis to the CNS via all of the following neurotransmitters EXCEPT:**
 - A. Serotonin
 - B. Stress hormones
 - C. Neuropeptide Y
 - D. Monoamines
- 6. Stress, in particular chronic stress, may increase depressive symptoms through which of the following mechanisms?**
 - A. Increase in deleterious habits and a concomitant decrease in overall self-esteem
 - B. Stress induces a systemic hyperinflammatory state which may, in susceptible individuals, lead to depression
 - C. Poor coping mechanisms may worsen both stress and depression
 - D. All of the above have been purported to cause the interaction between stress and depression.
- 7. Studies have shown that depression is associated with higher rates of caries in patients 35-54.**
 - A. True
 - B. False

- 8. In patients undergoing active periodontal maintenance therapy, baseline levels of depression were predictive of increased BOP and probing depths between 1 and 5 years after initial therapy. Occupational stress was not associated with increased clinical attachment loss over 5 years.**
- A. Both statements are true.
 - B. Both statements are false.
 - C. The first statement is true, the second statement is false.
 - D. The first statement is false, the second statement is true.
- 9. In both animal and human models, antidepressant use was associated with decreased clinical attachment loss, bone loss, and BOP.**
- A. True
 - B. False
- 10. Assessment of mental health disorders in the dental office allows for a fuller understanding of overall risk factors. Which of the following is a tool that can be used to assess depressive symptoms in the dental office?**
- A. PHQ8
 - B. SPQR
 - C. NEWT
 - D. OWL
- 11. Interdisciplinary care for individuals with depression and anxiety disorders is critical. If a patient is diagnosed with depression and/or anxiety, which of the following should be undertaken by his/her medical healthcare provider to improve oral health outcomes?**
- A. Discussion of the interaction between psychosocial disorders and oral health
 - B. Discussion of possible oral side effects of anti-depressive medications and mechanisms to mitigate their deleterious effects
 - C. Early referral for dental care
 - D. All of the above.

References

1. Ritchie H, Roser Max. Mental Health. OurWorldinData.org. 2018 Apr. Accessed November 11, 2022.
2. Pratt LA, Brody DJ. Depression in the US Household Population, 2009-2012. CDC. NCHS Data Brief No. 172, 2014 Dec. November 11, 2022.
3. Cademartori MG1, Gastal MT1, Nascimento GG1,. Is dls depression associated with oral health outcomes in adults and elders? A systematic review and meta-analysis.Clin Oral Investig. 2018 Nov;22(8):2685-2702. doi: 10.1007/s00784-018-2611-y. Epub 2018 Sep 6.
4. Genco RJ, Ho AW, Grossi SG, et al. Relationship of stress, distress and inadequate coping behaviors to periodontal disease. J Periodontol. 1999 Jul;70(7):711-23. doi: 10.1902/jop.1999.70.7.711.
5. Warren KR, Postolache TT, Groer ME, et al. Role of chronic stress and depression in periodontal diseases. Periodontol 2000. 2014 Feb;64(1):127-38. doi: 10.1111/prd.12036.
6. Hwang SH, Park SG. The relationship between depression and periodontal diseases. Community Dent Health. 2018 Mar 1;35(1):23-29. doi: 10.1922/CDH_4150Hwang07.7.
7. McFarland M, Ingelhart M. Depression, self efficacy, and oral health – an exploration. Oral Health Dent Manag 2010; 9: 214-222. November 11, 2022.
8. Delgado-Angulo EK, Sabbah W, Suominen AL, et al. The association of depression and anxiety with dental caries and periodontal disease among Finnish adults. Community Dent Oral Epidemiol. 2015 Dec;43(6):540-9. doi: 10.1111/cdoe.12179. Epub 2015 Jul 1.
9. Muniz FWMG, Melo IM, Rösing CK, et al. Use of antidepressive agents as a possibility in the management of periodontal diseases: A systematic review of experimental studies. J Investig Clin Dent. 2018 Feb;9(1). doi: 10.1111/jicd.12291. Epub 2017 Sep 1.
10. Ciancio SG. Medications' impact on oral health. J Am Dent Assoc. 2004 Oct;135(10):1440-8; quiz 1468-9. doi: 10.14219/jada.archive.2004.0055.
11. Kessler RC, Berglund P, Demler O, et al. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. Arch Gen Psychiatry. 2005 Jun;62(6):593-602. doi: 10.1001/archpsyc.62.6.593.
12. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition. Arlington, VA. American Psychiatric Publishing. 2013.
13. Kaufman J, Charney D. Comorbidity of mood and anxiety disorders. Depress Anxiety. 2000;12 Suppl 1:69-76. doi: 10.1002/1520-6394(2000)12:1+<69::AID-DA9>3.0.CO;2-K.
14. Hsu WY, Tsai HJ, Yu SH, et al. Association of Periodontitis and Subsequent Depression: A Nationwide Population-Based Study. Medicine (Baltimore). 2016 Aug;95(31):e4419. doi: 10.1097/MD.0000000000004419.
15. Rugulies R. Depression as a predictor for coronary heart disease. A review and meta-analysis. Am J Prev Med. 2002 Jul;23(1):51-61. doi: 10.1016/s0749-3797(02)00439-7.
16. Sullivan MD, O'Connor P, Feeney P, et al. Depression predicts all-cause mortality: epidemiological evaluation from AACORD HRQL substudy. Diabetes Care. 2012 Aug;35(8):1708-15. doi: 10.2337/dc11-1791. Epub 2012 May 22.
17. Wulsin LR, Valliant GE, Wells VE. A systematic review of the mortality of depression. Psychosom Med. 1999 Jan-Feb;61(1):6-17. doi: 10.1097/00006842-199901000-00003.
18. Lawrence D, Hancock KJ, Kisely S. The gap in life expectancy from preventable physical illness in psychiatric patients in Western Australia: retrospective analysis of population based registers. BMJ. 2013 May 21;346:f2539. doi: 10.1136/bmj.f2539.
19. Banasr M, Valentine GW, Li XY, et al. Chronic unpredictable stress decreases cell proliferation in the cerebral cortex of the adult rat. Biol Psychiatry. 2007 Sep 1;62(5):496-504. Epub 2007 Jun 21. doi: 10.1016/j.biopsych.2007.02.006.
20. Duman RS, Monteggia LM. A neurotropic model for stress-related mood disorders. Biol Psychiatry. 2006 Jun 15;59(12):1116-27. Epub 2006 Apr 21.

21. Duman RS, Voleti B. Signaling pathways underlying the pathophysiology and treatment of depression: novel mechanisms for rapid-acting agents. *Trends Neurosci.* 2012 Jan;35(1):47-56. doi: 10.1016/j.tins.2011.11.004.
22. Krishnan V, Nestler EJ. The molecular neurobiology of depression. *Nature.* 2008 Oct 16;455(7215):894-902. doi: 10.1038/nature07455.
23. McEwen BS. Central effects of stress hormones in health and disease: understanding the protective and damaging effects of stress and stress mediators. *Eur J Pharmacol.* 2008 Apr 7;583(2-3):174-85. doi: 10.1016/j.ejphar.2007.11.071. Epub 2008 Jan 30.
24. Shansky RM, Morrison JH. Stress-induced dendritic remodeling in the medial prefrontal cortex: effects of circuit, hormones, and rest. *Brain Res.* 2009 Oct 13;1293:108-13. doi: 10.1016/j.brainres.2009.03.062. Epub 2009 Apr 8.
25. Koob GF. Stress, corticotropin-releasing factor and drug addiction. *Ann N Y Acad Sci.* 1999;897:27-45. doi: 10.1111/j.1749-6632.1999.tb07876.x.
26. Liu F, Wen YF, Zhou Y, et al. A meta-analysis of emotional disorders as possible risk factors for chronic periodontitis. *Medicine (Baltimore).* 2018 Jul;97(28):e11434. doi: 10.1097/MD.00000000000011434.
27. Baxter AJ, Scott KM, Vos T, et al. Global prevalence of anxiety disorders: a systematic review and meta-regression. *Psychol Med.* 2013 May;43(5):897-910. doi: 10.1017/S003329171200147X. Epub 2012 Jul 10.
28. World Health Organization. Depression and other common mental disorders: Global health estimates. Accessed November 11, 2022.
29. United States: Institute for Health Metrics and Evaluation (IHME), 2018. Global Burden of Disease Collaborative Network. Global Burden of Disease Study 2017 (GBD 2017) Results. Seattle, WA. Accessed November 11, 2022.
30. Bandelow B, Michaelis S, Wedekind D. Treatment of anxiety disorders. *Dialogues Clin Neurosci.* 2017 Jun;19(2):93-107.
31. Thibaut F. Anxiety disorders: a review of current literature. *Dialogues Clin Neurosci.* 2017 Jun;19(2):87-88.
32. Musazzi L, Tornese P, Sala N, et al. Acute or Chronic? A Stressful Question. *Trends Neurosci.* 2017 Sep;40(9):525-535. doi: 10.1016/j.tins.2017.07.002. Epub 2017 Aug 1.
33. Marshall RD, Garakani A. Psychobiology of the acute stress response and its relationship to the psychobiology of post-traumatic stress disorder. *Psychiatr Clin North Am.* 2002 Jun;25(2):385-95. doi: 10.1016/s0193-953x(01)00005-3.
34. McEwen BS. Central effects of stress hormones in health and disease: Understanding the protective and damaging effects of stress and stress mediators. *Eur J Pharmacol.* 2008 Apr 7;583(2-3):174-85. doi: 10.1016/j.ejphar.2007.11.071. Epub 2008 Jan 30.
35. Sterlemann V, Ganea K, Liebl C, et al. Long-term behavioral and neuroendocrine alterations following chronic social stress in mice: implications for stress-related disorders. *Horm Behav.* 2008 Feb;53(2):386-94. Epub 2007 Nov 19. doi: 10.1016/j.yhbeh.2007.11.001.
36. McEwen BS. Neurobiological and Systemic Effects of Chronic Stress. *Chronic Stress (Thousand Oaks).* 2017 Jan-Dec;1. doi: 10.1177/2470547017692328. Epub 2017 Apr 10.
37. American Psychiatric Association. Anxiety Disorders. What Are Anxiety Disorders? Accessed November 11, 2022.
38. National Institute of Mental Health. Anxiety Disorders: Overview. Accessed November 11, 2022.
39. National Institute of Mental Health. Major Depression: Definitions. Accessed November 11, 2022.
40. Kessler RC, Bromet EJ. The epidemiology of depression across cultures. *Annu Rev Public Health.* 2013;34:119-38. doi: 10.1146/annurev-publhealth-031912-114409.
41. Centers for Disease Control and Prevention. Mental Health Conditions: Depression and Anxiety. Accessed November 11, 2022.
42. National Alliance on Mental Illness (NAMI). Depression: Treatment. Accessed November 11, 2022.

43. Lu XT, Zhao YX, Zhang Y, et al. Psychological stress, vascular inflammation and atherogenesis: potential roles of circulating cytokines. *J Cardiovasc Pharmacol*. 2013 Jul;62(1):6-12. doi: 10.1097/FJC.0b013e3182858fac.
44. Kaufman J, Charney D. Comorbidity of mood and anxiety disorders. *Depress Anxiety* 2000; 12(Suppl 1): 69-76. doi: 10.1002/1520-6394(2000)12:1+<69::AID-DA9>3.0.CO;2-K.
45. Strarr LR, Hammen C, Connolly NP, et al. Does relational dysfunction mediate the association between anxiety disorders and later depression? Test an interpersonal model of comorbidity. *Depress Anxiety*. 2014 Jan;31(1):77-86. doi: 10.1002/da.22172. Epub 2013 Aug 26.
46. Wittchen HU, Beesdo K, Bittner A, et al. Depressive episodes—evidence for a causal role of primary anxiety disorders? *Eur Psychiatry*. 2003 Dec;18(8):384-93. doi: 10.1016/j.eurpsy.2003.10.001.
47. Howren MB, Lamkin DM, Suls J. Associations of depression with C-reactive protein, IL-1 and IL-6: a meta-analysis. *Psychosom Med*. 2009 Feb;71(2):171-86. doi: 10.1097/PSY.0b013e3181907c1b. Epub 2009 Feb 2.
48. Singh T, Newman AB. Inflammatory markers in population studies of aging. *Ageing Res Rev*. 2011 Jul;10(3):319-29. doi: 10.1016/j.arr.2010.11.002. Epub 2010 Dec 8.
49. Hannestad J, DellaGioia N, Bloch M. The effect of antidepressant medication treatment on serum levels of inflammatory cytokines: a meta-analysis. *Neuropsychopharmacology*. 2011 Nov;36(12):2452-9. doi: 10.1038/npp.2011.132. Epub 2011 Jul 27.
50. Marshall EC, Zvolensky MJ, Vujanovic AA, et al. Evaluation of smoking characteristics among community-recruited daily smokers with and without posttraumatic stress disorder and panic psychopathology. *J Anxiety Disord*. 2008 Oct;22(7):1214-26. doi: 10.1016/j.janxdis.2008.01.003. Epub 2008 Jan 11.
51. Lawrence D, Hancock KJ, Kisely S. The gap in life expectancy from preventable physical illness in psychiatric patients in Western Australia: retrospective analysis of population-based registers. *BMJ*. 2013 May 21;346:f2539. doi: 10.1136/bmj.f2539.
52. Lawrence D, Kisley S, Pais J. The epidemiology of excess mortality in people with mental illness. *Can J Psychiatry*. 2010 Dec;55(12):752-60. doi: 10.1177/070674371005501202.
53. Kisley S, Sawyer E, Siskind D, et al. The oral health of people with anxiety and depressive disorders – a systematic review and meta-analysis. *J Affect Disord*. 2016 Aug;200:119-32. doi: 10.1016/j.jad.2016.04.040. Epub 2016 Apr 21.
54. Okoro, Catherine A.; Strine, Tara W.; Eke, Paul I.; Dhingra, Satvinder S.; and Balluz, Lina S., The association between depression and anxiety and use of oral health services and tooth loss. *Public Health Resources* 2012; 264.
55. Longley AJ, Doyle PE. Mental illness and the dental patient. *J Dent Hyg* 2003; 77(3):190-204; quiz 204-6. Erratum in: *J Dent Hyg* 2004; 78(1): 4.
56. The Cleveland Clinic Dental Phobia. Accessed November 11, 2022.
57. Matevosyan NR. Oral health of adults with serious mental illness: A review. *Comm Ment Health J*. 2009
58. Lam PD, John DA, Galfalvy H, Kunzel C, Lewis-Fernandez R. Oral Health-Related Quality of Life Among Publicly Insured Mental Health Service Outpatients with Serious Mental Illness. *Psych Services* 2019; 70(12): 1101-1109.
59. Freeman R, Goss S. Stress measures as predictors of periodontal disease--a preliminary communication. *Community Dent Oral Epidemiol*. 1993 Jun;21(3):176-7. doi: 10.1111/j.1600-0528.1993.tb00748.x.
60. Gree LW, Tryon WW, Marks B, et al. Periodontal disease as a function of life events stress. *J Human Stress*. 1986 Spring;12(1):32-6. doi: 10.1080/0097840X.1986.9936764.
61. Mannem S, Chava VK. The effect of stress on periodontitis: A clinicobiochemical study. *J Indian Soc Periodontol*. 2012 Jul;16(3):365-9. doi: 10.4103/0972-124X.100912.
62. Rai B, Kaur J, Anand SC, et al. Salivary stress markers, stress, and periodontitis: a pilot study. *J Periodontol*. 2011 Feb;82(2):287-92. doi: 10.1902/jop.2010.100319. Epub 2010 Aug 19.

63. Peruzzo DC, Benatti BB, Ambrosano GM, et al. A systematic review of stress and psychological factors as possible risk factors for periodontal disease. *J Periodontol*. 2007 Aug;78(8):1491-504. doi: 10.1902/jop.2007.060371.
64. Nascimento GG, Gastal MT, Leite FRM, et al. Is there an association between depression and periodontitis? A birth cohort study? *J Clin Periodontol*. 2019 Jan;46(1):31-39. doi: 10.1111/jcpe.13039.
65. Khambaty T, Steward JC. Associations of depressive and anxiety disorders with periodontal disease prevalence in young adults: analysis of 1999-2004 National Health and Nutrition Examination Survey (NHANES) Data. *Ann Behav Med*. 2013 Jun;45(3):393-7. doi: 10.1007/s12160-013-9471-0.
66. Kurushima Y, Bowyer R, Ide M, et al. Genetic and environmental contributions to the association between mood disorder and periodontal disease: A cross-sectional study among female twins in the UK. *J Clin Periodontol*. 2019 Jan;46(1):40-50. doi: 10.1111/jcpe.13045.
67. Shannon IL, Kilgore WG, O'Leary TJ. Stress as a predisposing factor in necrotizing ulcerative gingivitis. *J Periodontol*. 1969 Apr;40(4):240-2. doi: 10.1902/jop.1969.40.4.240.
68. Shields WD. Acute necrotizing ulcerative gingivitis. A study of some of the contributing factors and their validity in an Army population. *J Periodontol*. 1977 Jun;48(6):346-9. doi: 10.1902/jop.1977.48.6.346.
69. Maupin CC, Bell WB. The relationship of 17-hydroxycorticosteroid to acute necrotizing ulcerative gingivitis. *J Periodontol*. 1975 Dec;46(12):721-2. doi: 10.1902/jop.1975.46.12.721.
70. Davies RM, Smith RG, Porter SR. Destructive forms of periodontal disease in adolescents and young adults. *Br Dent J*. 1985 Jun 22;158(12):429-36. doi: 10.1038/sj.bdj.4805622.
71. Page RC, Altman LC, Ebersole JL, et al. Rapidly progressive periodontitis. A distinct clinical condition. *J Periodontol*. 1983 Apr;54(4):197-209.
72. Anttila S, Knuuttila M, Ylostalo P, et al. Symptoms of depression and anxiety in relation to dental health behavior and self-perceived dental treatment need. *Eur J Oral Sci*. 2006 Apr;114(2):109-14. doi: 10.1111/j.1600-0722.2006.00334.x.
73. Kurer JR, Watts TL, Weinman J, et al. Psychological mood of regular dental attenders in relation to oral hygiene behaviour and gingival health. *J Clin Periodontol*. 1995 Jan;22(1):52-5. doi: 10.1111/j.1600-051x.1995.tb01770.x.
74. Boyapati L, Wang HL. The role of stress in periodontal disease and wound healing. *Periodontol 2000*. 2007;44:195-210. doi: 10.1111/j.1600-0757.2007.00211.x.
75. Anttila SS, Knuuttila ML, Sakki TK. Depressive symptoms favor abundant growth of salivary lactobacilli. *Psychosom Med*. 1999 Jul-Aug;61(4):508-12. doi: 10.1097/00006842-199907000-00015.
76. Anttila SS, Knuuttila ML, Sakki TK. Relationship of depressive symptoms to edentulousness, dental health, and dental health behavior. *Acta Odontol Scand*. 2001 Dec;59(6):406-12. doi: 10.1080/000163501317153275.
77. Cademartori MG, Gastal MT, Nascimento GG, Demarco FF, Correa MB. Is depression associated with oral health outcomes in adults and elders? A systematic review and meta-analysis. *Clin Oral Invest* 2018; 22(8): 2685-2702.
78. Elter JR, White BA, Gaynes BN, et al. Relationship of clinical depression to periodontal treatment outcome. *J Periodontol*. 2002 Apr;73(4):441-9. doi: 10.1902/jop.2002.73.4.441.
79. Kamma JJ, Baehni PC. Five-year maintenance follow-up of early-onset periodontitis patients. *J Clin Periodontol*. 2003 Jun;30(6):562-72. doi: 10.1034/j.1600-051x.2003.00289.x.
80. Bhatia A, Sharma RK, Tewari S, et al. Effect of fluoxetine on periodontal status in patients with depression: a cross-sectional observational study. *J Periodontol*. 2015 Aug;86(8):927-35. doi: 10.1902/jop.2015.140706. Epub 2015 Mar 27.
81. Bhatia A, Sharma RK, Tewari S, et al. Periodontal status in chronic periodontitis depressed patients on desvenlafaxine: An observational study. *J Indian Soc Periodontol*. 2018 Sep-Oct;22(5):442-446. doi: 10.4103/jisp.jisp_219_18.

82. Anguiar JCA, Gomes EPP, Fonseca-Silva T, et al. Fluoxetine reduces periodontal disease progression in conditioned fear stress model in rats. *J Periodontal Res.* 2013 Oct;48(5):632-7. doi: 10.1111/jre.12049. Epub 2013 Feb 21.
83. Branco-de-Almeida LS, Franco GC, Castro ML, et al. Fluoxetine inhibits inflammatory response and bone loss in a rat model of ligature-induced periodontitis. *J Periodontol.* 2012 May;83(5):664-71. doi: 10.1902/jop.2011.110370. Epub 2011 Oct 3.
84. Kroenke K, Strine TW, Spitzer RL, et al. The PHQ-8 as a measure of current depression in the general population. *J Affect Disord.* 2009 Apr;114(1-3):163-73. doi: 10.1016/j.jad.2008.06.026. Epub 2008 Aug 27.
85. Villalobos Sancho SM. Effect of antidepressants on inflammatory markers in gingival tissue and serum samples in patients with chronic periodontitis. A Thesis. University of Alabama at Birmingham. Accessed November 11, 2022.
86. Self Management Resource Center (Open Source). Personal Health Questionnaire Depression Scale (PHQ-8). Accessed November 11, 2022.

Additional Resources

- No Additional Resources Available.

About the Author

Maria L. Geisinger, DDS, MS



Mia L. Geisinger, DDS, MS is a Professor and Director of Advanced Education in Periodontology in the Department of Periodontology in the University of Alabama at Birmingham (UAB) School of Dentistry. Dr. Geisinger received her BS in Biology from Duke University, her DDS from Columbia University School of Dental Medicine, and her MS and Certificate in Periodontology and Implantology from the University of Texas Health Science Center at San Antonio. Dr. Geisinger is a Diplomate in the American Board of Periodontology and a Fellow in the International Team for Implantology. She has served as the

President of the American Academy of Periodontology Foundation, as the Chair of the American Dental Association's Council on Scientific Affairs, and on multiple national and regional organized dentistry committees. She currently serves as the Vice President of the AAP, as a Board member for the ADA Science and Research Institute, and on numerous AAP and ADA committees and task forces. She has authored over 60 peer-reviewed publications and her research interests include periodontal and systemic disease interaction, implant dentistry in the periodontally compromised dentition, and novel treatment strategies for oral soft and hard tissue regeneration. She lectures nationally and internationally on topics in periodontology and oral healthcare.

Email: miagdds@uab.edu