

Lung cancer



Average chemotherapy use by sex (2012–2017):
stages 1–3 or local and regional



Australia

New South Wales



0 20% 40% 60% 80% 100%



Canada

Alberta



Atlantic Canada



British Columbia



Ontario



Saskatchewan-Manitoba



0% 20% 40% 60% 80% 100%



Norway

Norway



0 20% 40% 60% 80% 100%



UK

England



Northern Ireland



Scotland



Wales



0% 20% 40% 60% 80% 100%



TNM and SEER stages do not always align: UK and Canada used TNM staging, Norway and Australia used SEER.

Atlantic Canada includes Nova Scotia and Prince Edward Island.

Some jurisdictions had a large % of missing stage data which could introduce bias: Norway (12%); New South Wales (14%).

Lung cancer




Average radiotherapy use by sex (2012–2017):
stages 1–3 or local and regional



Australia

New South Wales



Canada

Alberta



Atlantic Canada



British Columbia



Ontario



Saskatchewan-Manitoba



Norway

Norway



UK

England



Northern Ireland



Scotland



Wales



TNM and SEER stages do not always align: UK and Canada used TNM staging, Norway and Australia used SEER.

Atlantic Canada includes Newfoundland & Labrador, New Brunswick, Nova Scotia and Prince Edward Island.

Some jurisdictions had a large % of missing stage data which could introduce bias: Northern Ireland (10%), Norway (12%), New South Wales (14%).

Lung cancer




Average chemotherapy use by sex (2012–2017):
stage 4 or distant




Australia

New South Wales




Canada

Alberta
Atlantic Canada
British Columbia
Ontario
Saskatchewan-Manitoba




Norway

Norway



UK

England
Northern Ireland
Scotland
Wales



Female
Male

TNM and SEER stages do not always align: UK and Canada used TNM staging, Norway and Australia used SEER.

Atlantic Canada includes Nova Scotia and Prince Edward Island.

Some jurisdictions had a large % of missing stage data which could introduce bias: Norway (12%); New South Wales (14%).

Lung cancer




Average radiotherapy use by sex (2012–2017):
stage 4 or distant



Australia

New South Wales



Canada

Alberta



Atlantic Canada



British Columbia



Ontario



Saskatchewan-Manitoba



0% 20% 40% 60% 80% 100%



Norway

Norway



0 20% 40% 60% 80% 100%



UK

England



Northern Ireland



Scotland



Wales



0% 20% 40% 60% 80% 100%



TNM and SEER stages do not always align: UK and Canada used TNM staging, Norway and Australia used SEER.

Atlantic Canada includes Newfoundland & Labrador, New Brunswick, Nova Scotia and Prince Edward Island.

Some jurisdictions had a large % of missing stage data which could introduce bias: Northern Ireland (10%), Norway (12%), New South Wales (14%).