PATIENT AGREEMENT TO SYSTEMIC ANTICANCER THERAPY: Gemcitabine

A A A I P I WILL A I WILL I P	•
CANCER THERAPY:	PATIENT'S FIRST NAME(S):
Gemcitabine	DATE OF BIRTH:
	NHS NUMBER:(or other identifier)
HOSPITAL NAME/STAMP:	☐ MALE ☐ FEMALE
	SPECIAL REQUIREMENTS: (e.g. other language/other communication method)
RESPONSIBLE HEALTH PROFESSIONAL:	
Name:	
Job title:	
NAME OF PROPOSED COURSE OF TREATME	NT (include brief explanation if medical term not clear)
Gemcitabine for the treatment of cutaneous lymparts.	
☐ Given intravenously on day 1, 8 and 15, every 28 c	
WHERE THE TREATMENT WILL BE GIVEN:	
☐ outpatient ☐ day unit/case ☐ inpatient	Other:
CTATEMENT OF LIFALTLI BROSE	ECCIONIAL
STATEMENT OF HEALTH PROFI appropriate knowledge of proposed procedure, as specified in t	ESSIGNAL (to be filled in by health professional with the hospital/Trust's consent policy)
I have explained the procedure/treatment to the patie	
☑ all relevant boxes	
THE INTENDED BENEFITS	
\square CURATIVE – to give you the best possible chance o	
■ DISEASE CONTROL/PALLIATIVE – the aim is not to The aim is to improve both quality of life and surviva	

☐ **ADJUVANT** – therapy given after surgery to reduce the risk of the cancer coming back.

□ **NEO-ADJUVANT** – therapy given before surgery/radiotherapy to shrink the cancer, allow radical treatment

PATIENT DETAILS

PATIENT'S SURNAME/FAMILY NAME:

and reduce the risk of the cancer coming back.

STATEMENT OF HEALTH PROFESSIONAL (continued)

Patient ide	entifier/label		

SIGNIFICANT, UNAVOIDABLE OR FREQUENTLY OCCURRING RISKS

(vomiting), temporary changes in the way the liver works, flu-like symptoms, fluid retention (you may notice you gain weight and/or your face, ankles and legs swell), breathlessness, skin rash, thinning of the hair and small amounts of blood or protein in the urine. An increased risk of getting an infection from a drop in white blood cells - it is harder to fight infections and you can become very ill. If you have a severe infection this can be life threatening. Contact you doctor or hospital straight away if: • your temperature goes over 37.5°C (99.5°F) or over 38°C (100.4°F), depending on the advice given by your chemotherapy team • you suddenly feel unwell (even with a normal temperature) OCCASIONAL SIDE EFFECTS: Between 1 and 10 in every 100 (1-10%) people have one or more of these effects: Diarrhoea, constipation, drowsiness, sore mouth and ulcers, loss of appetite, muscle and back pain, sweating, runny nose, and itchy skin. There may be a risk of tumour lysis syndrome in the presence of a high tumour burden. Symptoms are caused by the breakdown of cancer cells when treatment is started, causing changes in blood electrolytes. Patients at a high risk will be prescribed medicines to reduce the risk of complications.	COMMON SIDE EFFECTS:	OTI	HER RISKS:			
(you may notice you gain weight and/or your face, ankles and legs swell), breathlessness, skin rash, thinning of the hair and small amounts of blood or protein in the urine. ☐ An increased risk of getting an infection from a drop in white blood cells - it is harder to fight infections and you can become very ill. If you have a severe infection this can be life threatening. Contact you doctor or hospital straight away if: • your temperature goes over 37.5°C (99.5°F) or over 38°C (100.4°F), depending on the advice given by your chemotherapy team • you suddenly feel unwell (even with a normal temperature) OCCASIONAL SIDE EFFECTS: Between 1 and 10 in every 100 (1-10%) people have one or more of these effects: ☐ Diarrhoea, constipation, drowsiness, sore mouth and ulcers, loss of appetite, muscle and back pain, sweating, runny nose, and itchy skin. ☐ There may be a risk of tumour lysis syndrome in the presence of a high tumour burden. Symptoms are caused by the breakdown of cancer cells when treatment is started, causing changes in blood electrolyses. Patients at a high risk will be prescribed medicines to reduce the risk of complications. ☐ We provide the vein whilst being given; this is called extravasation. It can damage the tissue aroun the vein. Tell the nurse if you have any stingin pain, redness or swelling around the vein. Ell the nurse if you have any stingin pain, redness or swelling around the vein. Ell the nurse if you have any stingin pain, redness or swelling around the vein. Ell the nurse if you have any stingin pain, redness or swelling around the vein. Ell the nurse if you have any stingin pain, redness or swelling around the vein. Ell the nurse if you have any stingin pain, redness or swelling around the vein. Ell the nurse if you have any stingin pain, redness or swelling around the vein. Ell the nurse if you have any stingin pain, redness or swelling around the vein. Ell the nurse if you have any stingin pain, redness or swelling around the vein. Elt the suit pain, redness or swelling ar	More than 10 in every 100 (>10%) people have one or more of the side effects listed: Anaemia (low number of red blood cells), tiredness and feeling weak (fatigue), bruising or bleeding, feeling sick (nausea) and being sick (vomiting), temporary changes in the way the liver works, flu-like symptoms, fluid retention (you may notice you gain weight and/or your face, ankles and legs swell), breathlessness, skin rash, thinning of the hair and small amounts of blood or protein in the urine. An increased risk of getting an infection from a drop in white blood cells - it is harder to fight infections and you can become very ill. If you have a severe infection this can be life threatening. Contact you doctor or hospital straight away if: your temperature goes over 37.5°C (99.5°F) or over 38°C (100.4°F), depending on the advice given by your chemotherapy team you suddenly feel unwell (even with a normal temperature)		Rare side effects include allergic reactions, skin rash on areas previously exposed to radiotherapy, changes in the lungs, changes in the way the heart works, and haemolytic uraemic syndrome (causing anaemia, kidney			
 An increased risk of getting an infection from a drop in white blood cells - it is harder to fight infections and you can become very ill. If you have a severe infection this can be life threatening. Contact you doctor or hospital straight away if: your temperature goes over 37.5°C (99.5°F) or over 38°C (100.4°F), depending on the advice given by your chemotherapy team you suddenly feel unwell (even with a normal temperature) OCCASIONAL SIDE EFFECTS: Between 1 and 10 in every 100 (1-10%) people have one or more of these effects: Diarrhoea, constipation, drowsiness, sore mouth and ulcers, loss of appetite, muscle and back pain, sweating, runny nose, and itchy skin. There may be a risk of tumour lysis syndrome in the presence of a high tumour burden. Symptoms are caused by the breakdown of cancer cells when treatment is started, causing changes in blood electrolytes. Patients at a high risk will be prescribed medicines to reduce the risk of complications. Extravasation is not common but if it happen important that it's dealt with quickly. Potential side-effects with the anti-sickness medication may include: constipation, headaches, indigestion, difficulty sleeping, an agitation. Cancer can increase your risk of developing a blood clot (thrombosis), and having treatment with anti-cancer medicines may increase this further. A blood clot may cause pain, redness swelling in a leg, or breathlessness and chest you must tell your doctor straight away if you have any of these symptoms. Some anti-cancer medicines can damage women's ovaries and men's sperm. This may lead to infertility in men and women and/or end the development of a baby in the womb. It is important to to become pregnant or father a child while you are having treatme			of the vein whilst being given; this is called extravasation. It can damage the tissue around the vein. Tell the nurse if you have any stinging,			
threatening. Contact you doctor or hospital straight away if: • your temperature goes over 37.5°C (99.5°F) or over 38°C (100.4°F), depending on the advice given by your chemotherapy team • you suddenly feel unwell (even with a normal temperature) OCCASIONAL SIDE EFFECTS: Between 1 and 10 in every 100 (1-10%) people have one or more of these effects: Diarrhoea, constipation, drowsiness, sore mouth and ulcers, loss of appetite, muscle and back pain, sweating, runny nose, and itchy skin. There may be a risk of tumour lysis syndrome in the presence of a high tumour burden. Symptoms are caused by the breakdown of cancer cells when treatment is started, causing changes in blood electrolytes. Patients at a high risk will be prescribed medicines to reduce the risk of complications. Headaches, indigestion, difficulty sleeping, an agitation. Cancer can increase your risk of developing a blood clot (thrombosis), and having treatment with an anti-cancer medicines may increase this further. A blood clot thrombosis), and having treatment with an treatment is eleounce medicines can deavelope may be a risk of leveloping and blood clot (thrombosis), and having treatment with an treatment is eleounce medicines may increase this further. A blood clot (thrombosis), and having treatment with an anti-cancer medicines may increase this further. A blood clot (thrombosis), and having treatment with an anti-cancer medicines may increase this further. A blood clot (thrombosis), and having treatment with an anti-cancer medicines may increase this further. A blood clot (thrombosis), and having treatment with an anti-cancer medicines and chest you must tell your doctor straight away if you have any of these symptoms. Some anti-cancer medicines may damage the development of a baby in the womb. It is important not to become pregnant or father a child while you are having treatment and for 6 months afterwards. It is important to use effective contraception during and for 6 months after treatment. You can talk to your doctor or nurse a			Extravasation is not common but if it happens it's important that it's dealt with quickly.			
over 38°C (100.4°F), depending on the advice given by your chemotherapy team • you suddenly feel unwell (even with a normal temperature) OCCASIONAL SIDE EFFECTS: Between 1 and 10 in every 100 (1-10%) people have one or more of these effects: Diarrhoea, constipation, drowsiness, sore mouth and ulcers, loss of appetite, muscle and back pain, sweating, runny nose, and itchy skin. There may be a risk of tumour lysis syndrome in the presence of a high tumour burden. Symptoms are caused by the breakdown of cancer cells when treatment is started, causing changes in blood electrolytes. Patients at a high risk will be prescribed medicines to reduce the risk of complications. blood clot (thrombosis), and having treatment with anti-cancer medicines may increase this further. A blood clot may cause pain, redness swelling in a leg, or breathlessness and chest you must tell your doctor straight away if you have any of these symptoms. Some anti-cancer medicines can damage women's ovaries and men's sperm. This may lead to infertility in men and women and/or emenopause in women. Some anti-cancer medicines may increase this further. A blood clot (thrombosis), and having treatment with anti-cancer medicines may chest you must tell your doctor straight away if you have any of these symptoms. Some anti-cancer medicines may damage the development of a baby in the womb. It is important not to become pregnant or father a child while you are having treatment and for 6 months afterwards. It is important to use effective contraception during and for 6 morths afterwards. It is important to use effective contraception during and for 6 morths afterwards. It is important to use effective contraception during and for 6 morths afterwards. It is important to use effective contraception during and for 6 morths afterwards. It is important to use effective contraception during and for 6 morths afterwards. It is important to use effective contraception during and for 6 morths afterwards. It is important to use effective contraception durin			headaches, indigestion, difficulty sleeping, and			
 temperature) OCCASIONAL SIDE EFFECTS: Between 1 and 10 in every 100 (1-10%) people have one or more of these effects: Diarrhoea, constipation, drowsiness, sore mouth and ulcers, loss of appetite, muscle and back pain, sweating, runny nose, and itchy skin. There may be a risk of tumour lysis syndrome in the presence of a high tumour burden. Symptoms are caused by the breakdown of cancer cells when treatment is started, causing changes in blood electrolytes. Patients at a high risk will be prescribed medicines to reduce the risk of complications. swelling in a leg, or breathlessness and chest - you must tell your doctor straight away if you have any of these symptoms. Some anti-cancer medicines can damage women's ovaries and men's sperm. This may lead to infertility in men and women and/or of menopause in women. Some anti-cancer medicines may damage the development of a baby in the womb. It is important not to become pregnant or father a child while you are having treatment and for 6 months afterwards. It is important to use effective contraception during and for 6 months after treatment. You can talk to your doctor or nurse about this. Very rarely complications of treatment with a cancer medicines can be life-threatening or eresult in death. The risks are different for ever individual. You can talk to your doctor or nurse individual. You can talk to your doctor or nurse individual. 			Cancer can increase your risk of developing a blood clot (thrombosis), and having treatment with anti-cancer medicines may increase this risk			
Between 1 and 10 in every 100 (1-10%) people have one or more of these effects: Diarrhoea, constipation, drowsiness, sore mouth and ulcers, loss of appetite, muscle and back pain, sweating, runny nose, and itchy skin. There may be a risk of tumour lysis syndrome in the presence of a high tumour burden. Symptoms are caused by the breakdown of cancer cells when treatment is started, causing changes in blood electrolytes. Patients at a high risk will be prescribed medicines to reduce the risk of complications. Some anti-cancer medicines may damage the development of a baby in the womb. It is important not to become pregnant or father a child while you are having treatment and for 6 months afterwards. It is important to use effective contraception during and for 6 months after treatment. You can talk to your doctor or nurse about this. Very rarely complications of treatment with a cancer medicines can be life-threatening or eresult in death. The risks are different for every individual. You can talk to your doctor or nurse individual. You can talk to your doctor or nurse individual. You can talk to your doctor or nurse individual. You can talk to your doctor or nurse individual. You can talk to your doctor or nurse individual. You can talk to your doctor or nurse individual. You can talk to your doctor or nurse individual. You can talk to your doctor or nurse individual. You can talk to your doctor or nurse individual. You can talk to your doctor or nurse individual. You can talk to your doctor or nurse individual. You can talk to your doctor or nurse individual. You can talk to your doctor or nurse individual. You can talk to your doctor or nurse individual. You can talk to your doctor or nurse individual. You can talk to your doctor or nurse individual you are having treatment and for 6 months after treatment and for 6 months after treatment. You can talk to your doctor or nurse in your doct			swelling in a leg, or breathlessness and chest pa - you must tell your doctor straight away if you			
and ulcers, loss of appetite, muscle and back pain, sweating, runny nose, and itchy skin. There may be a risk of tumour lysis syndrome in the presence of a high tumour burden. Symptoms are caused by the breakdown of cancer cells when treatment is started, causing changes in blood electrolytes. Patients at a high risk will be prescribed medicines to reduce the risk of complications. Some anti-cancer medicines may damage the development of a baby in the womb. It is important not to become pregnant or father a child while you are having treatment and for 6 months afterwards. It is important to use effective contraception during and for 6 months after treatment. You can talk to your doctor or nurse about this. Very rarely complications of treatment with a cancer medicines can be life-threatening or end individual. You can talk to your doctor or nurse individual. You can talk to your doctor or nurse in the development of a baby in the womb. It is important not to become pregnant or father a child while you are having treatment and for 6 months afterwards. It is important to use effective contraception during and for 6 months after treatment. You can talk to your doctor or nurse about this.	Between 1 and 10 in every 100 (1-10%) people have one or more of these effects:		women's ovaries and men's sperm. This may lead to infertility in men and women and/or ea			
changes in blood electrolytes. Patients at a high risk will be prescribed medicines to reduce the risk of complications. I described medicines to reduce the risk of complications. Very rarely complications of treatment with a cancer medicines can be life-threatening or expected to the result in death. The risks are different for every individual. You can talk to your doctor or nurse.	pain, sweating, runny nose, and itchy skin. There may be a risk of tumour lysis syndrome in the presence of a high tumour burden. Symptoms are caused by the breakdown of cancer cells when treatment is started, causing changes in blood electrolytes. Patients at a high risk will be prescribed medicines to reduce the		Some anti-cancer medicines may damage the development of a baby in the womb. It is important not to become pregnant or father a child while you are having treatment and for			
cancer medicines can be life-threatening or e result in death. The risks are different for ever individual. You can talk to your doctor or nurs						
about what this means for you.			Very rarely complications of treatment with anti- cancer medicines can be life-threatening or even result in death. The risks are different for every individual. You can talk to your doctor or nurse about what this means for you.			

STATEMENT OF HEALTH PROFESSIONAL (continued)

(continued)	: Patient Identiner/label
ANY OTHER RISKS:	
 I have discussed what the treatment is likely to invoof the treatment, blood and any additional tests, for I have discussed the intended benefits and risks of treatment) and any particular concerns of this patients. 	any available alternative treatments (including no
THE FOLLOWING LEAFLET HAS BEEN PROVIDE	D:
Information leaflet for gemcitabine chemotherapy.24 hour chemotherapy service contact details	Other, please state:
Signed:	Date:
Name (PRINT):	
Job title:	
STATEMENT OF INTERPRETER (v	
have interpreted the information above to the patient to s/he can understand.	
Signed:	Date:
Name (PRINT):	

Job title: _

STATEMENT OF PATIENT

Patient identifier/label

Please read this form carefully. If your treatment has been planned in advance, you should already have your own copy of the form which describes the benefits and risks of the proposed treatment. If not, you will be offered a copy now. If you have any further questions, do ask - we are here to help you. You have the right to change your mind at any time, including after you have signed this form.

I agree to the procedure and course of treatment described on this form.

I understand that you cannot give me a guarantee that a particular person will perform the procedure. The person will, however, have appropriate training and experience.

I understand that any procedure in addition to those described on this form will only be carried out if it is necessary to save my life or to prevent serious harm to my health.

I have been told about additional procedures which may become necessary during my treatment. I have listed below any procedures which I do not wish to be carried out without further discussion:

Patient's signature:	Date:
Name (PRINT):	
A witness should sign below if the patient is unable to sig may also like a parent to sign here (see notes).	n but has indicated his or her consent. Young people/children
Parent's/Witness' signature:	_ Date:
Name (PRINT):	

COPY ACCEPTED BY PATIENT: YES / NO

(please circle)

Job title: _

CONFIRMATION OF CONSENT

(health professional to complete when the patient attends for treatment, if the patient has signed the form in advance)

On behalf of the team treating the patient, I have confirmed with the patient that s/he has no further questions and wishes the procedure to go ahead.

Signed: Date: Name (PRINT):_

IMPORTANT NOTES: (tick if applicable)

	See a	lso ac	lvance (decision	to re	tuse	trea	tmen
--	-------	--------	----------	----------	-------	------	------	------

Patient has withdrawn consent (ask patient to sign /date here)

Signed: _

FURTHER INFORMATION FOR PATIENTS

CONTACT DETAILS (if patient wishes to discuss options later):

Contact your hospital team if you have any questions about cancer and treatment.

Cancer Research UK can also help answer your questions about cancer and treatment. If you want to talk in confidence, call our information nurses on freephone 0808 800 4040, Monday to Friday, 9am to 5pm. Alternatively visit www.cruk.org for more information.

These forms have been produced by Guy's and St. Thomas' NHS Foundation Trust as part of a national project to support clinicians in ensuring all patients are fully informed when consenting to SACT. The project is supported by Cancer Research UK. This does not mean you are taking part in a



GUIDANCE FOR HEALTH PROFESSIONALS

(to be read in conjunction with the hospital's consent policy)

WHAT A CONSENT FORM IS FOR

This form documents the patient's agreement to go ahead with the investigation or treatment you have proposed. It is not a legal waiver – if patients, for example, do not receive enough information on which to base their decision, then the consent may not be valid, even though the form has been signed. Patients are also entitled to change their mind after signing the form, if they retain capacity to do so. The form should act as an aide-memoir to health professionals and patients, by providing a checklist of the kind of information patients should be offered, and by enabling the patient to have a written record of the main points discussed. In no way, however, should the written information provided for the patient be regarded as a substitute for face-to-face discussions with the patient.

THE LAW ON CONSENT

See the Department of Health's Reference guide to consent for examination or treatment 2nd Edition for a comprehensive summary of the law on consent (also available at www.doh.gov.uk).

WHO CAN GIVE CONSENT

Everyone aged 16 or over is presumed to have the capacity to give consent for themselves, unless the opposite is demonstrated. If a child under the age of 16 has "sufficient understanding and intelligence to enable him or her to understand fully what is proposed", then he or she will have capacity to give consent for himself or herself. Young people aged 16 and 17, and younger children with capacity, may therefore sign this form for themselves, but may like a parent to countersign as well. If the child is not able to give consent for himself or herself, someone with parental responsibility may do so on their behalf and a separate form is available for this purpose. Even where a child is able to give consent for himself or herself, you should always involve those with parental responsibility in the child's care, unless the child specifically asks you not to do so. If a patient has the capacity to give consent but is physically unable to sign a form, you should complete this form as usual, and ask an independent witness to confirm that the patient has given consent orally or non-verbally.

WHEN NOT TO USE THIS FORM

If the patient is 18 or over and lacks the capacity to give consent, you should use an alternative form (form for adults who lack the capacity to consent to investigation or treatment) instead of this form. A patient lacks capacity if they have an impairment of the mind or brain or

Patient identifier/label

disturbance affecting the way their mind or brain works and they cannot:

- understand information about the decision to be made
- retain that information in their mind
- use or weigh this information as a part of their decision making process, or
- communicate their decision (by talking, using sign language or any other means)

You should always take all reasonable steps (for example involving more specialist colleagues) to support a patient in making their own decision, before concluding that they are unable to do so.

Relatives cannot be asked to sign a form on behalf of an adult who lacks capacity to consent for themselves, unless they have been given the authority to do so under a Lasting Power of Attorney or as a court deputy.

INFORMATION

Information about what the treatment will involve, its benefits and risks (including side-effects and complications) and the alternatives to the particular procedure proposed, is crucial for patients when making up their minds. The courts have stated that patients should be told about 'significant risks which would affect the judgement of a reasonable patient'. 'Significant' has not been legally defined, but the GMC requires doctors to tell patients about 'significant, unavoidable or frequently occurring' risks. In addition if patients make clear they have particular concerns about certain kinds of risk, you should make sure they are informed about these risks, even if they are very small or rare. You should always answer questions honestly. Sometimes, patients may make it clear that they do not want to have any information about the options, but want you to decide on their behalf. In such circumstances, you should do your best to ensure that the patient receives at least very basic information about what is proposed. Where information is refused, you should document this on the consent form or in the patient's notes.

REFERENCES

- Summary of Product Characteristics (SPCs) for individual drugs: https://www.medicines.org.uk/emc/
- Cancer Research UK: http://www.cancerresearchuk.org/ about-cancer/cancers-in-general/treatment/cancer-drugs/
- Macmillan Cancer Support, Cancer Information: http://www. macmillan.org.uk/Cancerinformation/Cancertreatment/ Treatmenttypes/Chemotherapy/Chemotherapy.aspx
- 4. Guy's and St. Thomas' NHS Foundation Trust, Chemotherapy consent forms.