

Lung cancer




Number of days to start chemotherapy, with the percentage of patients treated (2012–2017): **stages 1–3 or local and regional**



Australia

New South Wales



Canada

Ontario



Saskatchewan-Manitoba



Alberta



Atlantic Canada



British Columbia



0% 10% 20% 30% 40% 50% 60%



Norway

Norway



0 10% 20% 30% 40% 50% 60%



UK

Wales



England



Northern Ireland



Scotland



0 10% 20% 30% 40% 50% 60%

Number of days to start treatment (percentile)

Median



Percentage treated

TNM and SEER stages do not always align: UK and Canada used TNM staging, Norway and Australia used SEER.

Atlantic Canada includes Prince Edward Island and Nova Scotia.

Some jurisdictions had a large % of missing stage data which could introduce bias: Norway (12%), New South Wales (14%).

Lung cancer




Number of days to start chemotherapy, with the percentage of patients treated (2012–2017): **stage 4 or distant**



Australia

New South Wales



Canada

Ontario



British Columbia



Alberta



Saskatchewan-Manitoba



Atlantic Canada



Norway

Norway



UK

England



Wales




Scotland



Northern Ireland



Number of days to start treatment (percentile)



TNM and SEER stages do not always align: UK and Canada used TNM staging, Norway and Australia used SEER.

Atlantic Canada includes Prince Edward Island and Nova Scotia.

Some jurisdictions had a large % of missing stage data which could introduce bias: Norway (12%), New South Wales (14%).

Lung cancer




Number of days to start radiotherapy, with the percentage of patients treated (2012–2017): **stages 1–3 or local and regional**



Australia

New South Wales



Canada

Ontario



Atlantic Canada



British Columbia



Alberta



Saskatchewan-Manitoba



Norway

Norway



UK

Northern Ireland



Scotland




Wales



England



Number of days to start treatment (percentile)



TNM and SEER stages do not always align: UK and Canada used TNM staging, Norway and Australia used SEER.

Atlantic Canada includes Newfoundland & Labrador, New Brunswick, Nova Scotia and Prince Edward Island.

Some jurisdictions had a large % of missing stage data which could introduce bias: Norway (12%), New South Wales (14%), Northern Ireland (10%).

Lung cancer




Number of days to start radiotherapy, with the percentage of patients treated (2012–2017): **stage 4 or distant**



Australia

New South Wales



Canada

British Columbia



Alberta



Saskatchewan-Manitoba



Atlantic Canada



Ontario



Norway

Norway



UK

Northern Ireland



Wales




England



Scotland



Number of days to start treatment (percentile)



TNM and SEER stages do not always align: UK and Canada used TNM staging, Norway and Australia used SEER.

Atlantic Canada includes Newfoundland & Labrador, New Brunswick, Nova Scotia and Prince Edward Island.

Some jurisdictions had a large % of missing stage data which could introduce bias: Norway (12%), New South Wales (14%), Northern Ireland (10%).