


Lung cancer


Average chemotherapy use by age groups (2012–2017): **all stages**



Coverage for Australia and Canada are not complete. Australia includes New South Wales and Victoria.
Canada includes Alberta, British Columbia, Ontario, Manitoba, Nova Scotia, Prince Edward Island, Saskatchewan.
Northern Ireland and Saskatchewan-Manitoba (85–99 only) were suppressed and not included in the country averages.
Some jurisdictions had a large % of missing stage data which could introduce bias: Norway (12%); New South Wales (14%).

Lung cancer

Average chemotherapy use by age groups (2012–2017): **all stages**




Coverage for Australia and Canada are not complete. Australia includes New South Wales and Victoria.
Canada includes Alberta, British Columbia, Ontario, Manitoba, Nova Scotia, Prince Edward Island, Saskatchewan.
Northern Ireland and Saskatchewan-Manitoba (85–99 only) were suppressed and not included in the country averages.
Some jurisdictions had a large % of missing stage data which could introduce bias: Norway (12%); New South Wales (14%).

Lung cancer

Average chemotherapy use by age groups (2012–2017): **all stages**



UK




♦ Small numbers (between 1–9) were suppressed in Northern Ireland (85–99).

Lung cancer

Average chemotherapy use by age groups (2012–2017): **all stages**



Canada




Atlantic Canada includes Nova Scotia and Prince Edward Island.

♦ Small numbers (less than 30) were suppressed in Saskatchewan-Manitoba (85–99).

Very small numbers may have been suppressed in Atlantic Canada (85–99).

Lung cancer

Average chemotherapy use by age groups (2012–2017): **stages 1–3 or local and regional**



TNM and SEER stages do not always align: UK and Canada used TNM staging, Norway and Australia used SEER.


Coverage for Australia and Canada are not complete. Australia includes New South Wales only. Canada includes Alberta, British Columbia, Ontario, Manitoba, Nova Scotia, Prince Edward Island, Saskatchewan.

Scotland, Wales, and Alberta (85–99 only) were suppressed and not included in country averages.

Some jurisdictions had a large % of missing stage data which could introduce bias: Norway (12%); New South Wales (14%).

Lung cancer

Average chemotherapy use by age groups (2012–2017): **stages 1–3 or local and regional**



TNM and SEER stages do not always align: UK and Canada used TNM staging, Norway and Australia used SEER.

Coverage for Australia and Canada are not complete. Australia includes New South Wales only. Canada includes Alberta, British Columbia, Ontario, Manitoba, Nova Scotia, Prince Edward Island, Saskatchewan.

Scotland, Wales, and Alberta (85–99 only) were suppressed and not included in the country averages.


Some jurisdictions had a large % of missing stage data which could introduce bias: Norway (12%); New South Wales (14%).

Lung cancer

Average chemotherapy use by age groups (2012–2017): **stages 1–3**



UK




♦ Small numbers (between 1–9) were suppressed in Scotland and Wales (85–99).
Very small numbers may have been suppressed in Northern Ireland (85–99).

Lung cancer

Average chemotherapy use by age groups (2012–2017): **stages 1–3**



Canada




Atlantic Canada includes Nova Scotia and Prince Edward Island.

♦ Small numbers (between 1–9) were suppressed in Alberta (85–99).

Very small numbers may have been suppressed in Atlantic Canada and Saskatchewan-Manitoba (85–99).

Lung cancer

Average chemotherapy use by age groups (2012–2017): **stage 4 or distant**



TNM and SEER stages do not always align: UK and Canada used TNM staging, Norway and Australia used SEER.


Coverage for Australia and Canada are not complete. Australia includes New South Wales only. Canada includes Alberta, British Columbia, Ontario, Manitoba, Nova Scotia, Prince Edward Island, Saskatchewan.

Northern Ireland and Saskatchewan-Manitoba (85-99 only) were suppressed and not included in the country averages.

Some jurisdictions had a large % of missing stage data which could introduce bias: Norway (12%); New South Wales (14%).

Lung cancer

Average chemotherapy use by age groups (2012–2017): **stage 4 or distant**



TNM and SEER stages do not always align: UK and Canada used TNM staging, Norway and Australia used SEER.

Coverage for Australia and Canada are not complete. Australia includes New South Wales only. Canada includes Alberta, British Columbia, Ontario, Manitoba, Nova Scotia, Prince Edward Island, Saskatchewan.

Northern Ireland and Saskatchewan-Manitoba (85-99 only) were suppressed and not included in the country averages.


Some jurisdictions had a large % of missing stage data which could introduce bias: Norway (12%); New South Wales (14%).

Lung cancer

Average chemotherapy use by age groups (2012–2017): **stage 4**



UK




◆ Small numbers (between 1–9) were suppressed in Northern Ireland (85–99).

Lung cancer

Average chemotherapy use by age groups (2012–2017): **stage 4**



Canada



Atlantic Canada includes Nova Scotia and Prince Edward Island.

♦ Small numbers (less than 30) were suppressed in Saskatchewan-Manitoba (85–99).

Very small numbers may have been suppressed in Atlantic Canada (85–99).