

Ovarian cancer




Number of days to start chemotherapy, with the percentage of patients treated (2012–2017): **stages 1–2 or local and regional**



Australia

New South Wales



Canada

Manitoba



Nova Scotia



Ontario



British Columbia



Alberta



0 20% 40% 60% 80% 100%



Norway

Norway



0 20% 40% 60% 80% 100%



UK

Scotland



England



Northern Ireland



Wales



0 20% 40% 60% 80% 100%

Number of days to start treatment (percentile)

Median



Percentage treated

TNM and SEER stages do not always align: UK and Canada used TNM staging, Norway and Australia used SEER.

Some jurisdictions had a large % of missing stage data which could introduce bias: England (17%), Northern Ireland (12%), Scotland (13%), Wales (28%), British Columbia (34%), Ontario (33%).

Ovarian cancer




Number of days to start chemotherapy, with the percentage of patients treated (2012–2017): **stages 3–4 or distant**



Australia

New South Wales



Canada

Ontario



British Columbia



Alberta



Manitoba




Nova Scotia



Norway

Norway



UK

Scotland



Wales



England



Northern Ireland



Number of days to start treatment (percentile)

Median



Percentage treated

TNM and SEER stages do not always align: UK and Canada used TNM staging, Norway and Australia used SEER.

Some jurisdictions had a large % of missing stage data which could introduce bias: England (17%), Northern Ireland (12%), Scotland (13%), Wales (28%), British Columbia (34%), Ontario (33%).