# Science Review 2017

# Sourcebook

An annual compilation of
research data uses enabled by

GBIF

# About the *Science Review* Sourcebook 2017

Each year, the GBIF *Science Review* highlights research uses of primary biodiversity data accessed through the GBIF network global infrastructure.

Citations of GBIF-mediated data continue to grow and now average more than one substantive example each day. In view of the corresponding growth in the number of pages of the *Science Review*, the narrative highlights remain separated from the full annual bibliographic table of citations, which includes more than 400 individual articles. This sourcebook is an electronic companion volume to the printed GBIF *Science Review* 2017 (also available electronically) which includes summaries of selected examples of research using GBIF as a data source, as well as analysis of trends and geographic location of research teams.

Articles summarized in the *Science Review* are **marked in bold** in the following tables, and the symbol 'a' indicates open-access articles that are available to all readers with or without a journal subscription.

## CONTENTS

## biodiversity and human health

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| **Alimi TO et al.** | **A multi-criteria decision analysis approach to assessing malaria risk in northern South America** | **BMC Public Health** | **https://doi.org/10.1186/s12889-016-2902-7** | **United States, Colombia** |
| Baruah D & Neog B | Botanical , Phytochemical and Pharmacological Review of Flacourtia Jangomas (Lour.) Raeusch | International Journal of Current Medical and Pharmaceutical Research | http://bit.ly/2ugUnRE | India |
| Cao B et al. | Modeling habitat distribution of Cornus officinalis with Maxent modeling and fuzzy logics in China | Journal of Plant Ecology | https://doi.org/10.1093/jpe/rtw009 | China |
| Carlson CJ et al. | An Ecological Assessment of the Pandemic Threat of Zika Virus | PLoS Neglected Tropical Diseases 10(8) | https://doi.org/10.1371/journal.pntd.0004968 | United States, South Africa |
| **Escobar LE et al.** | **Forecasting Chikungunya spread in the Americas via data-driven empirical approaches** | **Parasites & Vectors 9:112** | **https://doi.org/10.1186/s13071-016-1403-y** | **United States, China** |
| **Estrada-Peña A & de la Fuente J** | **Species interactions in occurrence data for a community of tick-transmitted pathogens** | **Scientific Data 3:160056** | **https://doi.org/10.1038/sdata.2016.56** | **Spain, United States** |
| Giovannini P et al. | Medicinal plants used in the traditional management of diabetes and its sequelae in Central America: A review | Journal of Ethnopharmacology 184: 58-71 | https://doi.org/10.1016/j.jep.2016.02.034 | United Kingdom |
| Hutter SE et al. | Rabies in Costa Rica: Documentation of the Surveillance Program and the Endemic Situation from 1985 to 2014 | Vector-Borne and Zoonotic Diseases | https://doi.org/10.1089/vbz.2015.1906. | Austria, Costa Rica |
| Ke Z et al. | Drug discovery of neurodegenerative disease through network pharmacology approach in herbs | Biomedicine & Pharmacotherapy 78: 272-279 | https://doi.org/10.1016/j.biopha.2016.01.021 | China |
| Martin GA et al. | Climatic suitability influences species specific abundance patterns of Australian flying foxes and risk of Hendra virus spillover | One Health 2: 115-121 | https://doi.org/10.1016/j.onehlt.2016.07.004 | Australia, Mexico, United States |
| **Moyes CL et al.** | **Predicting the geographical distributions of the macaque hosts and mosquito vectors of Plasmodium knowlesi malaria in forested and non-forested areas** | **Parasites & Vectors** | **https://doi.org/10.1186/s13071-016-1527-0** | **United Kingdom, Malaysia, Canada, Thailand, Singapore, Japan, Kenya, Australia, United States, Indonesia** |
| **Pech-May A et al.** | **Assessing the importance of four sandfly species (Diptera: Psychodidae) as vectors of Leishmania mexicana in Campeche, Mexico** | **Medical and Veterinary Entomology** | **https://doi.org/10.1111/mve.12169** | **Argentina, Mexico** |
| Piedrahita-Cortés J & Soler-Tovar D | Geographical distribution of the red howler monkey (Alouatta seniculus) and yellow fever in Colombia | Biomedica 36(0): 116-21 | https://doi.org/10.7705/biomedica.v36i0.2929 | Colombia |
| **Pigott DM et al.** | **Updates to the zoonotic niche map of Ebola virus disease in Africa** | **eLife 5:e16412** | **https://doi.org/10.7554/eLife.16412** | **United Kingdom, United States, Australia** |
| **Redding DW et al.** | **Environmental-mechanistic modelling of the impact of global change on human zoonotic disease emergence: a case study of Lassa fever** | **Methods in Ecology and Evolution, 7: 646–655** | **https://doi.org/10.1111/2041-210X.12549** | **United Kingdom, United States** |
| **Rivera D et al.** | **Is there nothing new under the sun? The influence of herbals and pharmacopoeias on ethnobotanical traditions in Albacete (Spain)** | **Journal of Ethnopharmacology** | **https://doi.org/10.1016/j.jep.2016.11.040** | **Spain, Italy** |
| **Samy AM et al.** | **Climate Change Influences on the Global Potential Distribution of the Mosquito Culex quinquefasciatus, Vector of West Nile Virus and Lymphatic Filariasis** | **PLoS ONE 11(10)** | **https://doi.org/10.1371/journal.pone.0163863** | **Sudan, Egypt, Brazil, United States** |
| **Stephens CR et al.** | **Can You Judge a Disease Host by the Company It Keeps? Predicting Disease Hosts and Their Relative Importance: A Case Study for Leishmaniasis** | **PLoS Neglected Tropical Diseases 10(10): e0005004** | **https://doi.org/10.1371/journal.pntd.0005004** | **Mexico** |
| Wansi JD et al. | Phytochemistry and pharmacology of the genus Drypetes: A review | Journal of Ethnopharmacology | https://doi.org/10.1016/j.jep.2016.06.060 | Cameroon, United Kingdom, Germany |
| Yi Y-J et al. | Maxent modeling for predicting the potential distribution of endangered medicinal plant (H. riparia Lour) in Yunnan, China | Ecological Engineering 92: 260-269 | https://doi.org/10.1016/j.ecoleng.2016.04.010 | China |
| Zhang L et al. | Predicting suitable cultivation regions of medicinal plants with Maxent modeling and fuzzy logics: a case study of Scutellaria baicalensis in China | Environmental Earth Sciences 75:361 | https://doi.org/10.1007/s12665-015-5133-9 | China |

# data management & data papers

| | AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|---|
| a | Alexander NS et al. | The European Distribution of Sus Scrofa. Model Outputs from the Project Described within the Poster – Where are All the Boars? An Attempt to Gain a Continental Perspective | Journal of open health data 4 | https://doi.org/10.5334/ohd.24 | United Kingdom |
| | Asase A & Peterson AT | Completeness of Digital Accessible Knowledge of the Plants of Ghana | Biodiversity Informatics | https://doi.org/10.17161/bi.v11i0.5860 | Ghana, United States |
| | **Barve V & Otegui J** | **bdvis: visualizing biodiversity data in R** | **Bioinformatics** | **https://doi.org/10.1093/bioinformatics/btw333** | **United States** |
| | Becker D et al. | A Plugin to Interface Openmodeller from QGIS for Species' Potential Distribution Modelling | ISPRS Annals of teh Photogrammetry, Remote Sensing and Spatial Information Sciences 3(7) | https://doi.org/10.5194/isprs-annals-III-7-251-2016 | Germany |
| a | Dikow T & Agosti D | Utilizing online resources for taxonomy: a cybercatalog of Afrotropical apiocerid flies (Insecta: Diptera: Apioceridae) | Biodiversity Data Journal 3: e5707 | https://doi.org/10.3897/BDJ.3.e5707 | United States, Switzerland |
| a | Droege G et al. | The Global Genome Biodiversity Network (GGBN) Data Standard specification | Database | https://doi.org/10.1093/database/baw125 | Germany, United Staes, Denmark, United Kingdom, Australia, Italy |
| | **Franklin J et al.** | **Big data for forecasting the impacts of global change on plant communities** | **Global Ecology and Biogeography** | **https://doi.org/10.1111/geb.12501** | **United States** |
| a | Gattolliat J-L et al. | Toward a comprehensive COI DNA barcode library for Swiss Stoneflies (Insecta: Plecoptera) with special emphasis on the genus Leuctra | Zoosymposia 11: 135–155 | https://doi.org/10.11646/zoosymposia.11.1.15 | Switzerland |
| a | **Gilles D et al.** | **RAINBIO: a mega-database of tropical African vascular plants distributions** | **PhytoKeys 74: 1-18** | **https://doi.org/10.3897/phytokeys.74.9723** | **France, Portugal, Denmark, Netherlands, Belgium, United Kingdom, United States, Germany** |
| | **Ivanova NV & Shashkov MP** | **Biodiversity Databases in Russia: Towards a National Portal** | **Arctic Science** | **https://doi.org/10.1139/AS-2016-0050** | **Russian Federation** |
| | Kirchner AC & Aden C | HotSpotSDM - ein WPS-Prozess für die Modellierung ökologischer Artenfunde | AGIT - Journal für Angewandte Geoinformatik | https://doi.org/10.14627/537622077 | Germany |
| | **Michonneau F et al.** | **rotl: an R package to interact with the Open Tree of Life data** | **Methods in Ecology and Evolution** | **https://doi.org/10.1111/2041-210X.12593** | **United States** |
| | **Rambold G et al.** | **Geographic heat maps of lichen traits derived by combining LIAS light description and GBIF occurrence data, provided on a new platform** | **Biodiversity & Conservation** | **https://doi.org/10.1007/s10531-016-1199-2** | **Germany, United States** |
| a | Rekadwad BN & Khobragade CN | Digital data for quick response (QR) codes of alkalophilic Bacillus pumilus to identify and to compare bacilli isolated from Lonar Crator Lake, India | Data in Brief 7: 1306-1313 | https://doi.org/10.1016/j.dib.2016.03.103 | India |
| a | Thessen AE et al. | Emerging semantics to link phenotype and environment | PeerJ 3:e1470 | https://doi.org/10.7717/peerj.1470 | United States, Germany, Argentina, Netherlands |
| a | **Ytow N** | **Taxonaut: an application software for comparative display of multiple taxonomies with a use case of GBIF Species API** | **Biodiversity Data Journal 4: e9787** | **https://doi.org/10.3897/BDJ.4.e9787** | **Japan** |

## food & farming

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| **Bosch J et al.** | **A Cartographic Tool for Managing African Swine Fever in Eurasia: Mapping Wild Boar Distribution Based on the Quality of Available Habitats.** | **Transboundary and Emerging Diseases** | **https://doi.org/10.1111/tbed.12559** | **Spain** |
| Bunn C et al. | Multiclass Classification of Agro-Ecological Zones for Arabica Coffee: An Improved Understanding of the Impacts of Climate Change | PLoS ONE 10(10): e0140490 | https://doi.org/10.1371/journal.pone.0140490 | Colombia, Nicaragua, United States |
| Bush MB et al. | A 6900-year history of landscape modification by humans in lowland Amazonia | Quaternary Science Reviews 141: 52-64 | https://doi.org/10.1016/j.quascirev.2016.03.022 | United States, Mexico, Netherlands |
| Castañeda-Álvarez NP et al. | Global conservation priorities for crop wild relatives | Nature Plants 2 | https://doi.org/10.1038/nplants.2016.22 | Colombia, United Kingdom, Netherlands, Germany |
| da Silva RS et al. | Potential risk levels of invasive Neoleucinodes elegantalis (small tomato borer) in areas optimal for open field Solanum lycopersicum (tomato) cultivation in the present and under predicted climate change | Pest Management Science | https://doi.org/10.1002/ps.4344 | Brazil, Australia |
| Delgado-Baquerizo M et al. | Biogeographic bases for a shift in crop C : N : P stoichiometries during domestication | Ecology Letters | https://doi.org/10.1111/ele.12593 | Australia, United States, Spain |
| **Dunne JA et al.** | **The roles and impacts of human hunter-gatherers in North Pacific marine food webs** | **Scientific Reports 6: 21179** | **https://doi.org/10.1038/srep21179** | **United States, Canada** |
| **Dupin J et al.** | **Bayesian estimation of the global biogeographical history of the Solanaceae** | **Journal of Biogeography** | **https://doi.org/10.1111/jbi.12898** | **United States, Australia, United Kingdom** |
| Galluzzi G et al. | An Integrated Hypothesis on the Domestication of Bactris gasipaes | PLoS ONE 10(12): e0144644 | https://doi.org/10.1371/journal.pone.0144644 | Colombia, Costa Rica |
| **Halouani G et al.** | **Modelling food web structure using an end-to-end approach in the coastal ecosystem of the Gulf of Gabes (Tunisia)** | **Ecological Modelling 45:57** | **https://doi.org/10.1016/j.ecolmodel.2016.08.008** | **Tunisia, France, South Africa, Peru** |
| Kaur G & Sangha KS | Diversity of arthropod fauna associated with chilli (Capsicum annuum L.) in Punjab | Journal of Entomology and Zoology Studies 4(5): 390-396 | http://bit.ly/2uh30M7 | India |
| López-Martínez V et al. | Current and Potential Distribution of the Cactus Weevil, Cactophagus spinolae (Gyllenhal) (Coleoptera: Curculionidae), in Mexico | The Coleopterists Bulletin 70 (2): 327-334 | https://doi.org/10.1649/0010-065X-70.2.327 | Mexico |
| **Osawa T et al.** | **Trade-off relationship between modern agriculture and biodiversity: Heavy consolidation work has a long-term negative impact on plant species diversity** | **Land Use Policy 54: 78-84** | **https://doi.org/10.1016/j.landusepol.2016.02.001** | **Japan** |
| Plath M et al. | A novel bioenergy feedstock in Latin America? Cultivation potential of Acrocomia aculeata under current and future climate conditions | Biomass and Bioenergy 9: 186-195 | https://doi.org/10.1016/j.biombioe.2016.04.009 | Germany, United States, Kenya, Netherlands |
| **Samy AM & Peterson AT** | **Climate Change Influences on the Global Potential Distribution of Bluetongue Virus** | **PLoS ONE 11(3): e0150489** | **https://doi.org/10.1371/journal.pone.0150489** | **United States, Egypt** |
| **Sanyal A & Decocq G** | **Adaptive evolution of seed oil content in angiosperms: accounting for the global patterns of seed oils** | **BMC Evolutionary Biology 16: 187** | **https://doi.org/10.1186/s12862-016-0752-7** | **France, Sweden** |
| Shaik RS et al. | The biology and management of prickly paddy melon (Cucumis myriocarpus L.), an important summer annual weed in Australia | Crop Protection 92: 29-40 | https://doi.org/10.1016/j.cropro.2016.10.005 | Australia |
| Syfert MM et al. | Crop wild relatives of the brinjal eggplant (Solanum melongena): Poorly represented in genebanks and many species at risk of extinction | American Journal of Botany 103 (10) | https://doi.org/10.3732/ajb.1500539 | United Kingdom, Netherlands, Colombia, Spain, France |
| Thomas E et al. | An assessment of the conservation status of Mesoamerican crop species and their wild relatives in light of climate change | Chapter XX in Enhancing Crop Genepool Use: Capturing Wild Relative and Landrace Diversity for Crop Improvement | http://bit.ly/2uhvjdo | Colombia, Costa Rica |

## impacts of climate change

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| Aljaryian R & Kumar L | Changing global risk of invading greenbug Schizaphis graminum under climate change | Crop Protection 88: 137-148 | https://doi.org/10.1016/j.cropro.2016.06.008 | Australia |
| **Biber-Freudenberger L et al.** | **Future Risks of Pest Species under Changing Climatic Conditions** | **PLoS ONE 11(4): e0153237** | **https://doi.org/10.1371/journal.pone.0153237** | **Germany, Kenya** |
| de Oliveira IS et al. | Assessing future habitat availability for coastal lowland anurans in the Brazilian Atlantic rainforest | Studies on Neotropical Fauna and Environment 51(1): 45-55 | https://doi.org/10.1080/01650521.2016.1160610 | Brazil, Germany, Portugal, Iran |
| Dotchamou FT et al. | Density and spatial distribution of Parkia biglobosa pattern in Benin under climate change | Journal of Agriculture and Envornment for international Development 110 (1): 173-194 | https://doi.org/10.12895/jaeid.20161.447 | Benin |
| **Duffy GA & Chown SL** | **Urban warming favours C4 plants in temperate European cities** | **Journal of Ecology** | **https://doi.org/10.1111/1365-2745.12652** | **Australia** |
| Feldman RE et al. | Climate driven range divergence among host species affects range-wide patterns of parasitism | Global Ecology and Conservation 9: 1-10 | https://doi.org/10.1016/j.gecco.2016.10.001 | Mexico, United States, Canada |
| **Frishkoff LO et al.** | **Climate change and habitat conversion favour the same species** | **Ecology Letters** | **https://doi.org/10.1111/ele.12645** | **United States, Canada, United Kingdom, Costa Rica, Sweden** |
| García-Martínez YG et al. | Traditional Agroecosystems and Global Change Implications in Mexico | Bulgarian Journal of Agricultural Science 22(4): 548-565 | http://bit.ly/2uhmT5w | Mexico |
| González-Santos R et al. | Prediction of the effects of climate change on Sechium edule (Jacq.) Swartz varietal groups in Mexico | Genetic Resources and Crop Evolution | https://doi.org/10.1007/s10722-016-0401-4 | Mexico |
| Graham CH et al. | Winter conditions influence biological responses of migrating hummingbirds | Ecosphere 7(10) | https://doi.org/10.1002/ecs2.1470 | United States |
| Hannah L et al. | Climate change influences on pollinator, forest, and farm interactions across a climate gradient | Climatic Change | https://doi.org/10.1007/s10584-016-1868-x | United States, Costa Rica |
| Hargrave MS et al. | The effects of warming on the ecophysiology of two co-existing kelp species with contrasting distributions | Oecologia | https://doi.org/10.1007/s00442-016-3776-1 | United Kingdom |
| **Harsch MA & HilleRisLambers J** | **Climate Warming and Seasonal Precipitation Change Interact to Limit Species Distribution Shifts across Western North America** | **PLoS ONE 11(7): e0159184** | **https://doi.org/10.1371/journal.pone.0159184** | **United States** |
| **Hattab T et al.** | **Forecasting fine-scale changes in the food-web structure of coastal marine communities under climate change** | **Ecography** | **https://doi.org/10.1111/ecog.01937** | **France, Tunisia, Canada** |
| Hof AR et al. | Vulnerability of subarctic and arctic breeding birds | Ecological Applications | https://doi.org/10.1002/eap.1434 | Sweden, United States |
| **Ikeda DH et al.** | **Genetically informed ecological niche models improve climate change predictions** | **Global Change Biology** | **https://doi.org/10.1111/gcb.13470** | **United States** |
| Iloh AC & Ogundipe OT | Using ecological niche models to plan conservation in a changing environment: A case for the plant Chasmanthera dependens Hochst (Menispermaceae) in West Africa | Journal of Ecology and The Natural Environment 8(1): 1-8 | https://doi.org/10.5897/JENE2015.0525 | Nigeria |
| Jiang H et al. | Predicting the Potential Distribution of Polygala tenuifolia Willd. under Climate Change in China | PLoS ONE 11(9): e0163718 | https://doi.org/10.1371/journal.pone.0163718 | China |
| Moutouama JK et al. | Potential climate change favored expansion of a range limited species, Haematostaphis barteri Hook f. | Journal of Agriculture and Environment for International Development | https://doi.org/10.12895/jaeid.2016110.516 | Benin |
| Müller L-LB et al. | "Are 3°C too much?" – Thermal niche breadth in Bromeliaceae and global warming | Journal of Ecology | https://doi.org/10.1111/1365-2745.12681 | Germany, Panama |
| **Munson SM & Long AL** | **Climate drives shifts in grass reproductive phenology across the western USA** | **New Phytologist** | **https://doi.org/10.1111/nph.14327** | **United States** |
| **Pranovi F et al.** | **The Influence of the Spatial Scale on the Fishery Landings-SST Relationship** | **Frontiers in Marine Science 3:143** | **https://doi.org/10.3389/fmars.2016.00143** | **Italy** |
| Saeedi H et al. | Modelling present and future global distributions of razor clams (Bivalvia: Solenidae) | Helgoland Marine Research 70:23 | https://doi.org/10.1186/s10152-016-0477-4 | New Zealand |
| Sántiz EC et al. | Effect of climate change on the distribution of a critically threatened species | Therya | https://doi.org/10.12933/therya-16-358 | Mexico |

## impacts of climate change

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| **Schleuning M et al.** | **Ecological networks are more sensitive to plant than to animal extinction under climate change** | **Nature Communications 7** | **https://doi.org/10.1038/ncomms13965** | **Germany, Canada, Poland, Switzerland, New Zealand, United Kingdom, Belgium, Serbia, Australia** |
| Stuart-Smith J et al. | Southernmost records of two Seriola species in an Australian ocean-warming hotspot | Marine Biodiversity | https://doi.org/10.1007/s12526-016-0580-4 | Australia, United States |
| Taylor S & Kumar L | Will climate change impact the potential distribution of a native vine (Merremia peltata) which is behaving invasively in the Pacific region? | Ecology and Evolution | https://doi.org/10.1002/ece3.1915 | Australia |
| **Wauchope HS et al.** | **Rapid climate-driven loss of breeding habitat for Arctic migratory birds** | **Global Change Biology** | **https://doi.org/10.1111/gcb.13404** | **Australia, Norway, Russian Federation, Denmark, United States** |
| **Zhang J et al.** | **Extinction risk of North American seed plants elevated by climate and land-use change** | **Journal of Applied Ecology** | **https://doi.org/10.1111/1365-2664.12701** | **Canada, France, Denmark** |

## invasive alien species

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| Geerts S et al. | Lack of human-assisted dispersal means Pueraria montana var. lobata (kudzu vine) could still be eradicated from South Africa | Biological Invasions | https://doi.org/10.1007/s10530-016-1226-y | South Africa |
| Silva DP et al. | Contextualized niche shifts upon independent invasions by the dung beetle Onthophagus taurus | Biological Invasions | https://doi.org/10.1007/s10530-016-1204-4 | Brazil, Spain, Australia, United States |
| **Acosta AL et al.** | **Worldwide Alien Invasion: A Methodological Approach to Forecast the Potential Spread of a Highly Invasive Pollinator** | **PLoS ONE 11(2): e0148295** | **https://doi.org/10.1371/journal.pone.0148295** | **Brazil** |
| **Allen JM & Bradley BA** | **Out of the weeds? Reduced plant invasion risk with climate change in the continental United States** | **Biological Conservation 203: 306-312** | **https://doi.org/10.1016/j.biocon.2016.09.015** | **United States** |
| Attias N et al. | Acácias Australianas no Brasil: Histórico, Formas de Uso e Potencial de Invasão | Biodiversidade Brasileira 3(2): 74-96 | http://bit.ly/2uh3i5F | Brazil |
| Barbosa FG | The future of invasive African grasses in South America under climate change | Ecological Informatics 36: 114-117 | https://doi.org/10.1016/j.ecoinf.2016.10.006 | Brazil |
| **Bellard C et al.** | **Global patterns in threats to vertebrates by biological invasions** | **Proceedings of the Royal Society B 283 (1823)** | **https://doi.org/10.1098/rspb.2015.2454** | **France, Italy, Germany** |
| **Bellard C et al.** | **Major drivers of invasion risks throughout the world** | **Ecosphere 7(3):e01241** | **https://doi.org/10.1002/ecs2.1241** | **France, United Kingdom** |
| Branquart E et al. | A prioritization process for invasive alien plant species incorporating the requirements of EU Regulation no. 1143/2014 | EPPO Bulletin | https://doi.org/10.1111/epp.12336 | Belgium, Italy, Switzerland, United Kingdom, France, Germany, Netherlands |
| Brown KA et al. | Modeling co-occurrence between toxic prey and naïve predators in an incipient invasion | Biodiversity and Conservation | https://doi.org/10.1007/s10531-016-1198-3 | United Kingdom, United States, Madagascar, Canada |
| **Canavan S et al.** | **The global distribution of bamboos: assessing correlates of introduction and invasion** | **AOB Plants** | **https://doi.org/10.1093/aobpla/plw078** | **South Africa, United Kingdom** |
| **Cardador L et al.** | **Combining trade data and niche modelling improves predictions of the origin and distribution of non-native European populations of a globally invasive species** | **Journal of Biogeography** | **https://doi.org/10.1111/jbi.12694** | **Spain** |
| Chai S-L et al. | Using Risk Assessment and Habitat Suitability Models to Prioritise Invasive Species for Management in a Changing Climate | PLoS ONE 11(10) | https://doi.org/10.1371/journal.pone.0165292 | Canada |
| Cheek MD & Semple JC | First official record of naturalised populations of Solidago altissima L. var. pluricephala M.C. Johnst. (Asteraceae: Astereae) in Africa | South African Journal of Botany 105: 333-336 | https://doi.org/10.1016/j.sajb.2016.05.00 | South Africa, Canada |
| Cindi DD & Jaca TP | First record of Opuntia pubescens H.L.Wendland ex Pfeiffer, 1835 naturalised in South Africa | BioInvasions Records 5(4): 213-219 | https://doi.org/10.3391/bir.2016.5.4.04 | South Africa |
| Creed JC et al. | The invasion of the azooxanthellate coral Tubastraea (Scleractinia: Dendrophylliidae) throughout the world: history, pathways and vectors | Biological Invasions | https://doi.org/10.1007/s10530-016-1279-y | Brazil, United States |

# invasive alien species

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| Čuda J et al. | Juvenile biological traits of Impatiens species are more strongly associated with naturalization in temperate climate than their adult traits | Perspectives in Plant Ecology, Evolution and Systematics 20: 1-10 | https://doi.org/10.1016/j.ppees.2016.02.007 | Czech Republic |
| da Silva RS et al. | Potential risk levels of invasive Neoleucinodes elegantalis (small tomato borer) in areas optimal for open-field Solanum lycopersicum (tomato) cultivation in the present and under predicted climate change | Pest Management Science | https://doi.org/10.1002/ps.4344 | Brazil, Australia |
| Di Febbraro M et al. | Modelling the effects of climate change on the risk of invasion by alien squirrels | Hystrix, the Italian Journal of Mammalogy 27 (1) | https://doi.org/10.4404/hystrix-27.1-11776 | Italy |
| **Dullinger I et al.** | **Climate change will increase the naturalization risk from garden plants in Europe** | **Global Ecology and Biogeography** | **https://doi.org/10.1111/geb.12512** | **Austria, Germany, United Kingdom, Czech Republic, France** |
| Escobar LE et al. | Realized niche shift associated with the Eurasian charophyte Nitellopsis obtusa becoming invasive in North America | Scientific Reports 6 | https://doi.org/10.1038/srep29037 | United States, China |
| Figueroa-Castro P et al. | First Report of Peltophorus adustus (Fall) (Coleoptera: Curculionidae: Baridinae) in Mexico, with Two New Host Associations | The Coleopterists Bulletin 70 (3): 667-670 | https://doi.org/10.1649/0010-065X-70.3.667 | Mexico |
| FItecher DH et al. | Predicting global invasion risks: a management tool to prevent future introductions | Scientific Reports 6 | https://doi.org/10.1038/srep26316 | United Kingdom, France |
| Forsyth RG et al. | Distributional Status of an Introduced Land Snail Discus rotundatus (Rotund Disc, Mollusca: Discidae) in Canada | The Canadian Field-Naturalist 130(3): 236-247 | https://doi.org/10.22621/cfn.v130i3.1887 | Canada, Republic of Korea |
| Fried G et al. | MONOGRAPHS ON INVASIVE PLANTS IN EUROPE NO. 1 Monographs on Invasive Plants in Europe: Baccharis halimifolia L. | Botany Letters | https://doi.org/10.1080/23818107.2016.1168315 | France, Spain, Germany, Australia |
| **Gallien L et al.** | **Is There Any Evidence for Rapid, Genetically-Based, Climatic Niche Expansion in the Invasive Common Ragweed?** | **PLoS ONE 11(4): e0152867** | **https://doi.org/10.1371/journal.pone.0152867** | **France** |
| Godefroid M et al. | Is phylogeography helpful for invasive species risk assessment? The case study of the bark beetle genus Dendroctonus? | Ecography | https://doi.org/10.1111/ecog.01474 | France |
| **Harris RMB et al.** | **Unusual suspects in the usual places: a phylo-climatic framework to identify potential future invasive species** | **Biological Invasions** | **https://doi.org/10.1007/s10530-016-1334-8** | **Australia** |
| Hernández-Lambraño RE et al. | Towards the top: niche expansion of Taraxacum officinale and Ulex europaeus in mountain regions of South America | Austral Ecology | https://doi.org/10.1111/aec.12476 | Spain, United Kingdom |
| **Horvitz N et al.** | **Pervasive human-mediated large-scale invasion: analysis of spread patterns and their underlying mechanisms in 17 of China's worst invasive plants** | **Journal of Ecology 105: 85–94** | **https://doi.org/10.1111/1365-2745.12692** | **China, Israel** |
| Hrivnák R et al. | Alien wetland annual Lindernia dubia (Scrophulariaceae): the first recently mentioned localities in Slovakia and their central European context | Biologia 71 (3) | https://doi.org/10.1515/biolog-2016-0039 | Slovakia, Czech Republic, Hungary |
| Januario SM et al. | Combining environmental suitability and population abundances to evaluate the invasive potential of the tunicate Ciona intestinalis along the temperate South American coast | PeerJ 3:e1357 | https://doi.org/10.7717/peerj.1357 | Chile |
| Kanturski M et al. | Pine pest aphids of the genus Eulachnus (Hemiptera: Aphididae: Lachninae): how far can their range extend? | Agricultural and Forest Entomology | https://doi.org/10.1111/afe.12171 | Poland |
| Katsanevakis S et al. | Mapping the impact of alien species on marine ecosystems: the Mediterranean Sea case study | Diversity and Distributions | https://doi.org/10.1111/ddi.12429 | Italy, Greece |
| Keet J-H et al. | Assessing the invasiveness of Berberis aristata and B. julianae (Berberidaceae) in South Africa: Management options and legal recommendations | South African Journal of Botany 105: 288-298 | https://doi.org/10.1016/j.sajb.2016.04.012 | South Africa |
| Kenis M et al. | Assessing the ecological risk posed by a recently established invasive alien predator: Harmonia axyridis as a case study | BioControl | https://doi.org/10.1007/s10526-016-9764-x | Switzerland, Belgium, United Kingdom, United States |
| Laugen AT et al. | The Pacific Oyster (Crassostrea gigas) Invasion in Scandinavian Coastal Waters: Impact on Local Ecosystem Services | Biological Invasions in Changing Ecosystems Vectors, Ecological Impacts, Management and Predictions | https://doi.org/10.1515/9783110438666-015 | Sweden |
| López-Martínez V et al. | Environmental suitability for Agrilus auroguttatus (Coleoptera: Buprestidae) in Mexico using MaxEnt and database records of four Quercus (Fagaceae) species | Agricultural and Forest Entomology | https://doi.org/10.1111/afe.12174 | Mexico, United States |
| **Luizza MW et al.** | **Integrating subsistence practice and species distribution modeling: assessing invasive elodea's potential impact on Native Alaskan subsistence of Chinook salmon and whitefish** | **Environmental Management 1:20** | **https://doi.org/10.1007/s00267-016-0692-4** | **United States** |
| Marchioro CA | Global Potential Distribution of Bactrocera carambolae and the Risks for Fruit Production in Brazil | PLoS ONE 11(11) | https://doi.org/10.1371/journal.pone.0166142 | Brazil |

## invasive alien species

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| Medina AM | What Can an Invasive Dung Beetle Tell Us About Niche Conservatism? | Environmental Ecology | https://doi.org/10.1093/ee/nvw092 | Brazil |
| Merow C et al. | Improving niche and range estimates with Maxent and point process models by integrating spatially explicit information | Global Ecology and Biogeography | https://doi.org/10.1111/geb.12453 | United States |
| Nxumalo MM et al. | Hydrocleys nymphoides (Humb. & Bonpl. ex Willd.) Buchenau: first record of naturalisation in South Africa | BioInvasions Records 5 | https://doi.org/10.3391/bir.2016.5.1.01 | South Africa |
| Pârâu LG et al. | Rose-ringed Parakeet Psittacula krameri Populations and Numbers in Europe: A Complete Overview | The Open Ornithology Journal 9 | https://doi.org/10.2174/1874453201609010001 | Germany, Belgium, Italy, United Kingdom, Netherlands Spain, France, Turkey |
| **PertierraLR et al.** | **Global thermal niche models of two European grasses show high invasion risks in Antarctica** | **Global Change Biology** | **https://doi.org/10.1111/gcb.13596** | **Spain, Australia** |
| Rekha Sarma R et al. | Effect of Climate Change on Invasion Risk of Giant African Snail (Achatina fulica Férussac, 1821: Achatinidae) in India | PLoS ONE 10(11): e0143724 | https://doi.org/10.1371/journal.pone.0143724 | India |
| Rodrigues JFM et al. | Invasion risk of the pond slider turtle is underestimated when niche expansion occurs | Freshwater Biology | https://doi.org/10.1111/fwb.12772 | Brazil, Spain, Germany |
| Scott JK et al. | Methods to select areas to survey for biological control agents: An example based on growth in relation to temperature and distribution of the weed Conyza bonariensis | Biological Control 97: 21-30 | https://doi.org/10.1016/j.biocontrol.2016.02.014 | Australia |
| Su A et al. | Geographic range expansion of alien birds and environmental matching | IBIS | https://doi.org/10.1111/ibi.12418 | United Kingdom, Australia, South Africa |
| Taucare-Ríos A et al. | Using Global and Regional Species Distribution Models (SDM) to Infer the Invasive Stage of Latrodectus geometricus (Araneae: Theridiidae) in the Americas | Ennvironmental Entomology | https://doi.org/10.1093/ee/nvw118 | Chile |
| **Trumbo DR et al.** | **Mixed population genomics support for the central marginal hypothesis across the invasive range of the cane toad (Rhinella marina) in Australia** | **Molecular Ecology** | **https://doi.org/10.1111/mec.13754** | **United States, Australia** |
| **Vall-llosera M et al.** | **Improved surveillance for early detection of a potential invasive species: the alien Rose-ringed parakeet Psittacula krameri in Australia** | **Biological Invasions** | **https://doi.org/10.1007/s10530-016-1332-x** | **Australia, New Zealand** |
| Visser V et al. | Much more give than take: South Africa as a major donor but infrequent recipient of invasive non-native grasses | Global Ecology and Biogeography | https://doi.org/10.1111/geb.12445 | South Africa, Australia |
| **Wan J-Z et al.** | **Impacts of the spatial scale of climate data on the modeled distribution probabilities of invasive tree species throughout the world** | **Ecological Informatics 36: 42-49** | **https://doi.org/10.1016/j.ecoinf.2016.10.001** | **China, United Kingdom** |
| Wan JZ et al. | Risk hotspots for terrestrial plant invaders under climate change at the global scale | Environmental Earth Sciences 75: 1012 | https://doi.org/10.1007/s12665-016-5826-8 | China |
| **Wang Y & Xu Z** | **Where are the Alien Species? Predictions of Global Plant Species Invasions under Current Environmental Conditions and the Human Footprint** | **Polish Journal of Environmental Studies 25 (4): 1729-1738** | **https://doi.org/10.15244/pjoes/62094** | **China** |
| Wang Z-Q et al. | Kalanchoe (Crassulaceae) as invasive aliens in China – new records, and actual and potential distribution | Nordic Journal of Botany | https://doi.org/10.1111/njb.01052 | China, Spain |
| Weyl PSR & Martin GD | Have grass carp driven declines in macrophyte occurrence and diversity in the Vaal River, South Africa? | African Journal of Aquatic Science | https://doi.org/10.2989/16085914.2015.1137856 | South Africa |
| Zeng Y et al. | Novel methods to select environmental variables in MaxEnt: A case study using invasive crayfish | Ecological Modelling 341: 5-13 | https://doi.org/10.1016/j.ecolmodel.2016.09.019 | Singapore |
| Zhu G et al. | Incorporating anthropogenic variables into ecological niche modeling to predict areas of invasion of Popillia japonica | Journal of Pest Science | https://doi.org/10.1007/s10340-016-0780-5 | China |

## species conservation and protected areas

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| Abdou L et al. | Modeling the Geographic Distribution of Prosopis africana (G. and Perr.) Taub. in Niger | Environment and Natural Resources Research 6 (2): 136-144 | https://doi.org/10.5539/enrr.v6n2p136 | Niger |
| Aguilar-López JL et al. | Species Diversity, Distribution, and Conservation Status in a Mesoamerican Region, Amphibians of the Uxpanapa-Chimalapas Region, Mexico | Tropical Conservation Science 9(4) | https://doi.org/10.1177/1940082916670003 | Mexico |
| **Albuquerque F & Beier P** | **Predicted rarity-weighted richness, a new tool to prioritize sites for species representation** | **Ecology and Evolution** | **https://doi.org/10.1002/ece3.2544** | **United States** |
| Ayram CAC et al. | Anthropogenic impact on habitat connectivity: A multidimensional human footprint index evaluated in a highly biodiverse landscape of Mexico | Ecological Indicators 72: 895-909 | https://doi.org/10.1016/j.ecolind.2016.09.007 | Mexico, Colombia |
| **Boavida J et al.** | **Overlooked habitat of a vulnerable gorgonian revealed in the Mediterranean and Eastern Atlantic by ecological niche modelling** | **Scientific Reports 6: 36460** | **https://doi.org/10.1038/srep36460** | **Portugal** |
| Brummitt N et al. | Where are threatened ferns found? Global conservation priorities for pteridophytes | Journal of Systematics and Evolution | https://doi.org/10.1111/jse.12224 | United Kingdom |
| **Butterfield BJ et al.** | **Prestoration: using species in restoration that will persist now and into the future** | **Restoration Ecology** | **https://doi.org/10.1111/rec.12381** | **United States** |
| **Calderón L et al.** | **Genomic evidence of demographic fluctuations and lack of genetic structure across flyways in a long distance migrant, the European turtle dove** | **BMC Evolutionary Biology 16:237** | **https://doi.org/10.1186/s12862-016-0817-7** | **Germany, United States, France, United Kingdom, Spain, Bulgaria, Greece, Malta, Italy** |
| Carneiro LRdA et al. | Limitations to the Use of Species-Distribution Models for Environmental-Impact Assessments in the Amazon | PLoS ONE 11(1): e0146543 | https://doi.org/10.1371/journal.pone.0146543 | Brazil |
| Castellanos-Morales G et al. | Use of molecular and environmental analyses for integrated in situ and ex situ conservation: The case of the Mexican prairie dog | Biological Conservation | https://doi.org/10.1016/j.biocon.2016.10.036 | Mexico |
| Davis SL & Cipollini D | Range, genetic diversity and future of the threatened butterfly, Pieris virginiensis | Insect Conservation and Diversity | https://doi.org/10.1111/icad.12189 | United States |
| **Diesmos AC et al.** | **Amphibians of the Philippines, Part I: Checklist of the Species** | **Proceedings of the California Academy of Sciences 62(20): 457-539** | **http://bit.ly/2t9WQ06** | **Philippines, United States, Malaysia** |
| **dos Santos CF et al.** | **Queens become workers: pesticides alter caste differentiation in bees** | **Scientific Reports 6** | **https://doi.org/10.1038/srep31605** | **Brazil** |
| **Gray CL et al.** | **Local biodiversity is higher inside than outside terrestrial protected areas worldwide** | **Nature Communications 7: 12306** | **https://doi.org/10.1038/ncomms12306** | **United Kingdom, Australia** |
| Hällfors MH et al. | Assessing the need and potential of assisted migration using species distribution models | Biological Conservation 196: 60-68 | https://doi.org/10.1016/j.biocon.2016.01.031 | Finland, United States |
| **Huang D et al.** | **Conservation of reef corals in the South China Sea based on species and evolutionary diversity** | **Biodiversity and Conservation** | **https://doi.org/10.1007/s10531-016-1052-7** | **Singapore, Malaysia, China, Taiwan, Brunei Darussalam, Philippines, Cambodia, Vietnam, Thailand** |
| Idohou R et al. | Spatio-temporal dynamic of suitable areas for species conservation in West Africa: eight economically important wild palms under present and future climates | Agroforestry Systems | https://doi.org/10.1007/s10457-016-9955-6 | Benin, United States |
| Jenkins CN et al. | Patterns of Vertebrate Diversity and Protection in Brazil | PLoS ONE 10(12): e0145064 | https://doi.org/10.1371/journal.pone.0145064 | Brazil |
| Lancaster LT | Widespread range expansions shape latitudinal variation in insect thermal limits | Nature Climate Change | https://doi.org/10.1038/nclimate2945 | United Kingdom |
| **Marshall CAM et al.** | **Bioquality Hotspots in the Tropical African Flora** | **Current Biology 26(23): 3214-3219** | **https://doi.org/10.1016/j.cub.2016.09.045** | **United Kingdom, Netherlands** |
| Martínez-Calderas JM et al. | Potential distribution of margay (Leopardus wiedii, Schinz 1821) in Northeastern Mexico | THERYA 7 (2): 241-255 | https://doi.org/10.12933/therya-16-360 | Mexico |
| McCoshum SM et al. | Species distribution models for natural enemies of monarch butterfly (Danaus plexippus) larvae and pupae: distribution patterns and implications for conservation | Journal of Insect Conservation | https://doi.org/10.1007/s10841-016-9856-z | United States |
| Medina RG et al. | Environmental, land cover and land use constraints on the distributional patterns of anurans: Leptodactylus species (Anura, Leptodactylidae) from Dry Chaco | PeerJ 4 | https://doi.org/10.7717/peerj.2605 | Argentina |

## species conservation and protected areas

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| **Mendoza-González G et al.** | **Priority areas for conservation of beach and dune vegetation of the Mexican Atlantic coast** | **Journal for Nature Conservation** | **https://doi.org/10.1016/j.jnc.2016.04.007** | **Mexico, United States** |
| Miller MH | Status Review Report of 3 Species of Angelsharks: Squatina aculeata, S. oculata, and S. squatina | Report to National Marine Fisheries Service, Office of Protected Resources | http://bit.ly/2uhmy2K | United States |
| Obregón R et al. | Effects of climate change on three species of Cupido (Lepidoptera, Lycaenidae) with different biogeographic distribution in Andalusia, southern Spain | Animal Biodiversity and Conservation, 39.1: 115-128 | http://bit.ly/2uh1cCJ | Spain |
| Osipova L & Sangermano F | Surrogate species protection in Bolivia under climate and land cover change scenarios | Journal for Nature Conservation 34: 107-117 | https://doi.org/10.1016/j.jnc.2016.10.002 | United States |
| **Ostrowski M-F et al.** | **Potential Implications of Climate Change on Aegilops Species Distribution: Sympatry of These Crop Wild Relatives with the Major European Crop Triticum aestivum and Conservation Issues** | **PLoS ONE 11(4): e0153974** | **https://doi.org/10.1371/journal.pone.0153974** | **France** |
| Peterson AT & Navarro-Sigüenza AG | Bird conservation and biodiversity research in Mexico: status and priorities | Journal of Field Ornithology | https://doi.org/10.1111/jofo.12146 | United States, Mexico |
| Peterson MS et al. | Distribution, Abundance, and Habitat Characteristics of Fundulus jenkinsi (Evermann) (Saltmarsh Topminnow) in Coastal Mississippi Watersheds, with Comments on Range-Wide Occurrences Based on Non-Vouchered and Museum Records | Southeastern Naturalist 15(3):415-430 | https://doi.org/10.1656/058.015.0304 | United States |
| **Phillips J et al.** | **In situ and ex situ diversity analysis of priority crop wild relatives in Norway** | **Diversity and Distributions** | **https://doi.org/10.1111/ddi.12470** | **United Kingdom, Norway** |
| Pokharel KP et al. | Predicting potential distribution of poorly known species with small database: the case of four-horned antelope Tetracerus quadricornis on the Indian subcontinent. | Ecology and Evolution | https://doi.org/10.1002/ece3.2037 | Germany |
| Reichel K et al. | Genetic diversity in the locally declining Laserpitium prutenicum L. and the more common Selinum carvifolia (L.) L.: a "silent goodbye"? | Conservation Genetics | https://doi.org/10.1007/s10592-016-0827-4 | Germany, France, Poland |
| Rhoades PR et al. | Evidence for Bombus occidentalis (Hymenoptera: Apidae) Populations in the Olympic Peninsula, the Palouse Prairie, and Forests of Northern Idaho | Journal of Insect Science | https://doi.org/10.1093/jisesa/iev155 | United States |
| **Solberg SØ & Chou YY** | **Conservation of Indigenous Vegetables from a Hotspot in Tropical Asia: What Did We Learn from Vavilov?** | **Frontiers in Plant Science** | **https://doi.org/10.3389/fpls.2016.01982** | **Taiwan** |
| Speed JDM & Austrheim G | The importance of herbivore density and management as determinants of the distribution of rare plant species | Biological Conservation 205: 77-84 | https://doi.org/10.1016/j.biocon.2016.11.030 | Norway |
| Stratmann TSM et al. | Locating Suitable Habitat for a Rare Species: Evaluation of a Species Distribution Model for Bog Turtles (Glyptemys Muhlenbergii) in the Southeastern United States | Herpetological Conservation and Biology 11(1):199–213 | http://bit.ly/2ugZ65K | United States, Germany |
| Thompson DM et al. | Effects of life-history requirements on the distribution of a threatened reptile | Conservation Biology | https://doi.org/10.1111/cobi.12800 | United States |
| **Tolley KA et al.** | **Conservation status and threats for African reptiles** | **Biological Conservation** | **https://doi.org/10.1016/j.biocon.2016.04.006** | **South Africa, United States** |
| Toranza C et al. | Threatened amphibians of Uruguay: Effectiveness of protected areas face of climate change | Ecologia Austral 26(2):138-149 | http://bit.ly/2uhbiDK | Uruguay |
| **Trejo-Salazar R-E et al.** | **Save Our Bats, Save Our Tequila: Industry and Science Join Forces to Help Bats and Agaves** | **Natural Areas Journal 36(4):523-530** | **https://doi.org/10.3375/043.036.0417** | **Mexico, United States** |
| **Trowbridge CD et al.** | **Shallow subtidal octocorals in an Irish marine reserve** | **Marine Biodiversity** | **https://doi.org/10.1007/s12526-016-0450-0** | **United States, United Kingdom, Ireland** |
| Vroh BTA et al. | Comparing Suitable Habitat Models to Predict Rare and Endemic Plant Species Distributions: What are the Limits of the Niche of Cola lorougnonis (Malvaceae) in Cote d'Ivoire? | Environment and Natural Resources Research 6(3) | https://doi.org/10.5539/enrr.v6n3p1 | Ivory Coast |
| Wang CJ et al. | Identifying appropriate protected areas for endangered fern species under climate change | SpringerPlus 5: 904 | https://doi.org/10.1186/s40064-016-2588-4 | China |
| Wehrtmann IS et al. | The primary freshwater crabs of Guatemala (Decapoda: Brachyura: Pseudothelphusidae), with comments on their conservation status | Journal of Crustacean Biology | https://doi.org/10.1163/1937240X-00002478 | Costa Rica, Brazil, Guatemala |

## species conservation and protected areas

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| Zhu GP et al. | Mapping the ecological dimensions and potential distributions of endangered relic shrubs in western Ordos biodiversity center | Scientific Reports 6 | https://doi.org/10.1038/srep26268 | China |

## advancing biodiversity science

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| Aagesen L et al. | Macro-Climatic Distribution Limits Show Both Niche Expansion and Niche Specialization among C4 Panicoids | PLoS ONE 11(3): e0151075 | https://doi.org/10.1371/journal.pone.0151075 | Argentina |
| Aguirre-Santoro J et al. | Molecular phylogenetics of the Ronnbergia Alliance (Bromeliaceae, Bromelioideae) and insights into their morphological evolution | Molecular Phylogenetics and Evolution 100: 1-20 | https://doi.org/10.1016/j.ympev.2016.04.007 | United States |
| Aliabadian M et al. | Phylogeny, biogeography, and diversification of barn owls (Aves: Strigiformes) | Biological Journal of the Linnean Society | https://doi.org/10.1111/bij.12824 | Iran, United Kingdom, Switzerland |
| Almeida TE & Salino A | State of the art and perspectives on neotropical fern and lycophyte systematics | Journal of Systematics and Evolution | https://doi.org/10.1111/jse.12223 | Brazil |
| **André T et al.** | **Speciation dynamics and biogeography of Neotropical spiral gingers (Costaceae)** | **Molecular Phylogenetics and Evolution 103: 55-63** | **https://doi.org/10.1016/j.ympev.2016.07.008** | **United States, Brazil** |
| Araújo RM et al. | Status, trends and drivers of kelp forests in Europe: an expert assessment | Biodiversity and Conservation 25 (7): 1319-1348 | https://doi.org/10.1007/s10531-016-1141-7 | Portugal, Italy, Spain, Germany, Norwey, France, United Kingdom, Australia |
| Arias-Alzate A et al. | Wild Felid Range Shift Due to Climatic Constraints in the Americas: a Bottleneck Explanation for Extinct Felids? | Journal of Mammalian Evolution | https://doi.org/10.1007/s10914-016-9350-0 | Mexico, Colombia |
| Bacon CD et al. | Species limits, geographical distribution and genetic diversity in Johannesteijsmannia (Arecaceae) | The Linnean Society of London, Botanical Journal of the Linnean Society | https://doi.org/10.1111/boj.12470 | Sweden, Colombia, Singapore, Malaysia, United Kingdom |
| **Ballesteros-Mejia L et al.** | **Putting insects on the map: Near-global variation in sphingid moth richness along spatial and environmental gradients** | **Ecography** | **https://doi.org/10.1111/ecog.02438** | **Brazil, United Kingdom, United States, Switzerland** |
| Batalha-Filho H & Miyaki C Y | Late Pleistocene divergence and postglacial expansion in the Brazilian Atlantic Forest: multilocus phylogeography of Rhopias gularis (Aves: Passeriformes) | Journal of Zoological Systematics and Evolutionary Research | https://doi.org/10.1111/jzs.12118 | Brazil |
| **Bateman BL et al.** | **Potential breeding distributions of U.S. birds predicted with both short-term variability and long-term average climate data** | **Ecological Applications** | **https://doi.org/10.1002/eap.1416** | **United States, Australia** |
| Bean WT & Osborn SD | A Suitability Model for White-Footed Voles with Insights into Habitat Associations at the Southern Boundary of Their Range | Northwestern Naturalist 97(2):105-112 | https://doi.org/10.1898/NWN15-28.1 | United States |
| Beier P & Albuquerque F | Evaluating Diversity as a Surrogate for Species Representation at Fine Scale | PLoS ONE 11(3) | https://doi.org/10.1371/journal.pone.0151048 | United States |
| Bellot S et al. | Assembled plastid and mitochondrial genomes, as well as nuclear genes, place the parasite family Cynomoriaceae in the Saxifragales | Genome Biology and Evolution | https://doi.org/10.1093/gbe/evw147 | Germany, China, Iran, Austria |
| **Benedict JC et al.** | **Species diversity driven by morphological and ecological disparity: a case study of seeds of Zingiberales (bananas, gingers, and relatives)** | **AoB PLANTS** | **https://doi.org/10.1093/aobpla/plw063** | **United States, United Kingdom, Singapore, Switzerland** |
| Beukhof ED et al. | Records of five bryozoan species from offshore gas platforms rare for the Dutch North Sea | Marine Biodiversity Records 9:91 | https://doi.org/10.1186/s41200-016-0086-6 | Denmark, Netherlands, Belgium |
| Bocksberger G et al. | Climate and the distribution of grasses in West Africa | Journal of Vegetation Science | https://doi.org/10.1111/jvs.12360 | Germany, Switzerland, France, Burkina Faso |
| Bocsi T et al. | Plants' native distributions do not reflect climatic tolerance | Diversity and Distributions | https://doi.org/10.1111/ddi.12432 | United States |
| Boucher FC et al. | Evolution and biogeography of the cushion life form in angiosperms | Perspectives in Plant Ecology, Evolution and Systematics 20: 22-31 | https://doi.org/10.1016/j.ppees.2016.03.002 | France, Switzerland |
| Brandt LA et al. | Comparison of climate envelope models developed using expert-selected variables versus statistical selection | Ecological Modelling 345: 10-20 | https://doi.org/10.1016/j.ecolmodel.2016.11.016 | United States |
| Brewer MJ et al. | Plateau: a new method for ecologically plausible climate envelopes for species distribution modelling | Methods in Ecology and Evolution | https://doi.org/10.1111/2041-210X.12609 | United Kingdom, Germany, New Zealand |

## advancing biodiversity science

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| Brischoux F et al. | Oceanic circulation models help to predict global biogeography of pelagic yellow-bellied sea snake | Biology Letters | https://doi.org/10.1098/rsbl.2016.0436 | France, United States, Australia |
| **Brock JMR et al.** | **Tree fern ecology in New Zealand: A model for southern temperate rainforests** | **Forest Ecology and Management 375: 112-126** | **https://doi.org/10.1016/j.foreco.2016.05.030** | **New Zealand** |
| Bush MB & McMichael CNH | Holocene Variability of an Amazonian Hyperdominant | Journal of Ecology | https://doi.org/10.1111/1365-2745.12600 | United States, Netherlands |
| Butterfield BJ & Munson SM | Temperature is better than precipitation as a predictor of plant community assembly across a dryland region | Journal of Vegetation Science | https://doi.org/10.1111/jvs.12440 | United States |
| Cáceres NC et al. | Which Factors Determine Spatial Segregation in the South American Opossums (Didelphis aurita and D. albiventris)? An Ecological Niche Modelling and Geometric Morphometrics Approach | PLoS ONE 11(6): e0157723 | https://doi.org/10.1371/journal.pone.0157723 | Brazil, United Kingdom |
| Cadena CD et al. | Climate, habitat associations and the potential distributions of Neotropical birds: Implications for diversification across the Andes | Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales | https://doi.org/10.18257/raccefyn.280 | Colombia, United States |
| Cai L et al. | Phylogeny of Elatinaceae and the Tropical Gondwanan Origin of the Centroplacaceae(Malpighiaceae, Elatinaceae) Clade | PLoS ONE 11(9) | https://doi.org/10.1371/journal.pone.0161881 | United States |
| Calleja JA et al. | Epiphytic Bryophyte Communities of Prunus lusitanica Iberian Forests: Biogeographic Islands Shaped by Regional Climates | Cryptogamie, Bryologie 37(1):53-85 | https://doi.org/10.7872/cryb/v37.iss1.2016.53 | Spain |
| Caperta AD et al. | Biogeographical, ecological and ploidy variation in related asexual and sexual Limonium taxa (Plumbaginaceae) | Botanical Journal of the Linnean Society, | https://doi.org/10.1111/boj.12498 | Portugal, Morocco |
| Cardinal-McTeague WM et al. | Biogeography and diversification of Brassicales: A 103 million year tale | Molecular Phylogenetics and Evolution | https://doi.org/10.1016/j.ympev.2016.02.021 | United States, Canada |
| Cardoso AL et al. | Integrated Cytogenetic and Mitochondrial DNA Analyses Indicate That Two Different Phenotypes of Hypancistrus (L066 and L333) Belong to the Same Species | Zebrafish | https://doi.org/10.1089/zeb.2015.1213 | Brazil |
| Chakrabarty P et al. | Five Years Later: An Update on the Status of Collections of Endemic Gulf of Mexico Fishes Put at Risk by the 2010 Oil Spill | Biodiversity Data Journal 4: e8728 | https://doi.org/10.3897/BDJ.4.e8728 | United States |
| **Chandler M et al.** | **Contribution of citizen science towards international biodiversity monitoring** | **Biological Conservation** | **https://doi.org/10.1016/j.biocon.2016.09.004** | **United States, Austria, Denmark, Spain, Australia** |
| Charles-Dominique T et al. | Spiny plants, mammal browsers, and the origin of African savannas | PNAS | https://doi.org/10.1073/pnas.1607493113 | South Africa, Canada, United States |
| Cheddadi R & Khater C | Climate change since the last glacial period in Lebanon and the persistence of Mediterranean species | Quaternary Science Reviews 150: 146-157 | https://doi.org/10.1016/j.quascirev.2016.08.010 | France, Lebanon |
| Chernomor O et al. | Split Diversity: Measuring and Optimizing Biodiversity Using Phylogenetic Split Networks | in Biodiversity Conservation and Phylogenetic Systematics 14: 173-195 | https://doi.org/10.1007/978-3-319-22461-9_9 | Austria, New Zealand |
| Congrains C et al. | Genetic and paleomodelling evidence of the population expansion of the cattle egret Bubulcus ibis in Africa during the climatic oscillations of the Late Pleistocene | Journal of Avian Biology | https://doi.org/10.1111/jav.00972 | Brazil, South Africa, Australia, Nigeria, Guinea_Bissau, Ghana |
| **Coro G et al.** | **Estimating absence locations of marine species from data of scientific surveys in OBIS** | **Ecological Modelling 323:61-76** | **https://doi.org/10.1016/j.ecolmodel.2015.12.008** | **Italy, Belgium, Greece, Philippines** |
| **Cramer MD & Verboom GA** | **Measures of biologically relevant environmental heterogeneity improve prediction of regional plant species richness** | **Journal of Biogeography** | **https://doi.org/10.1111/jbi.1291** | **South Africa** |
| Cresson P et al. | Lost in the North: The first record of Diretmichthys parini (Post and Quéro, 1981) in the northern North Sea | Marine Pollution Bulletin | https://doi.org/10.1016/j.marpolbul.2016.10.074 | France |
| **Crowl AA et al.** | **A global perspective on Campanulaceae: Biogeographic, genomic, and floral evolution** | **American Journal of Botany** | **https://doi.org/10.3732/ajb.1500450** | **United States** |
| Cruz JA et al. | Reconstructing the paleoenvironment of Loltún Cave, Yucatán, Mexico, with Pleistocene amphibians and reptiles and their paleobiogeographic implications | Revista Mexicana de Ciencias Geologicas | http://bit.ly/2uh6Rsh | Mexico |
| Cuéllar-Martínez M & Sosa V | Diversity patterns of monocotiledonous geophytes in Mexico | Botanical Sciences 94(4): 1-13 | https://doi.org/10.17129/botsci.763 | Mexico |
| **Curtis CA & Bradley BA** | **Plant Distribution Data Show Broader Climatic Limits than Expert-Based Climatic Tolerance Estimates** | **PLoS ONE 11(11): e0166407** | **https://doi.org/10.1371/journal.pone.0166407** | **United States** |
| Dal Vechio F et al. | Taxonomic Status and the Phylogenetic Placement of Amphisbaena leucocephala Peters, 1878 (Squamata, Amphisbaenidae) | South American Journal of Herpetology 11(3):157-175 | https://doi.org/10.2994/SAJH-D-16-00027.1 | Brazil |

## advancing biodiversity science

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| Davies MR & Piontek S | The marine fishes of St. Eustatius, Dutch Caribbean | Marine Biodiversity | https://doi.org/10.1007/s12526-016-0575-1 | Netherlands |
| Davis TR | First records of three fishes, and southern records of a further four fishes, from New South Wales, Australia | Check List 12(6) | https://doi.org/10.15560/12.6.2008 | Australia |
| de Almeida DAC et al. | Taxonomy and molecular phylogeny of Diatrypaceae (Ascomycota, Xylariales) species from the Brazilian semi-arid region, including four new species | Mycological Progress | https://doi.org/10.1007/s11557-016-1194-8 | Brazil, United States |
| De La Cruz-Agüero J et al. | Unusual records of deepwater teleosteans trawled off the western coast of Mexico | Journal of Fish Biology | https://doi.org/10.1111/jfb.13087 | Mexico |
| de Melo WA et al. | Coalescent Simulation and Paleodistribution Modeling for Tabebuia rosealba Do Not Support South American Dry Forest Refugia Hypothesis | PLoS ONE 11(7): e0159314 | https://doi.org/10.1371/journal.pone.0159314 | Brazil |
| de Moura Bubadué J et al. | Character displacement under influence of Bergmann's rule in Cerdocyon thous (Mammalia: Canidae) | Hystrix, teh Italian Journal of Mammalogy | https://doi.org/10.4404/hystrix-27.2-11433 | Brazil, Italy |
| **Deblauwe V et al.** | **Remotely sensed temperature and precipitation data improve species distribution modelling in the tropics** | **Global Ecology and Biogeography** | **https://doi.org/10.1111/geb.12426** | **France, Belgium, Cameroon, United States, India, Denmark, Netherlands** |
| Depa Ł et al. | Do ants drive speciation in aphids? A possible case of ant-driven speciation in the aphid genus Stomaphis Walker (Aphidoidea, Lachninae) | Zoological Journal of the Linnean Society | https://doi.org/10.1111/zoj.12437 | Poland |
| DeVaney SC | Species Distribution Modeling of Deep Pelagic Eels | Integrative & Comparative Biology | https://doi.org/10.1093/icb/icw032 | United States |
| Devi KT et al. | Three new additions to the grass (Poaceae) flora of Manipur, India | Plant Science Today 3(3): 272-281 | https://doi.org/10.14719/pst.2016.3.3.237 | India |
| Dianat M et al. | Evolutionary history of the Persian Jird, Meriones persicus, based on genetics, species distribution modelling and morphometric data | Journal of Zoological Systematics and Evolutionary Research | https://doi.org/10.1111/jzs.12145 | Iran, France |
| Domínguez MC et al. | Vicariance events shaping Southern South American insect distributions | Zoologica Scripta | https://doi.org/10.1111/zsc.12167 | Argentina |
| Drees C et al. | Molecular analyses and species distribution models indicate cryptic northern mountain refugia for a forest-dwelling ground beetle | Journal of Biogeography | https://doi.org/10.1111/jbi.12828 | Germany, Netherlands, Kenya |
| Dufresnes C et al. | Evolutionary melting pots: a biodiversity hotspot shaped by ring diversifications around the Black Sea in the Eastern tree frog (Hyla orientalis) | Molecular Ecology | https://doi.org/10.1111/mec.13706 | Switzerland, Russian Federation, Ukraine, Germany |
| Dugdale TM et al. | The biology of Australian weeds 65. 'Tradescantia fluminensis' Vell | Plant Protection Quarterly 30(4) | http://bit.ly/2uhukKe | Australia |
| Feng X et al. | Climatic Similarity of Extant and Extinct Dasypus Armadillos | Journal of Mammalian Evolution | https://doi.org/10.1007/s10914-016-9336-y | United States, Brazil |
| Ferreira GE et al. | Subtropical species of Sinningia (Gesneriaceae): Distribution patterns and limiting environmental factors | Flora - Morphology, Distribution, Functional Ecology of Plants 222: 86-95 | https://doi.org/10.1016/j.flora.2016.04.002 | Brazil, Switzerland |
| Ferrer-Paris JR et al. | Indicative response of Oxysternon festivum Linné (Coleoptera: Scarabaidae) to vegetation condition in the basin of the Orinoco river, Venezuela | Journal of Insect Conservation 20 (3): 527-538 | https://doi.org/10.1007/s10841-016-9886-6 | Venezuela |
| Figueroa LL & Bergey EA | Bumble Bees (Hymenoptera: Apidae) of Oklahoma: Past and Present Biodiversity | Journal of the Kansas Entomological Society 88(4):418-429 | https://doi.org/10.2317/0022-8567-88.4.418 | United States |
| Foody GM et al. | The Sensitivity of Mapping Methods to Reference Data Quality: Training Supervised Image Classifications with Imperfect Reference Data | ISPRS International Journal of Geo-Information 5(11) | https://doi.org/10.3390/ijgi5110199 | United Kingdom, India, Italy, France |
| **Fourcade Y** | **Comparing species distributions modelled from occurrence data and from expert-based range maps. Implication for predicting range shifts with climate change** | **Ecological Informatics 36: 8-14** | **https://doi.org/10.1016/j.ecoinf.2016.09.002** | **Sweden** |
| **Fournier AMV et al.** | **Combining citizen science species distribution models and stable isotopes reveals migratory connectivity in the secretive Virginia rail** | **Journal of Applied Ecology** | **https://doi.org/10.1111/1365-2664.12723** | **United States, Sweden, Canada** |
| Fournier AMV et al. | Combining citizen science species distribution models and stable isotopes reveals migratory connectivity in the secretive Virginia rail | Journal of Applied Ecology | https://doi.org/10.1111/1365-2664.12723 | United States, Sweden |
| Fuchs J et al. | Long-distance dispersal and inter-island colonization across the western Malagasy Region explain diversification in brush-warblers (Passeriformes: Nesillas | Biological Journal of the Linnean Society, | https://doi.org/10.1111/bij.12825 | France, Colombia, Madagascar, United Kingdom, Switzerland, United States |

## advancing biodiversity science

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| Gaddis KD et al. | Dry-washes determine gene flow and genetic diversity in a common desert shrub | Landscape Ecology | https://doi.org/10.1007/s10980-016-0393-7 | United States |
| García-Redondo V et al. | First Record of Sexual Structures in Pterosiphonia Parasitica (Rhodomelaceae, Rhodophyta) from the Iberian Peninsula | Thalassas: An International Journal of Marine Sciences | https://doi.org/10.1007/s41208-016-0012-z | Spain |
| Gherghel I et al. | A revision of the distribution of sea kraits (Reptilia, Laticauda) with an updated occurrence dataset for ecological and conservation research | ZooKeys 569: 135-148 | https://doi.org/10.3897/zookeys.569.6975 | United States, France, Romania |
| Gómez C et al. | Niche-tracking migrants and niche-switching residents: evolution of climatic niches in New World warblers (Parulidae) | Proceedings of the Royal Society B 283 (1823) | https://doi.org/10.1098/rspb.2015.2458 | Colombia |
| **Gómez-Ruiz EP & Lacher TE** | **Modelling the potential geographic distribution of an endangered pollination corridor in Mexico and the United States** | **Diversity and Distributions** | **https://doi.org/10.1111/ddi.12499** | **Mexico, United States** |
| González JA | Brachyuran crabs (Crustacea: Decapoda) from the Canary Islands (eastern Atlantic): checklist, zoogeographic considerations and conservation | Scientia Marina 80(1) | https://doi.org/10.3989/scimar.04350.10A | Spain |
| González JA et al. | Spatio-temporal point process statistics: A review | Spatial Statistics | https://doi.org/10.1016/j.spasta.2016.10.002 | Spain, Sweden |
| Gouveia SF & Correia I | Geographical clines of body size in terrestrial amphibians: water conservation hypothesis revisited | Journal of Biogeography | https://doi.org/10.1111/jbi.12842 | Brazil |
| Greve M et al. | Realising the potential of herbarium records for conservation biology | South African Journal of Botany 105: 317-323 | https://doi.org/10.1016/j.sajb.2016.03.017 | Denmark, South Africa, Brazil, United States, United Kingdom, Belgium, Burundi |
| **Griffiths HJ & Waller CL** | **The first comprehensive description of the biodiversity and biogeography of Antarctic and Sub-Antarctic intertidal communities** | **Journal of Biogeography** | **https://doi.org/10.1111/jbi.12708** | **United Kingdom** |
| Grünwald CI et al. | A new species of Ophryacus (Serpentes: Viperidae: Crotalinae) from eastern Mexico with comments of the taxonomy of related pitvipers | Mesoamerican Herpetology 2: 388-416 | http://bit.ly/2dkggtr | Mexico, United States |
| Guan B-C et al. | Landscape connectivity of Cercidiphyllum japonicum, an endangered species and its implications for conservation | Ecological Informatics 33: 51-66 | https://doi.org/10.1016/j.ecoinf.2016.04.002 | China |
| **Gueta T & Carmel Y** | **Quantifying the value of user-level data cleaning for: A case study using mammal distribution models** | **Ecological Informatics 34: 139-145** | **https://doi.org/10.1016/j.ecoinf.2016.06.001** | **Israel** |
| Hantemirova EV et al. | A new Eurasian phylogeographical paradigm? Limited contribution of southern populations to the recolonization of high latitude populations in Juniperus communis L. (Cupressaceae) | Journal of Biogeography | https://doi.org/10.1111/jbi.12867 | Russian Federation, Austria, Sweden |
| Hargrave CW & Gary KP | Historical Distribution of Bluehead Shiner (Pteronotropis hubbsi) | Southeastern Naturalist 15(sp9):110-116 | https://doi.org/10.1656/058.015.0sp913 | United States |
| Henriques Antão L et al. | Prevalence of multimodal species abundance distributions is linked to spatial and taxonomic breadth | Global Ecology and Biogeography | https://doi.org/10.1111/geb.12532 | United Kingdom, Australia, Portugal |
| Henrot A-J et al. | Middle Miocene climate and vegetation models and their validation with proxy data | Palaeogeography, Palaeoclimatology, Palaeoecology | https://doi.org/10.1016/j.palaeo.2016.05.026 | Belgium, Germany, Hungary, Australia, United States |
| Hernández-Baz F et al. | Notes on the Behavior and Distribution of the Day-Flying Moth, Heterusia atalantata (Guenée, [1858]) (Lepidoptera, Geometridae, Larentiinae), with Special Reference to Mexico | Journal of the Lepidopterists' Society 70(4):283-288 | https://doi.org/10.18473/lepi.70i4.a6 | United States, Mexico |
| Hernández-Velasco A et al. | Occurrence of Holacanthus clarionensis (Pomacanthidae), Stegastes leucorus, and Stegastes acapulcoensis (Pomacentridae) at Magdalena Bay, B.C.S., Mexico | Marine Biodiversity Records 9:49 | https://doi.org/10.1186/s41200-016-0062-1 | Mexico |
| Hultine KR et al. | Stable isotope physiology of stem succulents across a broad range of volume-to-surface area ratio | Oecologia | https://doi.org/10.1007/s00442-016-3690-6 | United States |
| Ivory SJ et al. | Niche expansion and temperature sensitivity of tropical African montane forests | Global Ecology and Biogeography | https://doi.org/10.1111/geb.12446 | United States, United Kingdom |
| Jaffé R et al. | Beekeeping practices and geographic distance, not land use, drive gene flow across tropical bees | Molecular Ecology | https://doi.org/10.1111/mec.13852 | Brazil, United States, Spain, Portugal |
| **Janssens SB et al.** | **Evolutionary dynamics and biogeography of Musaceae reveal a correlation between the diversification of the banana family and the geological and climatic history of Southeast Asia** | **New Phytologist** | **https://doi.org/10.1111/nph.13856** | **Belgium, Netherlands, Tanzania** |
| JC Ganglo & SB Kakpo | Completeness of Digital Accessible Knowledge of Plants of Benin and Priorities for Future Inventory and Data Discovery | Biodiversity Informatics 11: 23-39 | https://doi.org/10.17161/bi.v11i0.5053 | Benin |

## advancing biodiversity science

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| Jia S-W et al. | Evolutionary history of Gymnocarpos (Caryophyllaceae) in the arid regions from North Africa to Central Asia | The Linnean Society of London, Biological Journal of the Linnean Society, | https://doi.org/10.1111/bij.12834 | China, Germany, Sweden |
| Jiménez RA & Ornelas JF | Historical and current introgression in a Mesoamerican hummingbird species complex: a biogeographic perspective | PeerJ 4:e1556 | https://doi.org/10.7717/peerj.1556 | Mexico |
| Jiménez-Alfaro B et al. | Anticipating extinctions of glacial relict populations in mountain refugia | Biological Conservation 201: 243-251 | https://doi.org/10.1016/j.biocon.2016.07.015 | Czech Republic, Spain |
| Jung J-M et al. | Insect distribution in response to climate change based on a model: Review of function and use of CLIMEX | Entomological Research | https://doi.org/10.1111/1748-5967.12171 | Korea |
| Keller B et al. | Both morph- and species-dependent asymmetries affect reproductive barriers between heterostylous species | Ecology and Evolution | https://doi.org/10.1002/ece3.2293 | Switzerland, United States, Australia, Canada |
| Kennedy MP et al. | Niche-breadth of freshwater macrophytes occurring in tropical southern African rivers predicts species global latitudinal range | Aquatic Botany 136: 21-30 | https://doi.org/10.1016/j.aquabot.2016.09.003 | United Kingdom, Zambia, South Africa, Argentina, Brazil |
| **Keppel G et al.** | **Habitat diversity predicts orchid diversity in the tropical south-west Pacific** | **Journal of Biogeography** | **https://doi.org/10.1111/jbi.12805** | **Australia, United States** |
| Kissling WD et al. | Historical colonization and dispersal limitation supplement climate and topography in shaping species richness of African lizards (Reptilia: Agaminae) | Scientific Reports 6 | https://doi.org/10.1038/srep34014 | Netherlands, Denmark, Germany, United States |
| Koch JB et al. | Range Extension of Two Bumble Bee Species (Hymenoptera: Apidae) into Olympic National Park | Northwest Science 90(2):228-234 | https://doi.org/10.3955/046.090.0212 | United States |
| **Koski et al.** | **Reproductive character displacement and environmental filtering shape floral variation between sympatric sister taxa** | **Evolution** | **https://doi.org/10.1111/evo.13042** | **United States** |
| Koski MH & Ashman T-L | Macroevolutionary patterns of ultraviolet floral pigmentation explained by geography and associated bioclimatic factors | New Phytologist | https://doi.org/10.1111/nph.13921 | United States |
| Kramer-Schadt S et al. | The Borneo carnovore database and the application of predictive dstribution modelling | Raffles Bulletin of Zoology 33: 18-41 | http://bit.ly/2ugZIIA | Germany |
| Kratochwil A | Review of the Icelandic bee fauna (Hymenoptera: Apoidea: Anthophila) | Stuttgarter Beiträge zur Naturkunde A 9(1):217-227 | https://doi.org/10.18476/sbna.v9.a14 | Germany |
| Krehenwinkel H et al. | Rapid genetic and ecological differentiation during the northern range expansion of the venomous yellow sac spider Cheiracanthium punctorium in Europe | Evolutionary Applications | https://doi.org/10.1111/eva.12392 | Germany, United States, Romania, Slovenia |
| Krehenwinkel H et al. | A phylogeographical survey of a highly dispersive spider reveals eastern Asia as a major glacial refugium for Palaearctic fauna | Journal of Biogeography | https://doi.org/10.1111/jbi.12742 | Germany, United States, Japan |
| Krishnan A & Tamma K | Divergent morphological and acoustic traits in sympatric communities of Asian barbets | Royal Society Open Science | https://doi.org/10.1098/rsos.160117 | United States, India |
| Kübler D et al. | Assessing the importance of topographic variables for the spatial distribution of tree species in a tropical mountain forest | Erdkunde 70 (1): 19-47 | https://doi.org/10.3112/erdkunde.2016.01.03 | Germany, Ecuador |
| **La Salle J, Williams KJ & Moritz C** | **Biodiversity analysis in the digital era** | **Philosophical Transactions B 371 (1702)** | **https://doi.org/10.1098/rstb.2015.0337** | **Australia** |
| Lagomarsino LP et al. | The abiotic and biotic drivers of rapid diversification in Andean bellflowers (Campanulaceae) | New Phytologist | https://doi.org/10.1111/nph.13920 | United States, Sweden, Germany |
| Langejans GHJ et al. | Pleistocene molluscs from Klasies River (South Africa): Reconstructing the local coastal environment | Quaternary International | https://doi.org/10.1016/j.quaint.2016.01.013 | Netherlands, South Africa |
| **Leach K et al.** | **Modelling the influence of biotic factors on species distribution patterns** | **Ecological Modelling 337: 96-106** | **https://doi.org/10.1016/j.ecolmodel.2016.06.008** | **United Kingdom** |
| **Lee-Yaw JA et al.** | **A synthesis of transplant experiments and ecological niche models suggests that range limits are often niche limits** | **Ecology Letters** | **https://doi.org/10.1111/ele.12604** | **Canada, Switzerland, United States, Ireland** |
| Leite YLR et al. | Neotropical forest expansion during the last glacial period challenges refuge hypothesis | PNAS, Proceedings of the National Academy of Sciences of the United States of America | https://doi.org/10.1073/pnas.1513062113 | Brazil, Portugal |
| Lens F et al. | Scalariform-to-simple transition in vessel perforation plates triggered by differences in climate during the evolution of Adoxaceae | Annals of Botany | https://doi.org/10.1093/aob/mcw151 | Netherlands, France, South Africa, Belgium |
| Lessmann J et al. | Freshwater vertebrate and invertebrate diversity patterns in an Andean-Amazon basin: implications for conservation efforts | Neotropical Biodiversity 2(1) | https://doi.org/10.1080/23766808.2016.1222189 | United States, Ecuador, Portugal |

## advancing biodiversity science

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| Li G et al. | Mapping the climatic suitable habitat of oriental arborvitae (Platycladus orientalis) for introduction and cultivation at a global scale | Scientific Reports 6: 30009 | https://doi.org/10.1038/srep30009 | China |
| Lin LH & Wiens JJ | Comparing macroecological patterns across continents: evolution of climatic niche breadth in varanid lizards | Ecography | https://doi.org/10.1111/ecog.02343 | China, United States |
| **Liu H-M et al.** | **Exploring the pteridophyte flora of the Eastern Afromontane biodiversity hotspot** | **Journal of Systematics and Evolution 54: 691–705** | **https://doi.org/10.1111/jse.12232** | **China, Kenya, United Kingdom, Germany** |
| Liu W et al. | Morphological and Genetic Variation along a North-to-South Transect in Stipa purpurea, a Dominant Grass on the Qinghai-Tibetan Plateau: Implications for Response to Climate Change | PLoS ONE 11(8): e0161972 | https://doi.org/10.1371/journal.pone.0161972 | China |
| Londei T | Piapiacs (Ptilostomus afer Linnaeus, 1766) and yellow-billed oxpeckers (Buphagus africanus Linnaeus, 1766) avoid proximity when on African buffaloes (Syncerus caffer Sparrman, 1779) | African Journal of Ecology | https://doi.org/10.1111/aje.12319 | Italy |
| López-Aguirre C et al. | Extinction of South American sparassodontans (Metatheria): environmental fluctuations or complex ecological processes? | Palaeontology | https://doi.org/10.1111/pala.12272 | Australia |
| López-Pujol J et al. | Speciation and genetic diversity in Centaurea subsect. Phalolepis in Anatolia | Scientific Reports 6 | https://doi.org/10.1038/srep37818 | Spain, Turkey, Saudi Arabia |
| **Lundgren MR & Christin P-A** | **Despite phylogenetic effects, C3–C4 lineages bridge the ecological gap to C4 photosynthesis** | **Journal of Experimental Botany** | **https://doi.org/10.1093/jxb/erw451** | **United Kingdom** |
| Lv X et al. | Continental Refugium in the Mongolian Plateau during Quaternary Glacial Oscillations: Phylogeography and Niche Modelling of the Endemic Desert Hamster, Phodopus roborovskii | PLoS ONE 11(2) | https://doi.org/10.1371/journal.pone.0148182 | China |
| Lv X et al. | Climatic niche conservatism and ecological opportunity in the explosive radiation of arvicoline rodents (Arvicolinae, Cricetidae) | Evolution | https://doi.org/10.1111/evo.12919 | China |
| Ma H et al. | Hypsodonty of Dipodidae (Rodentia) in Correlation with Diet Preferences and Habitats | Journal of Mammalian Evolution | https://doi.org/10.1007/s10914-016-9352-y | China, Israel, Belgium |
| Ma KCK et al. | Richness and zoogeography of ascidians (Tunicata: Ascidiacea) in eastern Canada | Canadian Journal of Zoology | https://doi.org/10.1139/cjz-2016-0087 | Canada |
| Macfarlane CBA et al. | Population genetic isolation and limited connectivity in the purple finch (Haemorhous purpureus) | Ecology and Evolution | https://doi.org/10.1002/ece3.2524 | Canada |
| Mahabal A et al. | Distribution records and extended range of the Sri Lanka Frogmouth Batrachostomus moniliger (Aves: Caprimulgiformes: Podargidae) in the Western Ghats: a review from 1862 to 2015 | Journal of Threattened Taxa 8 (11): 9289-9305 | https://doi.org/10.11609/jott.2775.8.11.9289-9305 | India |
| Mai P et al. | Taxonomic revision of Peperomia (Piperaceae) from Uruguay | PhytoTaxa 244(2): 125-144 | https://doi.org/10.11646/phytotaxa.244.2.2 | Uruguay, Brazil |
| Mandáková T et al. | How diploidization turned a tetraploid into a pseudotriploid | American Journal of Botany | https://doi.org/10.3732/ajb.1500452 | Czech Republic, United States |
| Marchant B et al. | Patterns of abiotic niche shifts in allopolyploids relative to their progenitors | New Phytologist | https://doi.org/10.1111/nph.14069 | United States |
| Marques I et al. | Past climate changes facilitated homoploid speciation in three mountain spiny fescues (Festuca, Poaceae) | Scientiic Reports 6 | https://doi.org/10.1038/srep36283 | Spain, Portugal, Russian Federation |
| Márquez-Borrás F et al. | First record of Ophiura ljungmani (Echinodermata: Ophiuroidea) from an anchialine cave in the Mexican Caribbean | Revista Mexicana de Biodiversidad 1127-1130 | https://doi.org/10.1016/j.rmb.2016.07.006 | Mexico, Australia |
| Marta S et al. | Deciphering range dynamics: effects of niche stability areas and post-glacial colonization on alpine species distribution | Journal of Biogeography | https://doi.org/10.1111/jbi.12771 | Italy, United States, Germany |
| Martínez-Méndez N et al. | Modelado de nicho ecológico de las especies del género Abies (Pinaceae) en México: Algunas implicaciones taxonómicas y para la conservación | Botanical Sciences 94 (1): 5-24 | https://doi.org/10.17129/botsci.508 | Mexico |
| Martínez-Vilalta J et al. | Dynamics of non-structural carbohydrates in terrestrial plants: a global synthesis | Ecological Monographs | https://doi.org/10.1002/ecm.1231 | Spain, United States, Switzerland, Germany, Chile |
| Matuszak S et al. | Key innovations and climatic niche divergence as drivers of diversification in subtropical Gentianinae in southeastern and eastern Asia | American Journal of Botany | https://doi.org/10.3732/ajb.1500352 | Germany |
| Maurin O et al. | Diversification into novel habitats in the Africa clade of Dioscorea (Dioscoreaceae): erect habit and elephant's foot tubers | BMC Evolutionary Biology 16:238 | https://doi.org/10.1186/s12862-016-0812-z | South Africa, United Kingdom, Spain, Russian Federation, France |

# advancing biodiversity science

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| Medina I & Langmore NE | The evolution of host specialisation in avian brood parasites | Ecology Letters | https://doi.org/10.1111/ele.12649 | Australia |
| **Meiri S** | **Small, rare and trendy: traits and biogeography of lizards described in the 21st century** | **Journal of Zoology** | **https://doi.org/10.1111/jzo.12356** | **Israel** |
| Mena-Portales J et al. | Taxonomy, distribution and conservation status of some interesting hyphomycetes (Anamrophic Fungi) from La Palma Biosphere Reserve, Canary Islands | Boletín de la Sociedad Micológica de Madrid 39 | http://bit.ly/2uhp0eA | Spain |
| Merow C et al. | Integrating occurrence data and expert maps for improved species range predictions | Global Ecology and Biogeography | https://doi.org/10.1111/geb.12539 | United States, United Kingdom |
| **Meyer C et al.** | **Range geometry and socio-economics dominate species-level biases in occurrence information** | **Global Ecology and Biogeography** | **https://doi.org/10.1111/geb.12483** | **Germany, United Kingdom, United States** |
| Mikula O et al. | Evolutionary history and species diversity of African pouched mice (Rodentia: Nesomyidae: Saccostomus) | Zoologica Script | https://doi.org/10.1111/zsc.12179 | Czech Republic, Kenya, Tanzania |
| Molina-Henao YF et al. | Ecological and Geographic Dimensions of Diversification in Piper subgenus Ottonia: A Lineage of Neotropical Rainforest Shrubs | Systematic Botany 41(2):253-262 | https://doi.org/10.1600/036364416X691777 | Colombia, United States |
| Morales A et al. | Environmental niche drives genetic and morphometric structure in a widespread bat. | Journal of Biogeography | https://doi.org/10.1111/jbi.12666 | Mexico, United States, Brazil |
| Müller CM et al. | Geropogon hybridus (L.) Sch.Bip. (Asteraceae) exhibits micro-geographic genetic divergence at ecological range limits along a steep precipitation gradient | Plant Systematics and Evolution | https://doi.org/10.1007/s00606-016-1354-y | Germany |
| Multimäki S et al. | Comparison of Temporally Classified and Unclassified Map Animations | Cartographic Perspectives | https://doi.org/10.14714/CP82.1296 | Finand |
| Nava-Bolaños A et al. | Isolation barriers and genetic divergence in non-territorial Argia damselflies | Biological Journal of the Linnean Society | https://doi.org/10.1111/bij.12916 | Mexico, Sweden |
| **Newbold T et al.** | **Global patterns of terrestrial assemblage turnover within and among land uses** | **Ecography** | **https://doi.org/10.1111/ecog.01932** | **United Kingdom, Norway** |
| Nicolas V et al. | Molecular data and ecological niche modelling reveal the evolutionary history of the common and Iberian blind moles (Talpidae) in Europe | Zoologica Scripta | https://doi.org/10.1111/zsc.12189 | France, Spain |
| **Nogué S et al.** | **Pollination service delivery for European crops: Challenges and opportunities** | **Ecological Economics 128: 1-7** | **https://doi.org/10.1016/j.ecolecon.2016.03.023** | **United Kingdom, Norway, Spain** |
| Nordén B & Jordal JB | Some interesting lichenized fungi from old Fraxinus excelsior and Ulmus glabra in Norway, including four new country records | Graphis Scripta 28 (1-2): 17-21 | http://bit.ly/2uh4BRZ | Norway |
| **Nylinder S et al.** | **From the Namib around the world: biogeography of the Inuleae–Plucheinae (Asteraceae)** | **Journal of Biogeography** | **https://doi.org/10.1111/jbi.12764** | **Sweden** |
| Oberlander KC et al. | Species-rich and polyploid-poor: Insights into the evolutionary role of whole-genome duplication from the Cape flora biodiversity hotspot | American Journal of Botany | https://doi.org/10.3732/ajb.1500474 | Czech Republic, South Africa, United States, Switzerland |
| **Oliveira U et al.** | **The strong influence of collection bias on biodiversity knowledge shortfalls of Brazilian terrestrial biodiversity** | **Diversity and Distributions** | **https://doi.org/10.1111/ddi.12489** | **Brazil, Singapore** |
| Onstein RE & Linder HP | Beyond climate: convergence in fast evolving sclerophylls in Cape and Australian Rhamnaceae predates the mediterranean climate | Journal of Ecology | https://doi.org/10.1111/1365-2745.12538 | France, Switzerland |
| Onstein RE et al. | Evolutionary radiations of Proteaceae are triggered by the interaction between traits and climates in open habitats | Global Ecology and Biogeography | https://doi.org/10.1111/geb.12481 | Switzerland, France, Australia |
| Ornelas JF et al. | A mistletoe tale: postglacial invasion of Psittacanthus schiedeanus (Loranthaceae) to Mesoamerican cloud forests revealed by molecular data and species distribution modeling | BMC Evolutionary Biology | https://doi.org/10.1186/s12862-016-0648-6 | Mexico, United States |
| Ornelas JF et al. | Molecular and iridescent feather reflectance data reveal recent genetic diversification and phenotypic differentiation in a cloud forest hummingbird | Ecology and Evolution | https://doi.org/10.1002/ece3.1950 | Mexico, Netherlands |
| Pérez-De la O B et al. | Model Simulation of Potential Distribution of Lobesia botrana (Denis & Schiffermüller)1 in Grape, Vitis vinifera (L.) Vineyard Regions of Mexico | Southwestern Entomologist 41(3):693-704 | https://doi.org/10.3958/059.041.0312 | Mexico, United States |
| **Pérez-Moreno H et al.** | **Climatic patterns in the establishment of wintering areas by North American migratory birds** | **Ecology and Evolution** | **https://doi.org/10.1002/ece3.1973** | **Mexico, United States** |
| Perktaş U et al. | Integrating morphology, phylogeography, and ecological niche modeling to explore population differentiation in North African Common Chaffinches | Journal of Ornithology | https://doi.org/10.1007/s10336-016-1361-3 | Turkey, United States |
| Peterson AT & Anamza T | Reexamining Phylloscopus trochiloides complex as a ring species: A refugial counter-hypothesis | Journal of Avian Biology | https://doi.org/10.1111/jav.01167 | United States |

## advancing biodiversity science

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| Peterson AT et al. | Assumption- versus data-based approaches to summarizing species' ranges | Conservation Biology | https://doi.org/10.1111/cobi.12801 | United States, Mexico |
| **Peterson AT et al.** | **Digital Accessible Knowledge and well-inventoried sites for birds in Mexico: baseline sites for measuring faunistic change** | **PeerJ 4** | **https://doi.org/10.7717/peerj.2362** | **United States, Mexico** |
| **Pierce S et al.** | **A global method for calculating plant CSR ecological strategies applied across biomes worldwide** | **Functional Ecology** | **https://doi.org/10.1111/1365-2435.12722** | **Italy, Brazil, Germany, Argentina, Canada, Panama, Netherlands, United States, Russian Federation, United Kingdom, Tanzania, Australia, China, France, Belgium, Spain** |
| Pires THS et al. | Traits of a lineage with extraordinary geographical range: ecology, behavior and life-history of the sailfin tetra Crenuchus spilurus | Environmental Biology of Fishes | https://doi.org/10.1007/s10641-016-0534-5 | Brazil |
| Pirie MD et al. | The biodiversity hotspot as evolutionary hot-bed: spectacular radiation of Erica in the Cape Floristic Region | BMC Evolutionary Biology 16: 190 | https://doi.org/10.1186/s12862-016-0764-3 | Germany, South Africa |
| Posso-Terranova et al. | Complex niche divergence underlies lineage diversification in Oophaga poison frogs | Journal of Biogeography | https://doi.org/10.1111/jbi.12799 | Canada, Colombia |
| Poveda-Martínez D et al. | Historical Biogeography of Five Characidium Fish Species: Dispersal from the Amazon Paleobasin to Southeastern South America | PLoS ONE 11(10): e0164902 | https://doi.org/10.1371/journal.pone.0164902 | Colombia |
| Prieto-Torres DA & Rojas-Soto OR | Reconstructing the Mexican Tropical Dry Forests via an Autoecological Niche Approach: Reconsidering the Ecosystem Boundaries | PLoS ONE 11(3): e015093 | https://doi.org/10.1371/journal.pone.0150932 | Mexico |
| **Pyle RL et al.** | **A comprehensive investigation of mesophotic coral ecosystems in the Hawaiian Archipelago** | **PeerJ 4:e2475** | **https://doi.org/10.7717/peerj.2475** | **United States** |
| Quintero-Díaz GE et al. | The Great Plains Ratsnake, Pantherophis emoryi Baird & Girard, 1853, (Squamata: Colubridae), a new state record from Aguascalientes, México | Checklist 12(3) | https://doi.org/10.15560/12.3.1903 | Mexico |
| **Rabosky ARD et al.** | **Coral snakes predict the evolution of mimicry across New World snakes** | **Nature Communications** | **https://doi.org/10.1038/ncomms11484** | **United States, Israel** |
| Ranc N et al. | Performance tradeoffs in target-group bias correction for species distribution models | Ecography | https://doi.org/10.1111/ecog.02414 | United States, Italy, France, Sweden |
| Ratnam J et al. | Savannahs of Asia: antiquity, biogeography, and an uncertain future | Phylosophical Transactions of the Royal Society B 371 (1703) | https://doi.org/10.1098/rstb.2015.0305 | India, China, United Kingdom |
| Ray D et al. | Improving spatial transferability of ecological niche model of Hevea brasiliensis using pooled occurrences of introduced ranges in two biogeographic regions of India | Ecological Informatics 34:153-163 | https://doi.org/10.1016/j.ecoinf.2016.06.003 | India |
| Razanajatovo M et al. | Plants capable of selfing are more likely to become naturalized | Nature Communications 7 | https://doi.org/10.1038/ncomms13313 | Germany, United Kingdom, Austria, South Africa, Czech Republic |
| Reginato M & Michelangeli FA | Diversity and constraints in the floral morphological evolution of Leandra s.str. (Melastomataceae) | Annals of Botany | https://doi.org/10.1093/aob/mcw116 | United States |
| Ribeiro V et al. | Ecological and historical views of the diversification of Geositta miners (Aves: Furnariidae: Sclerurinae) | Journal of Ornithology | https://doi.org/10.1007/s10336-016-1398-3 | Brazil, United States |
| Ribeiro V et al. | Distribution dynamics of South American savanna birds in response to Quaternary climate change | Austral Ecology | https://doi.org/10.1111/aec.12363 | Brazil |
| Ribeiro-Júnior MA & da Silva MB et al. | A New Species of Bachia Gray 1845 (Squamata: Gymnophthalmidae) from the Eastern Guiana Shield | Herpetologica 72 (2):148-156 | https://doi.org/10.1655/HERPETOLOGICA-D-15-00030 | Brazil |
| Rivkin LR et al. | Why is gynodioecy a rare but widely distributed sexual system? Lessons from the Lamiaceae | New Phytologist | https://doi.org/10.1111/nph.13926 | Canada, United States |
| Rodrigues A & Stefanović S | Present-Day Genetic Structure of the Holoparasite Conopholis americana (Orobanchaceae) in Eastern North America and the Location of Its Refugia during the Last Glacial Cycle | International Journal of Plant Sciences | https://doi.org/10.1086/684180 | Canada |
| Rodríguez-San Pedro A et al. | Update on the distribution of Myotis atacamensis (Chiroptera: Vespertilionidae): first record from central Chile and description of echolocation calls | Chiroptera Neotropical 21(2): 1342-1346 | http://bit.ly/2dkiwRl | Chile |
| **Roger N et al.** | **Impact of pollen resources drift on common bumble bees in NW Europe** | **Global Change Biology** | **https://doi.org/10.1111/gcb.13373** | **Belgium, Brazil, Portugal, Netherlands** |
| Saeedi H et al. | Bimodal latitudinal species richness and high endemicity of razor clams (Mollusca) | Journal of Biogeography | https://doi.org/10.1111/jbi.12903 | New Zealand |

## advancing biodiversity science

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| **Salas-Lopez A et al.** | **The discovery of devil's gardens: an ant–plant mutualism in the cloud forests of the Eastern Amazon** | **Journal of Tropical Ecology 32 (3): 264-268** | **https://doi.org/10.1017/S0266467416000195** | **France** |
| Sanchez A | Fidelity and Promiscuity in an Ant-Plant Mutualism: A Case Study of Triplaris and Pseudomyrmex | PLoS ONE 10(12): e0143535 | https://doi.org/10.1371/journal.pone.0143535 | Colombia |
| Sanín MJ et al. | The Neogene rise of the tropical Andes facilitated diversification of wax palms (Ceroxylon: Arecaceae) through geographical colonization and climatic niche separation | Botanical Journal of the Linnean Society | https://doi.org/10.1111/boj.12419 | Colombia, Netherlands, Sweden, Denmark, Peru, France |
| Santana-Moreno LD et al. | First record of Ambidexter symmetricus (Decapoda: Caridea: Processidae) from the Yucatán Peninsula | Revista Mexicana de Biodiversidad | https://doi.org/10.1016/j.rmb.2016.07.008 | Mexico |
| Santos Ferreira da Cruz RH et al. | Distribution Data of Bird's Nest Fungi in Japan: Nidula niveotomentosa and Crucibulum laeve | Bulletin of the National Museum of Nature and Science. Series B, Botany 42 (2): 49-55 | http://bit.ly/2uh0g0J | Brazil, Japan |
| Sarkar SK et al. | On the Mimela Kirby, 1823 (Rutelinae: Scarabaeidae) of Buxa Tiger Reserve (a forest under biodiversity hot spot zone), Dooars, West Bengal, India | World Scientific News 50: 95-105 | http://bit.ly/2tWWxX1 | India |
| Schiffer PH & Herbig H-G | Endorsing Darwin: global biogeography of the epipelagic goose barnacles Lepas spp. (Cirripedia, Lepadomorpha) proves cryptic speciation | Zoological Journal of the Linnean Society | https://doi.org/10.1111/zoj.12373 | Germany, United Kingdom, Italy |
| Schneider JV et al. | Water supply and demand remain coordinated during breakdown of the global scaling relationship between leaf size and major vein density | New Phytologist | https://doi.org/10.1111/nph.14382 | Germany |
| Scoffoni C et al. | Hydraulic basis for the evolution of photosynthetic productivity | Nature Plants 2 | https://doi.org/10.1038/nplants.2016.72 | United States |
| Sebasky ME et al. | Investigating past range dynamics for a weed of cultivation, Silene vulgaris | Ecology and Evolution | https://doi.org/10.1002/ece3.2250 | United States |
| Shabani F et al. | A comparison of absolute performance of different correlative and mechanistic species distribution models in an independent area | Ecology and Evolution | https://doi.org/10.1002/ece3.2332 | Australia, Iran |
| Shabani F et al. | Variances in the projections, resulting from CLIMEX, Boosted Regression Trees and Random Forests techniques | Theoretical and Applied Climatology | https://doi.org/10.1007/s00704-016-1812-z | Australia, Iran |
| Shabani F et al. | A comparison of absolute performance of different correlative and mechanistic species distribution models in an independent area | Ecology and Evolution | https://doi.org/10.1002/ece3.2332 | Australia, Iran |
| Shang H et al. | Likely allopatric origins of Adiantum × meishanianum (Pteridaceae) through multiple hybridizations | Journal of Systematics and Evolution | https://doi.org/10.1111/jse.12205 | China |
| Shelton K et al. | New Records of Entocytherid Ostracods from New York and New Jersey | Northeastern Naturalist 23(4):525-531 | https://doi.org/10.1656/045.023.0401 | United States |
| Shrestha N & Zhang X-C | Is Huperzia hamiltonii (Spreng.) Trevis. A Himalayan Endemic? An Empirical Evaluation using Species Distribution Modeling | Indian Fern J. 31: 154-161 | http://bit.ly/2tjxW0N | China |
| **Sikes DS et al.** | **Cryptic diversity in the New World burying beetle fauna: Nicrophorus hebes Kirby; new status as a resurrected name (Coleoptera: Silphidae: Nicrophorinae)** | **Arthropod Systematics & Phylogeny 74(3): 299-309** | **http://bit.ly/2rytn1P** | **United States, Canada** |
| Silva DP et al. | Assessing the distribution and conservation status of a long-horned beetle with species distribution models | Journal of Insect Conservation | https://doi.org/10.1007/s10841-016-9892-8 | Brazil |
| **Smith BE et al.** | **From GenBank to GBIF: Phylogeny-Based Predictive Niche Modeling Tests Accuracy of Taxonomic Identifications in Large Occurrence Data Repositories** | **PLoS ONE 11(3): e0151232** | **https://doi.org/10.1371/journal.pone.0151232** | **United States, Germany** |
| Smith DM & Finch DM | Riparian trees and aridland streams of the southwestern United States: An assessment of the past, present, and future | Journal of Arid Environments 135:120-131 | https://doi.org/10.1016/j.jaridenv.2016.08.016 | United States |
| Sniderman JMK et al. | Pliocene reversal of late Neogene aridification | PNAS, Proceedings of the National Academy of Sciences | https://doi.org/10.1073/pnas.1520188113 | Australia, France |
| Sosa V et al. | Historical reconstruction of climatic and elevation preferences and the evolution of cloud forest-adapted tree ferns in Mesoamerica | PeerJ 4 | https://doi.org/10.7717/peerj.2696 | Mexico |
| Sosa-Pivatto M et al. | Do 120,000 years of plant–pollinator interactions predict floral phenotype divergence in Calceolaria polyrhiza? A reconstruction using species distribution models | Arthropod-Plant Interactions | https://doi.org/10.1007/s11829-016-9490-4 | Argentina, United States |
| Sotka EE et al. | A molecular phylogeny of marine amphipods in the herbivorous family Ampithoidae | Zoologica Scripta | https://doi.org/10.1111/zsc.12190 | United States, Australia |
| **Spalink D et al.** | **Biogeography of the cosmopolitan sedges (Cyperaceae) and the area-richness correlation in plants** | **Journal of Biogeography** | **https://doi.org/10.1111/jbi.12802** | **United States, Canada** |

## advancing biodiversity science

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| **Stropp J et al.** | **Mapping ignorance: 300 years of collecting flowering plants in Africa** | **Global Ecology and Biogeography** | **https://doi.org/10.1111/geb.12468** | **Italy, Brazil, United Kingdom, Spain, Luxembourg, Belgium** |
| Sungani H et al. | Multiple colonisations of the Lake Malawi catchment by the genus Opsaridium (Teleostei: Cyprinidae) | Molecular Phylogenetics and Evolution | https://doi.org/10.1016/j.ympev.2016.09.027 | United Kingdom, Malawi, Tanzania, Austria, South Africa |
| Tassia MG et al. | The Global Diversity of Hemichordata. | PLoS ONE 11(10): e0162564 | https://doi.org/10.1371/journal.pone.0162564 | United States, Sweden, Israel |
| **ter Steege H et al.** | **The discovery of the Amazonian tree flora with an updated checklist of all known tree taxa** | **Scientific Reports 6: 29549** | **https://doi.org/10.1038/srep29549** | **Netherlands, Colombia, France, Sweden, United States, Brazil** |
| Thiede J | A Review of Agave mitis (Asparagaceae/Agavaceae) | Bradleya 34 | http://bit.ly/2ugXWr9 | Germany |
| Thomas E et al. | Genetic diversity of Enterolobium cyclocarpum in Colombian seasonally dry tropical forest: implications for conservation and restoration | Biodiversity and Conservation | https://doi.org/10.1007/s10531-016-1274-8 | Peru, Colombia, Italy |
| Tirira DG et al. | Genus Glyphonycteris Thomas, 1896 (Mammalia: Chiroptera) in Ecuador: first confirmed record of G. sylvestris Thomas, 1896 and a geographical review to G. daviesi (Hill, 1965) | Check List 12(5) | https://doi.org/10.15560/12.5.1965 | Ecuador |
| Tran LAP | Interaction between Digestive Strategy and Niche Specialization Predicts Speciation Rates across Herbivorous Mammals | The American Naturalist | https://doi.org/10.1086/685094 | United States |
| Triponez Y et al. | Genome fingerprinting confirms the species status of the Loosestrifes Lysimachia punctata and L. verticillata (Primulaceae) | Plant Ecology and Evolution 149 (3): 335–338 | https://doi.org/10.5091/plecevo.2016.1126 | Switzerland, Turkey |
| Turchetto-Zolet AC et al. | Phylogeography and ecological niche modelling in Eugenia uniflora (Myrtaceae) suggest distinct vegetational responses to climate change between the southern and the northern Atlantic Forest | Botanical Journal of the Linnean Society | https://doi.org/10.1111/boj.12473 | Brazil |
| Ullah Z et al. | Phytogeographic analysis and Diversity of grasses and sedges (poales) of Northern Pakistan | Pakistan Journal of Botany 47(SI): 93-104 | http://bit.ly/2ugZXmY | Pakistan |
| Varela-Hernández F et al. | New Records of Ants from Oaxaca, Southern Mexico | Southwestern Entomologist 41(3):705-714 | https://doi.org/10.3958/059.041.0313 | Mexico |
| Vasconcelos TS & Doro JLP | Assessing how habitat loss restricts the geographic range of Neotropical anurans | Ecological Research | https://doi.org/10.1007/s11284-016-1401-8 | Brazil |
| Vitorino LC et al. | Demographical history and palaeodistribution modelling show range shift towards Amazon Basin for a Neotropical tree species in the LGM | BMC Evolutionary Biology 16: 213 | https://doi.org/10.1186/s12862-016-0779-9 | Brazil |
| Wan J-Z et al. | Climate change may alter genetic diversity of Duchesnea indica, a clonal plant species | Biochemical Systematics and Ecology 66: 114-122 | https://doi.org/10.1016/j.bse.2016.03.002 | China |
| Wang HB et al. | Lignified woods of Pinus (Pinaceae) from the late Miocene of central Yunnan, China, and their biogeographic and paleoclimatic implications | Palaeoworld | https://doi.org/10.1016/j.palwor.2016.06.003 | China, South Africa, Russian Federation |
| **Wang J-C et al.** | **Evolution of morphological and climatic adaptations in Veronica L. (Plantaginaceae)** | **PeerJ 4:e2333** | **https://doi.org/10.7717/peerj.2333** | **China, Germany** |
| Wang S-H et al. | Contrasting genetic patterns between two coexisting Eleutherococcus species in northern China | Ecology and Evolution | https://doi.org/10.1002/ece3.2118 | China |
| Wepfer PH et al. | Influences of climate and historical land connectivity on ant beta diversity in East Asia | Journal of Biogeography | https://doi.org/10.1111/jbi.12762 | Japan, China |
| Wicaksono CY et al. | Contracting montane cloud forests: a case study of the Andean alder (Alnus acuminata) and associated fungi in the Yungas | Biotropica | https://doi.org/10.1111/btp.12394 | Netherlands, Argentina |
| Willis CG & Donohue K | The evolution of intrinsic reproductive isolation in the genus Cakile (Brassicaceae) | Journal of Evolutionary Biology | https://doi.org/10.1111/jeb.13011 | United States |
| Wilson CA & Calvin CL | Metadata provide insights on patterns of epiparasitism in mistletoes (Santalales), an overlooked topic in forest biology | Botany | https://doi.org/10.1139/cjb-2016-0264 | United States |
| Winterstein M et al. | A Range Extension for Carex sartwellii in Interior Alaska | The Canadian Field-Naturalist 130(3): 191-198 | https://doi.org/10.22621/cfn.v130i3.1878 | United States |
| **Wyse SV & Dickie JB** | **Predicting the global incidence of seed desiccation sensitivity** | **Journal of Ecology** | **https://doi.org/10.1111/1365-2745.12725** | **United Kingdom** |
| Yang J et al. | Phylogeography and evolution of two closely related oak species (Quercus) from north and northeast China | Tree Genetics & Genomes 12:89 | https://doi.org/10.1007/s11295-016-1044-5 | China |

## advancing biodiversity science

| AUTHORS | TITLE | JOURNAL | DOI/URL | AUTHOR COUNTRIES |
|---|---|---|---|---|
| Yang L-Q et al. | Molecular phylogeny, biogeography and ecological niche modelling of Cardiocrinum (Liliaceae): insights into the evolutionary history of endemic genera distributed across the Sino-Japanese floristic region | Annals of Botany | https://doi.org/10.1093/aob/mcw210 | China |
| Zhang D et al. | Pleistocene sea level fluctuation and host plant habitat requirement influenced the historical phylogeography of the invasive species Amphiareus obscuriceps (Hemiptera: Anthocoridae) in its native range | BMC Evolutionary Biology 16: 174 | https://doi.org/10.1186/s12862-016-0748-3 | China, Japan |
| Zhang L et al. | Using DEM to predict Abies faxoniana and Quercus aquifolioides distributions in the upstream catchment basin of the Min River in southwest China | Ecological Indicators 69: 91-99 | https://doi.org/10.1016/j.ecolind.2016.04.008 | China, Canada |
| Zhao J-L et al. | Ecological divergence of two closely related Roscoea species associated with late Quaternary climate change | Journal of Biogeography | https://doi.org/10.1111/jbi.12809 | China, United States |
| Zhao M et al. | Phylogenetic relationships, song and distribution of the endangered Rufous-headed Robin Larvivora ruficeps | IBIS | https://doi.org/10.1111/ibi.12426 | China, Sweden |
| **Zizka A et al.** | **Finding needles in the haystack: Where to look for rare species in the American tropics** | **Ecography** | **https://doi.org/10.1111/ecog.02192** | **Sweden, Netherlands, Brazil** |

## What's next

Download the GBIF *Science Review* 2017
gbif.org/science-review-2017

Read the latest data use summaries
gbif.org/newsroom/uses/summary

Explore the data available at GBIF.org
gbif.org/occurrence/search

Subscribe to news and updates from GBIF
gbif.org/newsroom/summary#sign-up



**GBIF Secretariat**
Universitetsparken 15
DK-2100 Copenhagen Ø
Denmark

*tel* +45 35 32 14 70
*email* communication@gbif.org
*web* www.gbif.org