


WILLOWOOD CLOMAZONE 3ME

Same active as *COMMAND® 3ME*



OVERVIEW

Willowood Clomazone 3ME is key herbicide used to control a wide range of broadleaf weeds and grasses in several crops.

KEY CROPS

Rice • Soybeans • Cotton • Sugarcane • Tobacco

PACKAGING

- 2.5 gal (x2)

PRODUCT SPECIFICATIONS

ACTIVE INGREDIENT

Clomazone (3.0lbs Active Ingredient per Gallon)

RATES

11.0 fl oz/A - 34.0 fl oz/A
(See label for rates by crop)

WEED SPECTRUM

Barnyard Grass • Signal Grass • Panicum Species

APPLICATION TIMING

Pre Emergence • Early Post Emergence

Technical Use Guidelines

| | |
|---|---|
| Mode of Action | <ul style="list-style-type: none"> • DOXP SYNTHASE INHIBITORS |
| Crop Rotation REFER TO LABEL FOR DETAILS | <ul style="list-style-type: none"> • Corn - 9 months • Rice: Vary by rate applied • Cotton: Vary by rate applied |
| Adjuvant Recommendation | <ul style="list-style-type: none"> • NIS: 2 pints per 100 gallons of spray mixture (post emergence) |
| Rainfastness | <ul style="list-style-type: none"> • Not Applicable |
| Water Volume (GPA) REFER TO LABEL FOR DETAILS | <ul style="list-style-type: none"> • Ground: 10GPA - 25GPA |
| Application Tips | <ul style="list-style-type: none"> • Use as a broadleaf herbicide as needed. • Avoid gusty or windless conditions. |

ALWAYS READ AND FOLLOW LABEL DIRECTIONS. It is a violation of federal and state/provincial law to use any pesticide product other than in accordance with its label. The distribution, sale and use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. We do not guarantee the accuracy of any information provided on this page or which is provided by us in any form. It is your responsibility to confirm prior to purchase and use that a product is labeled for your specific purposes, including, but not limited to, your target crop or pest and its compatibility with other products in a tank mix. This information should not be used as a replacement for consulting the applicable product label. Consult the label for the most complete and up-to-date information about any referenced product. Readers must have a valid applicator license to use restricted use pesticides. Please consult your state department of agriculture for complete rules and regulations on the use of restricted use pesticides, as some products require specific record-keeping requirements.



WILLOWOOD CLOMAZONE 3ME

Same active as *COMMAND® 3ME*

MIXING PROCEDURE

Mix Willowood Clomazone 3ME as follows:

- Fill spray tank 1/4 to 1/3 full with water and maintain agitation.
- Add the specified amount of ingredients using the following order:
 - a. Dry formulations (e.g., wettable powders, dry flowables)
 - b. Liquid suspensions (e.g., flowables)
 - c. Liquids (e.g., EC's)
- Mix thoroughly and fill tank 1/2 full and maintain agitation.
- Add Willowood Clomazone 3ME to tank and maintain agitation.
- Complete filling the sprayer tank with water.
- If a surfactant is added, add it as the last ingredient to the spray tank.
- Maintain agitation during filling, mixing and application.
- If using a drift reduction agent, follow specific product label instructions for order of addition to spray tank.

CROP STAGING

Cotton: apply as a pre-emergent banded or broadcast treatment in conventional tillage or conservation tillage systems to control annual grasses and broadleaf weeds. Do not incorporate or use in a tank mixture with any herbicide that requires soil incorporation.

Rice

- Pre Emergence Surface Broadcast: apply 14 days before planting or up to 7 days after planting. Apply prior to weed emergence.
- Early Post Emergence Application: apply after planting as an early post emergence treatment at the 1- to 5-leaf stage.

Soybeans: apply as a pre-emergence soil applied treatment to control annual grasses and broadleaf weeds. Tank mix with other pre-emergence soybean herbicides for greater weed control (see label).

Sugarcane: apply to newly planted sugarcane in late summer or early fall. Apply to newly planted or ratoon sugarcane in spring through layby.

Tobacco

- Pre Emergence (Prior to Transplant): apply immediately or up to 30 days before transplanting. Apply as a surface treatment prior to weed emergence. If weeds emerge prior to pre emergence application, cultivate before or immediately preceding treatment.
- Pre Emergence (Post-Transplant): make an over-the-top treatments to plants immediately, up to 7 days after, or at transplanting, but ensure that application is made prior to emergence of weeds. If weeds emerge prior to pre emergence application, cultivate before or immediately preceding treatment.

FIELD NOTES
