

WILLOWOOD THIOBEN 8EC

Same active as BOLERO® 8EC



OVERVIEW

Willowood Thioben 8EC Rice Herbicide provides excellent contact and residual control of key weeds including, watergrass, smallflower umbrellaplant and sprangletop. Used early post emergence in dry-seeded rice or preplant incorporated in water seeded rice, Willowood Thioben 8EC affords growers tank mix flexibility which can broaden the weed control spectrum and provide additional residual control.

KEY CROPS

RICE

AVAILABLE PACK SIZE

- 2.5 gal jug (x2)

PRODUCT SPECIFICATIONS

ACTIVE INGREDIENT

Thiobencarb (8.0 lbs Active Ingredient per Gallon)

RATES

2.5 pt/A - 4.0 pt/A
(See label for rates by crop)

WEED SPECTRUM

Red Rice • Sprangletop • Watergrass • Smallflower Umbrellaplant

APPLICATION TIMING

Preemergence • Post Emergence

Technical Use Guidelines

Mode of Action	<ul style="list-style-type: none"> • LIPID SYNTHESIS INHIBITOR (NOT ACCase)
Crop Rotation REFER TO LABEL FOR DETAILS	<ul style="list-style-type: none"> • 6 months on all rotational crops
Adjuvant Recommendation	<ul style="list-style-type: none"> • Do not mix with liquid nitrogen or surfactant when applied alone or when tank mixed with Propanil
Rainfastness	<ul style="list-style-type: none"> • 24 hours
Water Volume (GPA) REFER TO LABEL FOR DETAILS	<ul style="list-style-type: none"> • Ground: 10 GPA - 20 GPA • Aerial: 10 GPA
Application Tips	<ul style="list-style-type: none"> • Provides residual control of some listed weeds on label for up to 5 weeks • Flexible with several application timing options

ALWAYS READ AND FOLLOW LABEL DIRECTIONS. It is a violation of federal and state/provincial law to use any pesticide product other than in accordance with its label. The distribution, sale and use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. We do not guarantee the accuracy of any information provided on this page or which is provided by us in any form. It is your responsibility to confirm prior to purchase and use that a product is labeled for your specific purposes, including, but not limited to, your target crop or pest and its compatibility with other products in a tank mix. This information should not be used as a replacement for consulting the applicable product label. Consult the label for the most complete and up-to-date information about any referenced product. Readers must have a valid applicator license to use restricted use pesticides. Please consult your state department of agriculture for complete rules and regulations on the use of restricted use pesticides, as some products require specific record-keeping requirements.

WILLOWOOD THIOBEN 8EC

Same active as *BOLERO® 8EC*

MIXING PROCEDURE

Mix Willowood Thioben 8EC as follows:

- Fill the tank one-half full of clean water.
- Begin agitation.
- If foaming is anticipated, add defoamer prior to the addition of the surfactant. Add the required amount of Willowood Thioben 8EC.
- Add tank mix partner (if any) in the following order:
 - Water soluble packets (preferably added before the surfactant)
 - Water dispersible granules/wettable powder
 - Soluble powders/UAN
 - Suspension concentrate
 - Emulsifiable concentrate
- Fill the remainder of the tank.
- Mix only the amount of spray solution that can be applied the day of mixing.
- Willowood Thioben 8EC must be applied within 12 hours of mixing.

CROP STAGING

- **Dry-Seeded Rice**
 - Early Post Emergence Application Non-Flooded
 - Apply when watergrass has developed no further than the 3 leaf stage.
 - When making application to drill-seeded rice, application may be made any time after germination (usually 5 to 9 days after seeding).
- **Water-Seeded Rice**
 - Pre Plant Non-Incorporated Non-Flooded
 - For control of watergrass at the 3 leaf stage or less, sprangletop 2-1/2 leaf or 3/4 to 1 inch in height, whichever is smaller, and smallflower umbrellaplant up to 1/2 inch in height.

FIELD NOTES
