

# WILLOWOOD ETHOFUMESATE 4SC

Same active as NORTRON®



## OVERVIEW

Willowood Ethofumesate 4SC provides an excellent weed control foundation with multiple application options. In addition, Ethofumesate 4SC also controls a wide variety of annual grasses and broadleaf weeds. When used in conjunction with RoundUp Ready® sugarbeets, Ethofumesate 4SC provides a solid foundation for the development of a weed resistance management program.

## KEY CROPS

Sugarbeets • Grass Seed • Vegetables

## PACKAGING

- 110 gal tote (x1)
- 2.5 gal jug (x2)

## PRODUCT SPECIFICATIONS

### ACTIVE INGREDIENT

Ethofumesate (4.0 lbs Active Ingredient per Gallon)

### RATES

0.5 pt/A - 8 pt/A  
(See label for rates by crop)

### WEED SPECTRUM

Kochia • Lambsquarters • Pigweed

### APPLICATION TIMING

Preemergence • Post Emergence

## Technical Use Guidelines

<b>Mode of Action</b>	<ul style="list-style-type: none"> <li>• Seedling Shoot Growth Inhibitors</li> </ul>
<b>Crop Rotation</b> REFER TO LABEL FOR DETAILS	<ul style="list-style-type: none"> <li>• All Crops: 6-12 months except sugarbeets and ryegrass</li> </ul>
<b>Adjuvant Recommendation</b>	<ul style="list-style-type: none"> <li>• AMS</li> <li>• NIS</li> <li>• MSO</li> </ul>
<b>Rainfastness</b>	<ul style="list-style-type: none"> <li>• 6 hours</li> </ul>
<b>Water Volume (GPA)</b> REFER TO LABEL FOR DETAILS	<ul style="list-style-type: none"> <li>• Ground: 10 GPA - 60 GPA</li> </ul>
<b>Application Tips</b>	<ul style="list-style-type: none"> <li>• Do not rotate with any crops other than sugarbeets or ryegrass for:               <ul style="list-style-type: none"> <li>◦ 12 months following preplant incorporated, pre emergence, conventional post emergence applications, or split (low rate) applications totaling more than 12 fl oz (0.375 lb ai/A);</li> </ul> </li> <li>• Do not use a Crop Oil Concentrate.</li> </ul>

ALWAYS READ AND FOLLOW LABEL DIRECTIONS. It is a violation of federal and state/provincial law to use any pesticide product other than in accordance with its label. The distribution, sale and use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. We do not guarantee the accuracy of any information provided on this page or which is provided by us in any form. It is your responsibility to confirm prior to purchase and use that a product is labeled for your specific purposes, including, but not limited to, your target crop or pest and its compatibility with other products in a tank mix. This information should not be used as a replacement for consulting the applicable product label. Consult the label for the most complete and up-to-date information about any referenced product. Readers must have a valid applicator license to use restricted use pesticides. Please consult your state department of agriculture for complete rules and regulations on the use of restricted use pesticides, as some products require specific record-keeping requirements.

# WILLOWOOD ETHOFUMESATE 4SC

Same active as *NORTRON*®

## MIXING PROCEDURE

Mix Willowood Ethofumesate 4SC as follows:

- Fill the spray tank to 1/4 to 1/3 of the total spray volume required with water or liquid fertilizer.
- Start agitation.
- Add the required amount of product for the acreage being treated directly to the spray tank while agitating and allow time to disperse. If liquid fertilizer is being used as the spray carrier rather than water, pre-mix product as described on label before adding to the spray tank.
- After adding product, add different formulation types in the following order: other formulation(s) packaged in water soluble packets; any compatibility agent, if required; dry flowables; wettable powders; Aqueous suspension, flowable and liquids.
- Maintain agitation and fill spray tank to 3/4 of total spray volume and add:
  - Emulsifiable concentrates; Solutions; Adjuvants. Allow time for complete mixing and dispersion after each addition.
- Finish filling the spray tank. Maintain continuous agitation during mixing and throughout application.

## CROP STAGING

### Sugar Beets (Pre Emergence):

- Grown Under Rainfall: apply alone or in a tank mix pre emergence at time of planting or shortly after, but prior to weed germination.
- Grown Under Furrow Irrigation: apply along or in a tank mix to the soil surface preplant or at time of planting, but prior to weed germination, and incorporate into the soil.
- Under Sprinkler Irrigation: apply alone or in a tank mix pre emergence at time of planting or shortly after, and irrigate prior to crop and weed germination.

### Sugar Beets (Post Emergence):

- Apply Willowood Ethofumesate 4SC + glyphosate to sugar beets having greater than 2 true leaves.

**Grass Seed:** See label for application timing instructions.

**Vegetables:** See label for specific application timing instructions.

## FIELD NOTES

---



---



---



---



---



---



---