

GROUP

2

HERBICIDE

WILLOWOOD CHLORIM 25 WDG

Same active as CLASSIC®



OVERVIEW

Willowood Chlorim 25WDG is a pre emergence and post emergence soybean herbicide used alone or in combination with other compatible tank-mix partners for the control of a number of key weeds including but not limited to: velvetleaf, morning glory, hemp sesbania and yellow nutsedge.

KEY CROPS

Soybeans • Peanuts • Non-Crop Areas

PACKAGING

- 8 x 10 oz. cases

PRODUCT SPECIFICATIONS

ACTIVE INGREDIENT

Chlorimuron ethyl (25%)

RATES

0.33 fl oz/A - 0.75 fl oz/A
(See label for rates by crop)

WEED SPECTRUM

Ragweed • Pigweed • Marestail • Dandelion

APPLICATION TIMING

Preplant (including burndown) • Pre Emergence • Post Emergence

Technical Use Guidelines

Mode of Action	<ul style="list-style-type: none"> • ALS INHIBITORS
Crop Rotation REFER TO LABEL FOR DETAILS	<ul style="list-style-type: none"> • Soybean - no rotation restrictions • Corn - 9 months • Grain Sorghum - 9 -15 months (depends on REI)
Adjuvant Recommendation	<ul style="list-style-type: none"> • COC: 1%V/V • NIS: .25% V/V • UAN: 32 oz/A • AMS: 2 lb/A
Rainfastness	<ul style="list-style-type: none"> • 1 hour
Water Volume (GPA) REFER TO LABEL FOR DETAILS	<ul style="list-style-type: none"> • Ground: 10 GPA - 25 GPA • Aerial: 3 GPA - 5 GPA
Application Tips	<ul style="list-style-type: none"> • Broadcast Application: <ul style="list-style-type: none"> ◦ Apply by ground equipment. ◦ Use flat fan nozzles at 25 to 40 PSI. ◦ Use 10 to 25 gal of spray volume per acre.

ALWAYS READ AND FOLLOW LABEL DIRECTIONS. It is a violation of federal and state/provincial law to use any pesticide product other than in accordance with its label. The distribution, sale and use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. We do not guarantee the accuracy of any information provided on this page or which is provided by us in any form. It is your responsibility to confirm prior to purchase and use that a product is labeled for your specific purposes, including, but not limited to, your target crop or pest and its compatibility with other products in a tank mix. This information should not be used as a replacement for consulting the applicable product label. Consult the label for the most complete and up-to-date information about any referenced product. Readers must have a valid applicator license to use restricted use pesticides. Please consult your state department of agriculture for complete rules and regulations on the use of restricted use pesticides, as some products require specific record-keeping requirements.



WILLOWOOD CHLORIM 25 WDG

Same active as CLASSIC®

MIXING PROCEDURE

Mix Willowood Chlorim 25 WDG as follows:

- Follow these steps for spray tank preparation:
 1. Fill the spray tank $\frac{1}{4}$ to $\frac{1}{3}$ full of water.
 2. While agitating, add the required amount of Willowood Chlorim 25WDG.
 3. Continue adequate agitation.
 4. Willowood Chlorim 25WDG must be thoroughly mixed with water in the spray tank before adding any other material (in order: tank mix herbicide, surfactant, crop oil concentrate, or nitrogen-based fertilizer). Agitation is required for uniform mixing and application.
 5. Use the Willowood Chlorim 25WDG spray preparation within 24 hours of product mixing, or product degradation may occur.
 6. If the mixture has settled, thoroughly re-agitate before using

CROP STAGING

- **Timing to Crop Stage:** application of Willowood Chlorim 25WDG may be made any time after the first trifoliolate has opened but no later than 60 days before soybean maturity.
- **Timing to Weeds:** treatment should be made when weeds are young and actively growing (after the first true leaves have expanded, but before the weeds are larger than the size listed below). Treatment made to weeds larger than the sizes indicated below, or to weeds under stress may result in unsatisfactory control.

FIELD NOTES
