

WILLOWOOD SULFEN 4SC

Same active as SPARTAN® and AUTHORITY®



OVERVIEW

Willowood Sulfen 4SC is a preemergence herbicide for the control of broadleaf weeds, grasses, and sedges.

KEY CROPS

Corn • Soybeans • Citrus • Wheat • Sorghum

PACKAGING

- 2.5 gal jug (x2)
- 265 gal tote (x1)

PRODUCT SPECIFICATIONS

ACTIVE INGREDIENT

Sulfentrazone (4.0 lbs Active Ingredient per Gallon)

RATES

4.5 fl oz/A - 12.0 fl oz/A
(See label for rates by crop)

WEED SPECTRUM

Kochia • Marestalk • Dandelions • Sedge

APPLICATION TIMING

Preemergence

Technical Use Guidelines

Mode of Action	<ul style="list-style-type: none"> • PPO INHIBITORS
Crop Rotation REFER TO LABEL FOR DETAILS	<ul style="list-style-type: none"> • Barley, Wheat: 4 months • Corn: 10 months • Cotton: 18 months • Canola: 24 months
Adjuvant Recommendation	<ul style="list-style-type: none"> • COC: 1% v/v + AMS for burndown
Rainfastness	<ul style="list-style-type: none"> • 1 hour
Water Volume (GPA) REFER TO LABEL FOR DETAILS	<ul style="list-style-type: none"> • Ground: 10 GPA (Minimum) • Aerial: 5 GPA (Minimum)
Application Tips	<ul style="list-style-type: none"> • Use 15 GPA for coverage during burndown application. • Great Tank mix partner with Dual for preemergence weed control. • Do not spray if the Soybeans are emerged or cracking.

ALWAYS READ AND FOLLOW LABEL DIRECTIONS. It is a violation of federal and state/provincial law to use any pesticide product other than in accordance with its label. The distribution, sale and use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. We do not guarantee the accuracy of any information provided on this page or which is provided by us in any form. It is your responsibility to confirm prior to purchase and use that a product is labeled for your specific purposes, including, but not limited to, your target crop or pest and its compatibility with other products in a tank mix. This information should not be used as a replacement for consulting the applicable product label. Consult the label for the most complete and up-to-date information about any referenced product. Readers must have a valid applicator license to use restricted use pesticides. Please consult your state department of agriculture for complete rules and regulations on the use of restricted use pesticides, as some products require specific record-keeping requirements.

WILLOWOOD SULFEN 4SC

Same active as SPARTAN® and AUTHORITY®

MIXING PROCEDURE

Willowood Sulfen 4SC may be applied on its own or in combination with other herbicides for a broader spectrum of weed control.

Mix Willowood Sulfen 4SC as follows:

- Fill a clean spray tank ½ full of water required for treatment.
- Begin agitation.
- Use a clean container to create a slurry of Willowood Sulfen 4SC and water.
- Add the slurry slowly to the spray tank, continuing agitation throughout.
- Rinse the slurry mix container and add rinsate solution to spray tank.
- Finish filling spray tank to required level.
- Maintain agitation throughout. The Willowood Sulfen 4SC/water slurry must be mixed thoroughly prior to application.

CROP STAGING

- **Corn (Field, Seed, Popcorn):**
 - Apply pre-plant to stubble or soil surface in the fall before spring planting .
 - Apply early pre-plant, pre-plant incorporated, pre-emergence in the spring 45 days prior to planting or up to 3 days after planting if seed furrow is closed and seedlings have not broken soil surface.
- **Soybeans:**
 - Apply pre-emergence or pre-plant incorporated in the spring, or in the fall before planting.
 - Apply to the soil surface in the spring either pre-plant incorporated or pre-emergence up to 3 days after planting.
- **Citrus:**
 - Make a broadcast soil application when weeds are not present and/or a post-emergence herbicide has been tank mixed with Willowood Sulfen 4SC.
 - Make applications to crops that have been established for one full growing season and are healthy and vigorous.
 - Apply when soil is moist and application will be followed by 1/2" rainfall or irrigation within two weeks after application.
 - Time applications to take advantage of normal rainfall and cool temperatures, especially if using drip or micro sprinkler irrigation is used which may not provide uniform incorporation into the soil.
- **Wheat (Spring):**
 - Make on-pre-plant or pre-emergence application 40-60 days before forage cutting and 120 days before grain harvest.

FIELD NOTES