

BENZ

Contains bentazon and imazamox, the active ingredients in Viper® ADV



OVERVIEW

Broad-spectrum control of most annual grasses and broadleaf weeds in field and succulent peas, dry faba beans, soybeans, dry beans, established red and alsike clover for seed production, and seedling and established alfalfa for seed, forage and hay.

KEY CROPS

Soybeans • Field peas • Dry edible beans

Packaging

- 8.1 L Jug (x2)
- 129.6 L Drum

For sale for use in Manitoba, Saskatchewan, Alberta, Peace River Region and Interior of British Columbia Only.

PRODUCT SPECIFICATIONS

ACTIVE INGREDIENT

Bentazon and Imazamox

RATES

0.4 L per Acre
(Refer to label for rates by crop)

WEED SPECTRUM

Broad-spectrum weed control of broadleaf weeds & grasses

Technical Use Guidelines

Mode of Action	<ul style="list-style-type: none"> • Group 2, 6
Crop Rotation	<ul style="list-style-type: none"> • Winter wheat: 3 months (3 MAT) • <i>Avoid spray overlap as yield reduction may result.</i> • <i>Initial crop injury to non-Clearfield Canola may be observed.</i>
Adjuvants	<ul style="list-style-type: none"> • 0.8 L/Acre of UAN is required with Benz
Rainfastness	<ul style="list-style-type: none"> • Rainfast in 6 hours
Water Volume	<ul style="list-style-type: none"> • Minimum 10 US gal/Ac
Application Tips	<ul style="list-style-type: none"> • The following crops may be grown safely the year following an application: • <i>Canary seed, Field corn, Field peas, Soybeans, Clearfield® Canola</i> • <i>(e.g., canola varieties with the Clearfield® trait), Non-Clearfield canola, Lentils, Spring wheat, Durum wheat, Spring barley, Sunflower, Tame oats, Flax, Chickpeas, Clearfield® Sunflowers (e.g., sunflower varieties with the Clearfield® trait)</i>

BENZ

Contains benzaton and imazamox, the active ingredients in Viper® ADV



MIXING PROCEDURE

Tank Mix Partners:

1) 0.4 L/Ac of Benz + (360-720 gm AI/Ac of glyphosate) in RR soybeans

2) 0.4 L/Ac of Benz + 0.144 L /Ac of Basagran Forte + 0.8 L/Ac UAN 28% on Dry Edible beans

CROP STAGING

- Application should be made from the 3-6 node stage of field peas and dry faba beans and the cotyledon to 4-leaf stage of soybeans and after weeds have emerged
- Apply when broadleaf weeds are from the cotyledon to 4-leaf stage and when grassy weeds are at the 1-4 true leaf or early tillering
- For field peas and dry faba beans, initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield
- DO NOT make more than one application per season

FIELD NOTES
