

WILLOWOOD IMAZETHAPYR 2SL

Same active as PURSUIT®



OVERVIEW

Willowood Imazethapyr 2SL is a contact preplant, preplant incorporated, pre emergence, and post emergence herbicide for the control of certain broadleaf weeds and yellow nutsedge.

KEY CROPS

Soybeans • Alfalfa • Peas • Edible Beans

PACKAGING

- 1 gal jug (x4)

PRODUCT SPECIFICATIONS

ACTIVE INGREDIENT

Imazethapyr (2.0 lbs Active Ingredient per Gallon)

RATES

3.0 fl oz/A - 6.0 fl oz/A
(See label for rates by crop)

WEED SPECTRUM

Broadleaf Weeds • Yellow Nutsedge • Grasses

APPLICATION TIMING

Pre Emergence • Post Emergence

Technical Use Guidelines

Mode of Action	<ul style="list-style-type: none"> • ALS INHIBITORS
Crop Rotation REFER TO LABEL FOR DETAILS	<ul style="list-style-type: none"> • Corn & Barley: Next season • Alfalfa, Clover, Wheat: 4 months • Cotton, Oats, Sorghum: 18 months
Adjuvant Recommendation	<ul style="list-style-type: none"> • AMS • COC: 1% V/V • MSO: 1% V/V • NIS: .25% V/V
Rainfastness	<ul style="list-style-type: none"> • 1 hour
Water Volume (GPA) REFER TO LABEL FOR DETAILS	<ul style="list-style-type: none"> • Ground: 10 GPA (minimum) • Aerial: 5 GPA (minimum)
Application Tips	<ul style="list-style-type: none"> • For optimum weed control, cultivate the field 7 – 10 days after a post emergence Willowood Imazethapyr 2SL application. • Timely cultivation will improve residual weed control, particularly under dry conditions. • For consideration, add metolachlor as a pre and fomesafen as a post.

ALWAYS READ AND FOLLOW LABEL DIRECTIONS. It is a violation of federal and state/provincial law to use any pesticide product other than in accordance with its label. The distribution, sale and use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. We do not guarantee the accuracy of any information provided on this page or which is provided by us in any form. It is your responsibility to confirm prior to purchase and use that a product is labeled for your specific purposes, including, but not limited to, your target crop or pest and its compatibility with other products in a tank mix. This information should not be used as a replacement for consulting the applicable product label. Consult the label for the most complete and up-to-date information about any referenced product. Readers must have a valid applicator license to use restricted use pesticides. Please consult your state department of agriculture for complete rules and regulations on the use of restricted use pesticides, as some products require specific record-keeping requirements.

WILLOWOOD IMAZETHAPYR 2SL

Same active as PURSUIT®

MIXING PROCEDURE

Mix Willowood Imazethapyr 2SL as follows:

- Fill spray tank 1/2 full with clean water.
- Add water soluble packet products. Mix thoroughly.
- Add WP (wettable powder), DG (dispersible granule), DF (dry flowable), or liquid flowable formulations (not in soluble packets).
- Add Willowood Imazethapyr 2SL. Mix thoroughly.
- Add other water soluble products.
- Add EC (emulsifiable concentrate) products.
- Add crop oil concentrates or surfactants.
- Add liquid fertilizer.
- Fill the remainder of the tank with water while maintaining agitation.

CROP STAGING

- **Alfalfa**
 - Seedling: make application when the seedling alfalfa is in the second (2nd) trifoliate stage or larger and when the majority of the weeds are 1 to 3 inches in height.
 - Established Stands: make application to established alfalfa in the fall; to dormant, or semi-dormant alfalfa in the spring (less than 3 inches of re-growth); or between cuttings. Applications must be made before significant alfalfa (less than or equal to 3 inches) to allow the product to reach the target weeds.
- **Soybeans**
 - Application must be made before bloom.
- **Edible Beans and Peas**
 - See label for specific application instructions.
- **Peas**
 - See label for specific application instructions.

FIELD NOTES
