

PREACH FUNGICIDE

Contains *Pyraclostrobin*, the active ingredient in *Headline® EC Fungicide*



OVERVIEW

Helps prevent crops from disease by inhibiting spore germination and offers a curative-eradivative effect by inhibiting the growth and sporulation of the fungus on the leaf surface.

KEY CROPS

Lentils | Field peas | Canola | Dry beans
Corn | Cereals | Flax | Potatoes

PACKAGING

6.5 L jets (x2)

PRODUCT SPECIFICATIONS

ACTIVE INGREDIENT

Pyraclostrobin

RATE RANGE

- 0.12-0.36 L/Acre

DISEASE SPECTRUM

Leaf rust • Tan spot • Septoria leaf spot
• Spot blotch • Stripe rust • Scald Rye • Leaf rust
• Powdery mildew • Crown rust
• Ascochyta blight • Black spot • Blackleg
• Anthracnose

APPLICATION TIMING

Apply Preach prior to disease development or when weather conditions are favourable for disease development

Technical Use Guidelines

Mode of Action	• Group 11
Method of Application	• Ground, Air
REI	• 12 hours
Rainfastness	• Avoid application when heavy rain is forecast
Water Volume	• Use a minimum of 10 US gal/Ac
Pre-Harvest Interval	• Ranges between 7-30 days prior to harvest depending on the crop