

WILLOWOOD THIO ULTRAMAX

Same active as BOLERO® ULTRAMAX



OVERVIEW

Willowood Thio UltraMax is a dry thiobencarb formulation that provides outstanding contact and residual weed control in rice similar to Willowood Thioben 8EC, but in a convenient non-VOC containing dry formulation.

KEY CROPS

Rice

REGISTERED STATE

- California only

PRODUCT SPECIFICATIONS

ACTIVE INGREDIENT

Thiobencarb (0.15lb Active Ingredient per Pound)

RATES

23.3 lbs/A - 26.7 lbs/A
(See label for rates by crop)

WEED SPECTRUM

Barnyardgrass • Smallflower Umbrellaplant • Sprangletop

APPLICATION TIMING

Apply after flooding when rice is in the expanded, 2-leaf stage

Technical Use Guidelines

Mode of Action	<ul style="list-style-type: none"> • LIPID SYNTHESIS INHIBITOR (NOT ACCase)
Crop Rotation <small>REFER TO LABEL FOR DETAILS</small>	<ul style="list-style-type: none"> • 6 months on all rotational crops
Adjuvant Recommendation	<ul style="list-style-type: none"> • Not Applicable
Rainfastness	<ul style="list-style-type: none"> • Not Applicable
Water Volume (GPA) <small>REFER TO LABEL FOR DETAILS</small>	<ul style="list-style-type: none"> • Ground: 10 GPA - 20 GPA • Aerial: 10 GPA
Application Tips	<ul style="list-style-type: none"> • Do not apply if rice exhibits symptoms of stress. • The variety Cal Pearl may react adversely to some stress conditions more than other varieties. • Do not drain fields that have been treated with Thio UltraMax for a minimum of 30 days following treatment. • Do not make more than one application per year.

ALWAYS READ AND FOLLOW LABEL DIRECTIONS. It is a violation of federal and state/provincial law to use any pesticide product other than in accordance with its label. The distribution, sale and use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. We do not guarantee the accuracy of any information provided on this page or which is provided by us in any form. It is your responsibility to confirm prior to purchase and use that a product is labeled for your specific purposes, including, but not limited to, your target crop or pest and its compatibility with other products in a tank mix. This information should not be used as a replacement for consulting the applicable product label. Consult the label for the most complete and up-to-date information about any referenced product. Readers must have a valid applicator license to use restricted use pesticides. Please consult your state department of agriculture for complete rules and regulations on the use of restricted use pesticides, as some products require specific record-keeping requirements.

GROUP

8

HERBICIDE

WILLOWOOD THIO ULTRAMAX

Same active as BOLERO® ULTRAMAX

CROP STAGING

- Apply after rice has been flooded
 - Rice is in the expanded, 2-leaf stage (first open leaf)
 - Sprangletop population is 2 leaf or less
 - Smallflower umbrellaplant is 3 leaf or less

FIELD NOTES
