

WILLOWOOD THIONIL EC

Same active as RICEBEAUX®



OVERVIEW

Willowood Thionil EC is an excellent herbicide for the control of broadleaf and grass weed species in rice. An EC formulation with multiple modes of action provides both contact and residual weed control on a broad spectrum of weeds. An excellent weed control solution in rice with both Newpath® herbicide in the Clearfield® production system as well as the conventional rice acres.

KEY CROPS

Rice

AVAILABLE PACK SIZE

- 30 gal drum (x1)
- 265 gal tote (x1)

PRODUCT SPECIFICATIONS

ACTIVE INGREDIENT

- Thiobencarb (3.0 lbs Active Ingredient per Gallon)
- Propanil (3.0 lbs Active Ingredient per Gallon)

RATES

2.0 qt/A - 5.3 qt/A
(See label for rates by crop)

WEED SPECTRUM

Signalgrass • Sprangletop Broadleaf • Hemp Sesbania • Dayflower • Pitted Morningglory

APPLICATION TIMING

Pre Emergence • Post Emergence

Technical Use Guidelines

Mode of Action	<ul style="list-style-type: none"> • LIPID SYNTHESIS INHIBITOR (NOT ACCase)
Crop Rotation <small>REFER TO LABEL FOR DETAILS</small>	<ul style="list-style-type: none"> • 6 months on all rotational crops
Adjuvant Recommendation	<ul style="list-style-type: none"> • NIS @ 0.25% v/v
Rainfastness	<ul style="list-style-type: none"> • 24 hours
Water Volume (GPA) <small>REFER TO LABEL FOR DETAILS</small>	<ul style="list-style-type: none"> • Ground: 20GPA - 30GPA • Aerial: 10GPA
Application Tips	<ul style="list-style-type: none"> • This product normally requires 8 hours of DIRECT sunlight after application for absorption into target weeds. • Recommend application of product be planned so that the applied product remains in contact with the leaf surfaces for at least 48 hours prior to rainfall or flooding. • Morning applications have produced better results in weed control.

ALWAYS READ AND FOLLOW LABEL DIRECTIONS. It is a violation of federal and state/provincial law to use any pesticide product other than in accordance with its label. The distribution, sale and use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. We do not guarantee the accuracy of any information provided on this page or which is provided by us in any form. It is your responsibility to confirm prior to purchase and use that a product is labeled for your specific purposes, including, but not limited to, your target crop or pest and its compatibility with other products in a tank mix. This information should not be used as a replacement for consulting the applicable product label. Consult the label for the most complete and up-to-date information about any referenced product. Readers must have a valid applicator license to use restricted use pesticides. Please consult your state department of agriculture for complete rules and regulations on the use of restricted use pesticides, as some products require specific record-keeping requirements.

WILLOWOOD THONIL EC

Same active as RICEBEAUX®

MIXING PROCEDURE

Mix Willowood Thionil EC as follows:

- Fill the spray tank $\frac{1}{4}$ to $\frac{1}{3}$ with clean water as the carrier.
- Begin agitation and add the specified amount of Willowood Thionil EC.
- Maintain agitation until the product is fully dispersed for at least 5 minutes.
- While maintaining agitation, fill the tank with water. The product should be thoroughly mixed prior to adding any other material.
- While the tank is filling, add any tank mix partner (other labeled rice herbicides, adjuvants, drift control agents, etc.).
- Settling may occur if the constant agitation is not maintained. If settling does occur, thoroughly mix before using.
- Make application within 24 hours of product mixing, or the product may degrade.
- If product and tank mix partner will be applied in multiple loads, pre-slurry the product in clean water before adding to the tank. This will prevent the tank mix partner from affecting the dissolution of the Willowood Thionil EC.

CROP STAGING

- Best results for selective weed control are obtained when most grasses have reached the 1 to 3-leaf stage

FIELD NOTES
