FBN[®] PROTEB 250 EC

Contains prothioconazole and tebuconazole, the active ingredients in Prosaro® Fungicide



OVERVIEW

Improve Crop Grades by reducing DON (deoxynivalenol) levels, and reducing FDK (fusarium damaged kernels), while also improving Bushel Weight. Acts as both a protective and a curative fungicide.

BENEFITS

Among the most effective Fusarium Head Blight fungicides available, *FBN* PROTEB offers both leaf and head disease control, helping improve yield, and quality.

Key Crops

Spring Wheat | Winter Wheat | Durum Wheat Barley | Oats

PACKAGING

- 2 x 6.5L case (40 Acres)
- 104 L drum (320 Acres)

PRODUCT SPECIFICATIONS

ACTIVE INGREDIENT

Prothioconazole (125 g/L) and Tebuconazole (125 g/L)

APPLICATION RATE

324 mL/AC

DISEASE SPECTRUM

Rusts | Powdery Mildew | Blotches | Scald | Septoria | Tan Spot | Fusarium Head Blight | Powdery Mildew | Blotches | Septoria | Tan Spot | Fusarium Head Blight

APPLICATION TIMING

For suppression of Fusarium Head Blight in wheat, apply when at least 75% of the heads on the main stem are fully emerged, to when to when 50% of the heads on the main stem are in full flower. Application at this time will also control the listed leaf diseases.

For suppression of Fusarium Head Blight in barley, apply when 70 - 100 % of the heads on the main stem are fully emerged, to 3 days after full head emergence.

In Oats apply as a preventative spray when the earliest leaf disease symptoms appear on the leafs and stems.

© 2015 - 2023 Farmer's Business Network Canada, Inc. All rights reserved. The sprout logo, "Farmers Business Network," and "FBN," are trademarks or registered trademarks of Farmer's Business Network, Inc. or its affiliates. Prosaro® is a registered trademarks of Bayer Intellectual Property GmbH or its affiliates. All other trademarks are the property of their respective owners. FBN Direct products and services and other products distributed by FBN Direct are offered by Farmer's Business Network Canada, Inc. all canada, Inc. all





FBN[®] PROTEB 250 EC

Contains prothioconazole and tebuconazole, the active ingredients in Prosaro® Fungicide



Technical Use Guidelines		
Mode of Action	• Group 3	
Method of Application	Ground, Air	
REI	• 12 hours	
PHI	• 36 days	
Rainfastness	Avoid application when heavy rain is forecast	
Water Volume	Ground: 10.7 US GAL/AC Air: 5.5 US GAL/AC	
Minimum Treatment Interval	• 7 days	

PRODUCT SPECIFICATIONS

APPLICATION TIMING

- FBN PROTEB 250 Fungicide must be applied with properly calibrated, clean equipment.
- Prior to adding FBN PROTEB 250 Fungicide to the spray tank, ensure that the spray tank is thoroughly clean.
- Add one-half of the required amount of water to the spray or mixing tank and start agitation.
- Add the required quantity of FBN PROTEB 250
 Fungicide to the water and complete filling
 with water to the required total volume.
- Maintain agitation throughout mixing and spraying.

FIELD NOTES

© 2015 - 2023 Farmer's Business Network Canada, Inc. All rights reserved. The sprout logo, "Farmers Business Network," and "FBN," are trademarks or registered trademarks of Farmer's Business Network, Inc. or its affiliates. Prosaro® is a registered trademarks of Bayer Intellectual Property GmbH or its affiliates. All other trademarks are the property of their respective owners. FBN Direct products and services and other products distributed by FBN Direct are offered by Farmer's Business Network Canada, Inc. all canada, Inc. all

