

GROUP

1

HERBICIDE

WILLOWOOD CLETHODIM 2EC

Same active as *SELECT®*



OVERVIEW

Willowood Clethodim 2EC is a post emergence herbicide that controls a wide range of the toughest annual and perennial grasses. Clethodim improves the profitability of glyphosate tolerant corn-soybean rotations by controlling volunteer glyphosate-tolerant corn and flushes in glyphosate-tolerant soybeans.

KEY CROPS

Alfalfa • Citrus • Grapes • Tree Fruit & Nuts • Soybeans • Sugar Beets • Vegetables •

PACKAGING

- 265 gal tote (x1)
- 2.5 gal jug (x2)

PRODUCT SPECIFICATIONS

ACTIVE INGREDIENT

Clethodim (2.00 lbs Active Ingredient per gallon)

RATES

3.0 fl oz/A - 16.0 fl oz/A
(See label for rates by crop)

WEED SPECTRUM

Volunteer Corn • Annual Grasses • Perennial Grasses

APPLICATION TIMING

Post Emergence

Technical Use Guidelines

Mode of Action	<ul style="list-style-type: none"> • ACCase Inhibitors
Crop Rotation REFER TO LABEL FOR DETAILS	<ul style="list-style-type: none"> • Field Corn: 6 days • Wheat: 1 day • Canola: 1 day • Rye: 1 day
Adjuvant Recommendation	<ul style="list-style-type: none"> • AMS: 2.5-4.0 lb/Ac • NIS: 0.125% to 0.25% V/V
Rainfastness	<ul style="list-style-type: none"> • 1 hour
Water Volume (GPA) REFER TO LABEL FOR DETAILS	<ul style="list-style-type: none"> • Ground: 5 GPA - 40 GPA • Air: 3 GPA - 20 GPA
Application Tips	<ul style="list-style-type: none"> • Tank mixes easily with other broadleaf herbicides, glyphosate, and several insecticides. • For volunteer control, use COC. (1 gal/100 acres) + AMS (2.5lb - 4lb/100 acres). • Typical rate of 6 fl oz/A - 10 fl oz/A depending on Volunteer Corn height. • Will not control broadleaf weeds.

ALWAYS READ AND FOLLOW LABEL DIRECTIONS. It is a violation of federal and state/provincial law to use any pesticide product other than in accordance with its label. The distribution, sale and use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. We do not guarantee the accuracy of any information provided on this page or which is provided by us in any form. It is your responsibility to confirm prior to purchase and use that a product is labeled for your specific purposes, including, but not limited to, your target crop or pest and its compatibility with other products in a tank mix. This information should not be used as a replacement for consulting the applicable product label. Consult the label for the most complete and up-to-date information about any referenced product. Readers must have a valid applicator license to use restricted use pesticides. Please consult your state department of agriculture for complete rules and regulations on the use of restricted use pesticides, as some products require specific record-keeping requirements.



WILLOWOOD CLETHODIM 2EC

Same active as *SELECT*®

MIXING PROCEDURE

Mix Willowood Chethodim 2EC as follows:

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. While agitating, add the correct amount of Willowood Clethodim 2EC. Agitation should create a rippling or rolling action on the water surface.
3. If tank mixing Willowood Clethodim 2EC with other labeled herbicides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates, and then solutions. Prepare no more spray mixture than is required for the immediate spray operation.
4. Add any required adjuvants (crop oil concentrate, non-ionic surfactant and/or nitrogen solution).
5. Fill spray tank to desired level with water. Agitation should continue until all spray solution has been applied.

CROP STAGING

- Apply post emergence to actively growing grasses according to label. Applications made to grass plants stressed by insufficient moisture, or cold temperatures, or to grass plants exceeding specified growth stages may result in unsatisfactory control.
- In arid regions where irrigation is used to supplement limited rainfall, apply Clethodim 2EC as soon as possible, after an irrigation (within 7 days). In arid regions, a second application of will generally provide more effective control of perennial grass weeds than a single application. Make a second application to actively growing grasses 2-to-3 weeks after emergence of new growth.
- Cultivation of treated grasses 7 days prior to, or within 7 days after, application may reduce weed control. DO NOT APPLY Clethodim 2EC if rainfall is expected within one hour, since control may be reduced.

FIELD NOTES
