

WILLOWOOD PROPANIL 4SC

Same active as SUPERWHAM!®



OVERVIEW

Willowood Propanil 4SC is for post emergence weed control in rice. Willowood Propanil 4SC is most effective if applied when susceptible grasses and broadleaf weeds are small and growing actively under favorable soil moisture and weather conditions. Early weed control removes weed competition from the rice crop, saves moisture, and generally contributes to increased yields.

KEY CROPS

Rice

PACKAGING

- 30 gal tote (x1)

PRODUCT SPECIFICATIONS

ACTIVE INGREDIENT

Propanil (4.0 lbs Active Ingredient per Gallon)

RATES

3.0 qt/A - 4.0 qt/A

(See label for rates by crop)

WEED SPECTRUM

Annual Sedge • Barnyardgrass • Broadleaf Signalgrass • Crabgrass Species • Foxtail Species • Goosegrass • Coffebean • Mexicanweed • Redroot Pigweed • Smallflower Umbrella Plant • Texas Panicum • Watergrass

APPLICATION TIMING

Post Emergence

Technical Use Guidelines

Mode of Action	<ul style="list-style-type: none"> • PHOTOSYSTEM II INHIBITORS
Crop Rotation <small>REFER TO LABEL FOR DETAILS</small>	<ul style="list-style-type: none"> • 60 days on all rotational crops
Adjuvant Recommendation	<ul style="list-style-type: none"> • COC: 8.0 fl oz/A - 16.0 fl oz/A
Rainfastness	<ul style="list-style-type: none"> • 8 hours (48 hrs highly recommended)
Water Volume (GPA) <small>REFER TO LABEL FOR DETAILS</small>	<ul style="list-style-type: none"> • Ground: 15 GPA - 20 GPA • Aerial: 10 GPA - 15 GPA
Application Tips	<ul style="list-style-type: none"> • Thorough coverage of emerged weeds is essential for best results. • Willowood Propanil 4SC has no preemergence or residual herbicidal activity. • Apply Willowood Propanil 4SC only to fields that have been drained of floodwater.

ALWAYS READ AND FOLLOW LABEL DIRECTIONS. It is a violation of federal and state/provincial law to use any pesticide product other than in accordance with its label. The distribution, sale and use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. We do not guarantee the accuracy of any information provided on this page or which is provided by us in any form. It is your responsibility to confirm prior to purchase and use that a product is labeled for your specific purposes, including, but not limited to, your target crop or pest and its compatibility with other products in a tank mix. This information should not be used as a replacement for consulting the applicable product label. Consult the label for the most complete and up-to-date information about any referenced product. Readers must have a valid applicator license to use restricted use pesticides. Please consult your state department of agriculture for complete rules and regulations on the use of restricted use pesticides, as some products require specific record-keeping requirements.

WILLOWOOD PROPANIL 4SC

Same active as SUPERWHAM!®

MIXING PROCEDURE

Mix Willowood Propanil 4SC (wet spray) as follows:

- Fill the tank 1/4 to 1/3 full of clean water.
- While agitating, add the required amount of Propanil 4SC
- Continue agitation until the Propanil 4SC is fully dispersed, at least 5 minutes.
- Once the Propanil 4SC is fully dispersed, maintain agitation and continue filling the tank with water. The Propanil 4SC should be thoroughly mixed with water before adding any other material.
- As the tank is filling, add the required tank mix partner (other labeled rice herbicides, adjuvants, drift control agents, etc.).
- If the mixture is not continuously agitated, settling may occur. If settling occurs, thoroughly re-agitate before using.
- Apply Propanil 4SC spray preparations within 24 hours of product mixing, or the product may degrade.
- If Propanil 4SC and a tank mix partner are to be applied in multiple loads, pre-slurry the Propanil 4SC in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the Propanil 4SC

CROP STAGING

- **Early Timing:** apply when a satisfactory stand of rice has been established that will tolerate flooding (1 to early 4 leaf staging).
- **Mid-Timing:** apply to actively growing grasses in the 4-to-6 leaf and early tilling stage, or when they are in the 2-to-4 leaf stage but stressed under dry soil conditions.
- **Rescue Timing:** control of older tillering grass, usually 30-40 days after planting.

FIELD NOTES
